Supplementary Material for *Interaction of Light with Arbitrarily Shaped Dielectric Media: Spectral Analysis & Evolution Semigroups of Electromagnetic Scattering*

Yajun Zhou

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138

(Dated: November 24, 2009)

arXiv:0911.4540v1 [math-ph] 24 Nov 2009

Contents

S0. Background and Outline 1

S1. Spectral Properties of Electromagnetic Scattering 4
- S1.0. Shape Regularity, Energy Admissibility, and the Born Equation 5
    - S1.0.1. Regular Shapes and Energetically Admissible Vector Fields 5
    - S1.0.2. Born Equation for Energetically Admissible Electric Fields 7
- S1.1. Uniqueness Theorem for Electromagnetic Scattering 9
    - S1.1.1. Fourier Transform of the Born Operator 9
    - S1.1.2. A Generalized Optical Theorem 11
    - S1.1.3. The Uniqueness Theorem for Light Scattering 14
- S1.2. Existence Theorem for Electromagnetic Scattering 16
    - S1.2.1. Fredholm Decomposition of the Born operator 16
    - S1.2.2. The Existence Theorem for Light Scattering 18
    - S1.2.3. Stable Dependence of the Solution on Experimental Parameters 22
- S1.3. Additional Comments and Details 24
    - S1.3.1. Born Equation in the Discrete Dipole Approximation Method for Light Scattering 24
    - S1.3.2. Rigorous Derivations of Some Fourier Inversion Formulae 26
    - S1.3.3. Norm Estimates of the Born Operator 28
    - S1.3.4. Upper Bounds for the Total Scattering Cross-Section 32
    - S1.3.5. Analytic Continuation of the Helmholtz Equation 36
    - S1.3.6. Generalizations to Inhomogeneous and Anisotropic Media 38

S2. Evolution Semigroups for Electromagnetic Scattering 39
- S2.1. Integral Representation of the Solution to Light Scattering 40
    - S2.1.1. Evolution Semigroups, Mean Ergodicity, and Fixed Points 40
    - S2.1.2. Evolution of Physically Admissible Vector Fields — Extended Uniqueness Theorem 42
- S2.2. Smooth Solution, Optical Resonance and Strong Stability — Extended Existence Theorem 46
- S2.3. Additional Comments and Details 54
    - S2.3.1. Tubular Neighborhoods of a Smooth Surface 54
    - S2.3.2. Connectedness in the Perspectives of de Rham and Hodge 59
    - S2.3.3. Transverse Vector Fields in Star-Shaped Dielectrics 63
    - S2.3.4. Surface Geometry and Optical Resonance 64
    - S2.3.5. Optical Resonance in Mie Scattering 72
    - S2.3.6. Resonance Modes in Acoustic and Electromagnetic Scattering 79

References 83

S0. BACKGROUND AND OUTLINE

Following the line of thoughts established in Rayleigh-Gans scattering [1, 2, 3], Max Born (see Ref. [4], Kap. VII, “Molekulare Optik”) attacked the light scattering problem in general geometries by viewing the process as the radiation of induced dipoles in the dielectric medium (i.e. with dielectric constant $\epsilon = (1+\chi)\epsilon_0 = n^2\epsilon_0$, but otherwise non-magnetic, with magnetic permeability $\mu = \mu_0$ equal to that of the vacuum). Born’s approach formulates the light scattering problem as an integral equation, akin to the Lippmann-Schwinger equation in quantum scattering.

Adopting the convention to separate the time and space variables of monochromatic light radiation as $\boldsymbol{E}(\boldsymbol{r},t) = \boldsymbol{E}(\boldsymbol{r})e^{i\omega t}$,[1] writing the electric field of the incident light as $\boldsymbol{E}_{\text{inc}}(\boldsymbol{r},t) = \boldsymbol{E}_{\text{inc}}(\boldsymbol{r})e^{i\omega t}$, and using $\chi(\boldsymbol{r}) = n^2(\boldsymbol{r}) - 1$ to denote spatial distribution of dielectric susceptibility, we may put the Born equation (cf. Ref. [4], p. 315, Eq. 11) of light scattering in the form of

$$\boldsymbol{E}(\boldsymbol{r}) = \boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) + \nabla\times\nabla\times\iiint \frac{\chi(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' - \chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r}). \tag{1}$$

The three terms on the right-hand side represent the incident field, the radiation field due to oscillating dipoles, and the depolarization field, respectively. It is usually a physical requirement that the incident field be a source-free vibration in the entire space, i.e. $\nabla\cdot\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) = 0, (\nabla^2 + k^2)\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r}\in\mathbb{R}^3$. This is a natural consequence of the homogeneous Maxwell equations [9] satisfied by the incident electric field $\boldsymbol{E}_{\text{inc}}(\boldsymbol{r},t)$ and magnetic (induction) field $\boldsymbol{B}_{\text{inc}}(\boldsymbol{r},t)$ in a space devoid of charges and currents:[2]

$$\nabla\cdot\boldsymbol{E}_{\text{inc}} = 0,\quad \nabla\times\boldsymbol{B}_{\text{inc}} = \mu_0\epsilon_0\frac{\partial\boldsymbol{E}_{\text{inc}}}{\partial t},\quad \nabla\cdot\boldsymbol{B}_{\text{inc}} = 0,\quad \nabla\times\boldsymbol{E}_{\text{inc}} = -\frac{\partial\boldsymbol{B}_{\text{inc}}}{\partial t}.$$

Here, the relation $(\mu_0\epsilon_0)^{-1/2} = \omega/k = c$ gives the speed of light in the vacuum.

Following the spirit in Born and Wolf’s book (Ref. [10], see also [11], pp. 33-34), one may derive the Born equation from the Maxwell equations as follows. Let $\boldsymbol{D}(\boldsymbol{r},t) = [1+\chi(\boldsymbol{r})]\epsilon_0\boldsymbol{E}(\boldsymbol{r},t)$ be the electric displacement, $\boldsymbol{A}(\boldsymbol{r},t)$ be the vector potential satisfying $\boldsymbol{B}(\boldsymbol{r},t) = \nabla\times\boldsymbol{A}(\boldsymbol{r},t)$, then we have

$$i\omega[\boldsymbol{E}(\boldsymbol{r}) + \chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})]e^{i\omega t} = \frac{1}{\epsilon_0}\frac{\partial\boldsymbol{D}(\boldsymbol{r},t)}{\partial t} = c^2\nabla\times\boldsymbol{B}(\boldsymbol{r},t) = c^2\nabla\times\nabla\times\boldsymbol{A}(\boldsymbol{r},t),$$

according to one of the Maxwell equations. Here, the total electric field $\boldsymbol{E}(\boldsymbol{r},t) = \boldsymbol{E}_{\text{inc}}(\boldsymbol{r},t) + \boldsymbol{E}_{\text{sc}}(\boldsymbol{r},t)$ and total magnetic field $\boldsymbol{B}(\boldsymbol{r},t) = \boldsymbol{B}_{\text{inc}}(\boldsymbol{r},t) + \boldsymbol{B}_{\text{sc}}(\boldsymbol{r},t)$ can be attributed to the superposition of the incident field $\boldsymbol{E}_{\text{inc}}(\boldsymbol{r},t), \boldsymbol{B}_{\text{inc}}(\boldsymbol{r},t)$ and the scattering field $\boldsymbol{E}_{\text{sc}}(\boldsymbol{r},t), \boldsymbol{B}_{\text{sc}}(\boldsymbol{r},t)$. The scattering electromagnetic field, with $O(1/|\boldsymbol{r}|)$ asymptotic behavior at large distances $|\boldsymbol{r}|\to+\infty$, satisfies the inhomogeneous Maxwell equations

$$\nabla\cdot\boldsymbol{E}_{\text{sc}} = 0,\quad \nabla\times\boldsymbol{B}_{\text{sc}} = \mu_0\left(\boldsymbol{J}_P + \epsilon_0\frac{\partial\boldsymbol{E}_{\text{sc}}}{\partial t}\right),\quad \nabla\cdot\boldsymbol{B}_{\text{sc}} = 0,\quad \nabla\times\boldsymbol{E}_{\text{sc}} = -\frac{\partial\boldsymbol{B}_{\text{sc}}}{\partial t}.$$

Here, the polarization current density $\boldsymbol{J}_P(\boldsymbol{r},t) = [\epsilon(\boldsymbol{r}) - \epsilon_0]\partial\boldsymbol{E}(\boldsymbol{r},t)/\partial t = i\omega\epsilon_0\chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})e^{i\omega t}$ (the density of electric current generated by oscillating dipoles in the dielectric medium) acts as a source for electromagnetic vibration, giving rise to the d’Alembert equation for $\boldsymbol{A}_{\text{sc}}(\boldsymbol{r},t)$ as

$$\nabla^2\boldsymbol{A}_{\text{sc}}(\boldsymbol{r},t) - \frac{1}{c^2}\frac{\partial^2\boldsymbol{A}_{\text{sc}}(\boldsymbol{r},t)}{\partial t^2} = -\mu_0\boldsymbol{J}_P(\boldsymbol{r},t) = -\frac{\chi(\boldsymbol{r})}{c^2}\frac{\partial\boldsymbol{E}(\boldsymbol{r},t)}{\partial t}.$$

Suppose that the susceptibility vanishes outside a large sphere $\chi(\boldsymbol{r}) = 0, |\boldsymbol{r}| > R_M$, then the solution to the above d'Alembert equation satisfies (See Ref. [12], Eq. 14.10)

---

[1] This is the convention set up in the presentation of Mie scattering theory in the original paper by Gustav Mie [5], and is thus followed by various other sources of literature on light scattering (e.g. Refs. [6, 7]) as well. Unfortunately, the usual convention in electrodynamics (e.g. Refs. [8, 9]) is just the opposite $\boldsymbol{E}(\boldsymbol{r},t) = \boldsymbol{E}(\boldsymbol{r})e^{-i\omega t}$. This discrepancy causes a sign difference in the imaginary part of the susceptibility $\operatorname{Im}\chi(\boldsymbol{r})$, among other things. In what follows, whenever necessary, we quote a literature result after paraphrasing it to accommodate to the $e^{i\omega t}$ convention.

[2] This is an idealization of the scenario in reality, where the charges and currents responsible for the incident light beam are sufficiently far from the scattering medium, and neither the charge distribution nor the current distribution of the beam source is seriously affected by the scattered light.

$$\begin{aligned}\boldsymbol{A}_{\rm sc}(\boldsymbol{r},t) &= \frac{\mu_0}{4\pi}\iiint_{|\boldsymbol{r}|\leq R_M}\frac{\boldsymbol{J}_P\left(\boldsymbol{r}',t-\frac{|\boldsymbol{r}-\boldsymbol{r}'|}{c}\right)}{|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'-\frac{1}{4\pi}\oiint_{|\boldsymbol{r}|=R_M}\left[\frac{2\frac{\partial}{\partial t}\boldsymbol{A}_{\rm sc}\left(\boldsymbol{r}',t-\frac{|\boldsymbol{r}-\boldsymbol{r}'|}{c}\right)}{c|\boldsymbol{r}-\boldsymbol{r}'|}(\boldsymbol{n}'\cdot\nabla')|\boldsymbol{r}-\boldsymbol{r}'|\right]\mathrm{d}\,S'\\ &\quad -\frac{1}{4\pi}\oiint_{|\boldsymbol{r}|=R_M}\left[\frac{(\boldsymbol{n}'\cdot\nabla')\boldsymbol{A}_{\rm sc}\left(\boldsymbol{r}',t-\frac{|\boldsymbol{r}-\boldsymbol{r}'|}{c}\right)}{|\boldsymbol{r}-\boldsymbol{r}'|}-\boldsymbol{A}_{\rm sc}\left(\boldsymbol{r}',t-\frac{|\boldsymbol{r}-\boldsymbol{r}'|}{c}\right)(\boldsymbol{n}'\cdot\nabla')\frac{1}{|\boldsymbol{r}-\boldsymbol{r}'|}\right]\mathrm{d}\,S' \qquad (2)\end{aligned}$$

It is a physical requirement that the scattering field has the asymptotic behavior $\boldsymbol{A}_{\rm sc}(\boldsymbol{r},t)=\boldsymbol{F}(\boldsymbol{r}/|\boldsymbol{r}|)e^{i\omega t-ik|\boldsymbol{r}|}/|\boldsymbol{r}|(1+O(1/|\boldsymbol{r}|))$ as $|\boldsymbol{r}|\to+\infty$, so the surface terms[3] in Eq. 2 vanishes in the $R_M\to+\infty$ limit.

In other words, the scattering magnetic field can be represented by the following retarded potential formula[4]:

$$\begin{aligned}\boldsymbol{B}_{\rm sc}(\boldsymbol{r},t)=\nabla\times\boldsymbol{A}_{\rm sc}(\boldsymbol{r},t) &= \frac{1}{c^2}\nabla\times\iiint\frac{\chi(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\frac{\partial}{\partial t}\boldsymbol{E}\left(\boldsymbol{r}',t-\frac{|\boldsymbol{r}-\boldsymbol{r}'|}{c}\right)\mathrm{d}^3\boldsymbol{r}'\\ &= \frac{i\omega e^{i\omega t}}{c^2}\nabla\times\iiint\frac{\chi(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'. \qquad (3)\end{aligned}$$

Now, the Born equation (Eq. 1) for light scattering emerges after we substitute the expression of $\boldsymbol{B}_{\rm sc}(\boldsymbol{r},t)$ (Eq. 3) into the relation $\nabla\times\boldsymbol{B}_{\rm sc}=\mu_0\left(\boldsymbol{J}_P+\epsilon_0\partial\boldsymbol{E}_{\rm sc}/\partial t\right)$.

As shown above, the Born equation can be rigorously derived from the Maxwell equations, and it gives a clear physical picture of the microscopic mechanism of light scattering – it directly attributes the distortion of the incident wave via the light irradiation emitted from the electrically-driven molecules that constitute the dielectric medium. For light scattered by a homogeneous medium occupying a bounded volume[5] satisfying certain geometric and topological constraints, with refractive index satisfying $n=\sqrt{1+\chi}>0$ or $\operatorname{Im} n<0$, Claus Müller has proved (see pp. 297-320 in Ref. [8] for an abridged presentation) that the Maxwell equations for scattering problems are well-posed. In other words, when the incident field $\boldsymbol{E}_{\rm inc}$ satisfies the Helmholtz equation $(\nabla^2+k^2)\boldsymbol{E}_{\rm inc}(\boldsymbol{r})=\boldsymbol{0}$ and the divergence-free condition $\nabla\cdot\boldsymbol{E}_{\rm inc}(\boldsymbol{r})=0$ (as dictated by the Maxwell equations[6]), the Born equation admits a unique solution $\boldsymbol{E}(\boldsymbol{r})$ (which also solves the Maxwell equations), and the output $\boldsymbol{E}(\boldsymbol{r})$ depends continuously on the input $\boldsymbol{E}_{\rm inc}(\boldsymbol{r})$. Müller has also proved that the Maxwell equations are well-posed if the susceptibility $\chi(\boldsymbol{r})$ is continuously differentiable in the entire space (pp. 285-297 in Ref. [8]), or if the homogeneous medium is a perfect conductor (pp. 320-326 in Ref. [8]).

However, in practice, one often needs to study the solvability of the Born equation for arbitrary (say, merely differentiable or square integrable) incident fields $\boldsymbol{E}_{\rm inc}$ that do not necessarily satisfy the Helmholtz equation, due to factors such as model imperfections and numerical approximations in the incident beam profile $\boldsymbol{E}_{\rm inc}(\boldsymbol{r})$, as well as iterative procedures (like successive Born approximations) that may not necessarily preserve the Helmholtz equation. Therefore, in general, the mathematical well-posedness of the Born equation (in the form of Eq. 1) is a much stronger result than the well-posedness of the Maxwell equations. When the spatial distribution of dielectric susceptibility $\chi(\boldsymbol{r})$ has bounded second-order partial derivatives in the entire space, optical scattering problems for square-integrable fields can be formulated in the framework of Schrödinger methods (see Ref. [13], pp. 197-199). In case the spatial distribution of dielectric susceptibility $\chi(\boldsymbol{r})$ is continuously differentiable in the entire space, and the gradient satisfies the Lipschitz-Hölder inequality $|\nabla\chi(\boldsymbol{r})-\nabla'\chi(\boldsymbol{r}')|\leq L_1|\boldsymbol{r}-\boldsymbol{r}'|^{\alpha},\forall\boldsymbol{r},\boldsymbol{r}'\in\mathbb{R}^3$ $(0<\alpha\leq 1)$, the electric field will be continuous everywhere, and the corresponding Born equation has a well-understood mathematical structure (See Ref. [14], Chap. 9). Recent literature [15] has also extended the mathematical well-posedness of the Born equation to cases where the susceptibility satisfies the Lipschitz-Hölder inequality $|\chi(\boldsymbol{r})-\chi(\boldsymbol{r}')|\leq L_0|\boldsymbol{r}-\boldsymbol{r}'|^\alpha,\forall\boldsymbol{r},\boldsymbol{r}'\in\mathbb{R}^3$ $(0<\alpha\leq 1)$. However, in typical experimental geometries for light scattering on dielectric media, the distributions of dielectric susceptibility $\chi(\boldsymbol{r})$ is piecewise constant, and the electric field has discontinuities. The presence of discontinuities complicates the mathematical structure of the Born equation, and its understanding has been limited to some numerical studies of the integral operator related to sphere scattering [16].

---

[3] The surface terms do not vanish for the incident field. Therefore, the incident field can assume non-zero values despite the fact that it satisfies a homogeneous set of partial differential equations.

[4] The choice of the retarded Green's function (with retarded time $t'=t-c^{-1}|\boldsymbol{r}-\boldsymbol{r}'|$) yields a far field as asymptotically proportional to $e^{-ik|\boldsymbol{r}|}/|\boldsymbol{r}|$. Although the choice of the advanced Green's function (with advanced time $t'=t+c^{-1}|\boldsymbol{r}-\boldsymbol{r}'|$) seems an ostensibly feasible solution to the d'Alembert equation, it violates the physical law of causality and ultimately undermines the well-posedness of the Born equation.

[5] Hereafter, we use the word "volume" to denote either a three-dimensional region depicting a solid shape or a number that describes the amount of space occupied by that region, the exact meaning of which is usually clear from context. We also use the abbreviation "dielectric volume" to mean "the region occupied by the dielectric medium"; similarly, "dielectric boundary" refers to the boundary of the region occupied by the dielectric medium.

[6] In Müller's proof, the Helmholtz equation and the divergence-free condition are both allowed to break down within a bounded volume outside the scatterer (where $\chi(\boldsymbol{r})=0$). In other words, it accommodates the situation where the incident beam is pumped by charges and currents lying in a bounded volume.

Later afterwards in this supplementary material, we will provide an in-depth analysis for light scattered by a homogeneous dielectric medium occupying a bounded volume V, inside which $\chi(\boldsymbol{r})=\chi$, outside which $\chi(\boldsymbol{r})=0$, then the triple integral on the right-hand side of the Born equation can be rewritten as the integral inside the volume V:

$$\iiint\frac{\chi(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'=\chi\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}',$$

and we will be mainly interested in the Born equation inside the volume V as well:[7]

$$(1+\chi)\boldsymbol{E}(\boldsymbol{r})=\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})+\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}',\quad \boldsymbol{r}\in V. \tag{4}$$

For a dissipative/absorptive medium, we have $\operatorname{Im}\chi<0$; for a non-dissipative/non-absorptive medium, we have $\operatorname{Im}\chi=0$.[8] We will not be discussing perfectly conducting material where $\operatorname{Im}\chi=-\infty$, as the boundary condition of electromagnetic field at a perfectly conducting metal surface is relatively simple, and the related scattering problems have been well understood [18, 19]. Besides the detailed analysis for the homogeneous dielectric media, we will also briefly mention natural generalizations of our arguments to inhomogeneous and anisotropic media.

In Section S1 of this supplementary material, we will establish the following result concerning the solvability of the Born equation (Eq. 4).

Theorem 0.1 *For dissipative dielectric media* $\operatorname{Im}\chi<0$*, the Born equation (Eq. 4) has a unique solution for every incident beam satisfying the square-integrability condition* $\iiint_V|\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}<+\infty$*, and the solution satisfies the bound estimate*

$$\iiint_V|\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}\leq\frac{|\chi|^2}{|\operatorname{Im}\chi|^2}\iiint_V|\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}. \tag{5}$$

Moreover, when $\chi\notin(-\infty,-1]$*, the Born equation (Eq. 4) has a solution for every square-integrable* $\boldsymbol{E}_{\mathrm{inc}}$ *if and only if the corresponding homogeneous equation* (*with* $\boldsymbol{E}_{\mathrm{inc}}=\boldsymbol{0}$) *admits only a trivial solution* $\boldsymbol{E}=\boldsymbol{0}$. ▌

To arrive at Theorem 0.1, we will show, in Section S1, that for light scattering problems formulated as above, a certain type of energy conservation law (or a form of "generalized optical theorem"[9]) holds

$$\sigma_{\mathrm{sc}}+\sigma_{\mathrm{abs}}=\frac{4\pi}{k}\operatorname{Im}\left[-\frac{\chi k^2}{4\pi}\iiint_V\boldsymbol{E}^*_{\mathrm{inc}}(\boldsymbol{r})\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right] \tag{6}$$

---

[7] There are two reasons why the electric field inside the dielectric is more essential than the field outside. From the perspective of the Born equation, the electric field outside the dielectric volume can be fully determined by the incident field and an integral involving the internal field $\boldsymbol{E}(\boldsymbol{r}),\boldsymbol{r}\in V$, thus playing a subordinate rôle in scattering problems. From the perspective of the Maxwell equations, the electromagnetic field in the exterior region is also completely determined by the tangential components of the electric field on the dielectric boundary and the radiation condition at infinity, according to William Saunders' theorem on the unique and existent solution to electromagnetic diffraction problems [17].

[8] The non-zero imaginary part in dissipative/absorptive media can arise from different microscopic mechanisms [9], such as (a) the radiation damping of bound electrons and non-radiative decay of excited states in insulators; (b) the Joule heating induced by the motion of free electrons in conductors; (c) a combination of mechanisms *a* and *b* in semiconductors, and so on. Nonetheless, in terms of the linear response to time-harmonic electromagnetic fields (i.e. monochromatic "light"), all these materials can be phenomenologically described as a "dielectric" with complex-valued susceptibility χ. In the light of this, the term "dielectric" will be used hereafter in a broad sense, to cover non-conducting insulators as well as conducting metals. For convenience, we may also write $\operatorname{Im}\chi=-\sigma_{\mathrm{cond}}/\omega$ where $\sigma_{\mathrm{cond}}\geq 0$ is the phenomenological electric conductivity.

[9] In the literature [20, 21, 22], "generalized optical theorem" was used as an umbrella term for a variety of formulas that were established in a similar vein as the classical optical theorem, which relates the total extinction cross-section to the imaginary part of the forward scattering amplitude. Indeed, when the incident field $\boldsymbol{E}_{\mathrm{inc}}$ represents a plane wave $\boldsymbol{e}_{\mathrm{inc}}e^{-ikz}$, then the right-hand side of Eq. 6 is equal to the imaginary part of the forward scattering amplitude $\frac{4\pi}{k}\operatorname{Im}\left[-\frac{\chi k^2}{4\pi}\iiint_V\boldsymbol{e}^*_{\mathrm{inc}}(\boldsymbol{r})\cdot\boldsymbol{E}(\boldsymbol{r})e^{ikz}\,\mathrm{d}^3\boldsymbol{r}\right]$, as in the classical optical theorem. When the incident field $\boldsymbol{E}_{\mathrm{inc}}$ is arbitrary, the right-hand side of Eq. 6 can be phenomenologically interpreted as the electromagnetic work done in unit time by interaction of the incident field $\boldsymbol{E}_{\mathrm{inc}}$ with the polarization current density $\boldsymbol{J}_P(\boldsymbol{r})=i\omega\epsilon_0\chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})$.

whenever the electric energy is finite: $\iiint_V \boldsymbol{E}^*(\boldsymbol{r})\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r} < +\infty, \iiint_V \boldsymbol{E}^*_{\mathrm{inc}}(\boldsymbol{r})\cdot\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r} < +\infty$. Here, the total scattering cross-section σ_{sc} (which may be regarded as electromagnetic energy that is radiated, in unit time, to infinite distances) is given by

$$\sigma_{\mathrm{sc}} := \frac{|\chi|^2 k^4}{16\pi^2} \oiint_{|\boldsymbol{n}|=1} \left[\boldsymbol{n}\times\iiint_V \boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right]\cdot\left[\boldsymbol{n}\times\iiint_V \boldsymbol{E}^*(\boldsymbol{r}')e^{-ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right]\mathrm{d}\,\Omega \geq 0$$

where $\mathrm{d}\,\Omega = \sin\theta\,\mathrm{d}\,\theta\,\mathrm{d}\,\phi$ is the infinitesimal solid angle; while the total absorption cross-section σ_{abs} (which may be phenomenologically interpreted as dissipation of electromagnetic energy in unit time due to Joule heating) is given by

$$\sigma_{\mathrm{abs}} := \operatorname{Im}\left[-\chi k\iiint_V \boldsymbol{E}^*(\boldsymbol{r})\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right] \geq 0.$$

Although our major arguments will be carried out for the homogeneous dielectric in a bounded volume (Eq. 4), most of our conclusions also have natural generalizations to the Born equation related to an arbitrary spatial distribution of dielectric susceptibility $\chi(\boldsymbol{r})$ that vanishes outside a bounded volume (Eq. 1).

In Section S2 of this supplementary material, we will extend the quantitative arguments in Section S1, and pave way for a practical solution to the light scattering problem using the evolution semigroups. The major conclusion of Section S2 can be summarized in the following theorem.

Theorem 0.2 *If the dielectric has a smooth boundary ∂V and connected exterior volume $\mathbb{R}^3\setminus(V\cup\partial V)$, with its susceptibility χ satisfying $\operatorname{Im}\chi \leq 0$ and $\chi \neq -2$, then the Born equation (Eq. 4) has a unique solution for every incident beam that satisfies both the energy condition $\iiint_V |\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r} < +\infty$ and the transversality condition $\nabla\cdot\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r}) = 0, \boldsymbol{r}\in V$. Moreover, the solution to the Born equation (Eq. 4) encounters only a discrete set of singularities ("optical resonance modes") in the entire complex χ-plane.* ▮

As a result, the light scattering problem can be robustly solved except in a neighborhood of $\chi = -2$ where intrinsic numerical instabilities may occasionally arise, due to unbounded electric enhancement ratio

$$\sup_{0<\iiint_V |\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}<+\infty}\left[\iiint_V |\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}\bigg/\iiint_V |\boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}\right] = +\infty$$

that may accompany the Born equation (Eq. 4) in the case of $\chi = -2$.

The rest part of this supplementary material serves as a technical presentation of the mathematical arguments involved in the analytical studies of the Born equation. The presentation will be organized in standard mathematical format, where a major claim is highlighted as a numbered **Lemma**, **Proposition**, **Theorem** or **Corollary**. The statements of these claims maintain the logic flow, which is further substantiated by the accompanying passages that start with the label **Proof** and end with the sign▮.

In what follows, the notation $n = \sqrt{1+\chi}$ ($\operatorname{Im} n \leq 0$) will be reserved for the refractive index, the symbol $k = |\boldsymbol{k}|$ for the wave number (the modulus of the wave vector $\boldsymbol{k}$), and the letter c for the speed of light — out of respect to their prime importance in formulating the physical problem of light scattering. Neither n nor k in italic faces will be used as integer indices in subscripts or superscripts, and the lower-case c in italic face will not be used to represent a generic constant. The notation $\boldsymbol{n}$ in bold italic face will always stand for a unit vector ($|\boldsymbol{n}| = 1$) in various contexts. We also differentiate the use of symbols ϵ and ε, the former being reserved for the dielectric constant $\epsilon = (1+\chi)\epsilon_0$ as in physical literature, while the latter stands for an infinitesimal quantity in various mathematical contexts. When we speak of "wavelength", we mean $\lambda_{\mathrm{wave}} = 2\pi/|\boldsymbol{k}|$, and as a compromise, the plain λ will be used in the spectral analysis of linear operators.

In this supplementary material, we strive to keep the presentation accessible to both the mathematically and physically inclined audience. In particular, for the sake of self-consistency, we will explain, in (more or less) pedagogical detail, some standard mathematical terminologies and tools that may appear foreign in the physical literature on electromagnetic scattering. Following a mathematically rigorous **Proof**, there may be a **Remark** that accommodates physical interpretations of the immediate results, as well as possible cross references.

Acknowledgments This work is part of the thesis project supervised and supported by Prof. Xiaowei Zhuang of Harvard University.

S1. SPECTRAL PROPERTIES OF ELECTROMAGNETIC SCATTERING

In this section, we will establish the claim that the Born equation for light scattering is mathematically well-posed for dissipative media, in terms of uniqueness, existence and stability, based on the generalized optical theorem.

We will prove in S1.1 and S1.2, that the "energetically admissible" solutions to the Born equation are unique and existent, when light is scattered by dielectric media occupying some "regular" regions[10] $V \subset \mathbb{R}^3$ in the three-dimensional space $\mathbb{R}^3$. Furthermore, in S1.2.3, we will show that the solution to the Born equation is also stable with respect to small perturbations in the incident field $\boldsymbol{E}_{\text{inc}}$, small deviations in the wavelength $\lambda_{\text{wave}} = 2\pi/k$, and small distortions in the regular shape V of the scattering medium, and the solution is also complex-analytic with respect to the susceptibility χ for $\operatorname{Im}\chi < 0$. In the meantime, we will also outline the spectrum of the Born equation, by pinpointing the values of complex susceptibilities χ that may either fail the uniqueness theorem or the existence theorem.

Before embarking on the proof of the well-posedness of the Born equation in S1.1 and S1.2, we need to precisely define "shape regularity" in terms of geometric and topological constraints on the dielectric volume, as well as specify the requirements for "energy admissibility" in terms of analytic properties of electric fields. We also need to demonstrate that our definitions about "regularity" and "admissibility" are compatible with the Born equation. These preparations form a separate subsection S1.0, as immediately follows.

S1.0. Shape Regularity, Energy Admissibility, and the Born Equation

S1.0.1. Regular Shapes and Energetically Admissible Vector Fields

First, we recapitulate some standard topological terminologies (cf. Ref. [23, 24]) that apply to a region $M \subset \mathbb{R}^3$.

Definition We say that a region $M \subset \mathbb{R}^3$ is *bounded* if there exists a finite constant $B_M < +\infty$ such that $|\boldsymbol{r}| \leq B_M, \forall \boldsymbol{r} \in M$. We say that the region $M \subset \mathbb{R}^3$ is *compact* if it is a bounded set satisfying $\operatorname{Cl}(M) = M$. Here, the *topological closure* of a non-empty subset $A \subset \mathbb{R}^3$ in $\mathbb{R}^3$ is defined as $\operatorname{Cl}(A) := \{\boldsymbol{r} \in \mathbb{R}^3 | \exists$ a sequence $\{\boldsymbol{r}_s \in A | s = 1, 2, \dots\}$, such that $\lim_{s\to\infty} |\boldsymbol{r}_s - \boldsymbol{r}| = 0\}$. In other words, $\operatorname{Cl}(A)$ is the totality of points that can be expressed as the limit of a sequence in A. Axiomatically, the topological closure of the empty set is defined to be itself: $\operatorname{Cl}(\emptyset) = \emptyset$. We write $A^\natural \Subset A$ (and say that $A^\natural$ is *compactly contained* in A) if $A^\natural \subset \operatorname{Cl}(A^\natural) \subset A$ and $\operatorname{Cl}(A^\natural)$ is bounded. The *topological boundary* ∂M of M is given by $\partial M = \operatorname{Cl}(M) \cap \operatorname{Cl}(\mathbb{R}^3 \setminus M)$, which also satisfies $\operatorname{Cl}(M) = M \cup \partial M$.

We say that M is *open* if $\mathbb{R}^3 \setminus M$ is *closed*, i.e. $\operatorname{Cl}(\mathbb{R}^3 \setminus M) = \mathbb{R}^3 \setminus M$. Equivalently, a non-empty subset $M \subset \mathbb{R}^3$ is said to be *open* if for every point $\boldsymbol{r}_\star \in M$, there exists a positive number $\varepsilon > 0$ such that the open-ball neighborhood $O(\boldsymbol{r}_\star, \varepsilon) := \{\boldsymbol{r} \in \mathbb{R}^3 | 0 \leq |\boldsymbol{r} - \boldsymbol{r}_\star| < \varepsilon\} \subset M$ is contained in M. The empty set $\emptyset$ is open, because $\mathbb{R}^3 \setminus \emptyset = \mathbb{R}^3$ is closed. We say that M is *connected* if it cannot be further decomposed into the union of two separated and non-empty sets, that is, whenever we have the decomposition $M = M_1 \cup M_2$, where $\operatorname{Cl}(M_1) \cap M_2 = M_1 \cap \operatorname{Cl}(M_2) = \emptyset$, then either $M_1 = \emptyset$ or $M_2 = \emptyset$.

For an arbitrary set $M \subset \mathbb{R}^3$, we say that its non-empty connected subset $X \subset M$ forms a *connected component* of M if any connected set Y satisfying $X \subset Y \subset M$ must also satisfy $X = Y$. When $M \subset \mathbb{R}^3$ is an open set, then any one of its connected components is also an open set. ▮

Next, we recapitulate some standard notations for differentiable and integrable mappings (cf. Ref. [25, 26, 27, 28, 29, 30]).

Definition In the following, a complex-valued vector field $\boldsymbol{F}(\boldsymbol{r}), \boldsymbol{r} \in M$ is identified with a mapping $\boldsymbol{F} : M \longrightarrow \mathbb{C}^3$, with its *modulus* at a certain point $\boldsymbol{r}$ given by $|\boldsymbol{F}(\boldsymbol{r})| := \sqrt{\boldsymbol{F}^*(\boldsymbol{r}) \cdot \boldsymbol{F}(\boldsymbol{r})} \geq 0$, and its *support* $\operatorname{supp}\boldsymbol{F}$ defined by $\operatorname{supp}\boldsymbol{F} := \operatorname{Cl}\{\boldsymbol{r} \in M || \boldsymbol{F}(\boldsymbol{r})| > 0\}$. We say that the vector field $\boldsymbol{F} : M \longrightarrow \mathbb{C}^3$ is *bounded* on M if there exists a finite constant $B_{\boldsymbol{F}} < +\infty$ such that $|\boldsymbol{F}(\boldsymbol{r})| \leq B_{\boldsymbol{F}}, \forall \boldsymbol{r} \in M$. In general, the least possible number $b \in [0, +\infty]$ that fulfills the inequality $|\boldsymbol{F}(\boldsymbol{r})| \leq b, \forall \boldsymbol{r} \in M$ will be denoted by $\sup_{\boldsymbol{r}\in M} |\boldsymbol{F}(\boldsymbol{r})|$, and bounded vector fields accordingly satisfy $\sup_{\boldsymbol{r}\in M} |\boldsymbol{F}(\boldsymbol{r})| < +\infty$. We say that the vector field $\boldsymbol{F} : M \longrightarrow \mathbb{C}^3$ is *continuous* at a given point $\boldsymbol{r} \in M$ if for any sequence $\{\boldsymbol{r}_s \in M | s = 1, 2, \dots\}$ that converges to $\boldsymbol{r}$ (i.e. $\lim_{s\to\infty} |\boldsymbol{r}_s - \boldsymbol{r}| = 0$), we have $\lim_{s\to\infty} |\boldsymbol{F}(\boldsymbol{r}_s) - \boldsymbol{F}(\boldsymbol{r})| = 0$.

We use the abbreviation $D^{\boldsymbol{\mu}}\boldsymbol{F}, |\boldsymbol{\mu}| = m \geq 0$ to refer to any of the 3^m possible partial derivatives (in the classical sense[11]) of the vector field $\boldsymbol{F} : M \longrightarrow \mathbb{C}^3$, and $|\boldsymbol{\mu}| = 0$ corresponds to the original vector field without differentiation. For $m = 0, 1, 2, \dots$, we define the family of m-th order *pointwise continuously differentiable vector fields* on M as

$$C^m(M; \mathbb{C}^3) := \{\boldsymbol{F} : M \longrightarrow \mathbb{C}^3 | \text{The vector field } D^{\boldsymbol{\mu}}\boldsymbol{F} \text{ is continuous at every point } \boldsymbol{r} \in M, \forall |\boldsymbol{\mu}| \leq m\}.$$

---

[10] In all that follows, $A \subset B$ denotes that A is a (not necessarily proper) subset of B, and hence $A \subset B$ does not preclude the possibility of $A = B$. The inclusion relation for proper subsets satisfying $A \subset B$ and $A \neq B$ will be denoted by $A \subsetneqq B$.

[11] For example, let $\boldsymbol{u} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$ be a unit vector, then the first-order *classical* derivative $(\boldsymbol{u} \cdot \nabla)\boldsymbol{F}(\boldsymbol{r})$ at the point $\boldsymbol{r} = \boldsymbol{r}_0$ is evaluated by $\lim_{\varepsilon\to 0} [\boldsymbol{F}(\boldsymbol{r}_0 + \varepsilon\boldsymbol{u}) - \boldsymbol{F}(\boldsymbol{r}_0)]/\varepsilon$, provided that the limit exists. Here, $\boldsymbol{e}_x \cdot \nabla$ refers to $\partial/\partial x$, and so forth.

In particular, $C^0(M;\mathbb{C}^3)$ denotes the family of *pointwise continuous vector fields* on M. Noting that a continuous mapping sends a compact set to a bounded set, we may conclude that for every $\boldsymbol{F} \in C^m(M;\mathbb{C}^3)$, the vector field $D^{\boldsymbol{\nu}}\boldsymbol{F} : M \longrightarrow \mathbb{C}^3, \forall|\boldsymbol{\nu}| \leq m$ is bounded on any subset $M^\natural \Subset M$ compactly contained in M, i.e. $\|\boldsymbol{F}\|_{C^m(M^\natural;\mathbb{C}^3)} := \sum_{|\boldsymbol{\nu}|\leq m} \sup_{\boldsymbol{r}\in M^\natural} |k^{-|\boldsymbol{\nu}|}D^{\boldsymbol{\nu}}\boldsymbol{F}(\boldsymbol{r})| < +\infty$. In a similar vein, we may define the family of *pointwise continuously differentiable mappings* $C^m(M';\mathbb{C}^d)$ where $M' \subset \mathbb{R}^{d'}$ is an arbitrary subset of a Euclidean space $\mathbb{R}^{d'}$ of dimension $d' = 1, 2, 3, \dots$, and $\mathbb{C}^d$ is the d-fold ($d = 1, 2, 3, \dots$) Cartesian product of the complex plane $\mathbb{C}$.

We define the family of *infinitely differentiable vector fields* on M as $C^\infty(M;\mathbb{C}^3) = \bigcap_{m\geq 0} C^m(M;\mathbb{C}^3)$. We define the family of *infinitely differentiable and compactly supported vector fields* on an open set M as $C_0^\infty(M;\mathbb{C}^3) := \{\boldsymbol{F} \in C^\infty(M;\mathbb{C}^3)|\operatorname{supp}\boldsymbol{F} \Subset M\}$. When M is bounded and open, every member of $C_0^\infty(M;\mathbb{C}^3)$ is infinitely differentiable in M and vanishes in a certain neighborhood of the boundary ∂M.

We define the family of *p-th power integrable*[12] *vector fields* on M as

$$L^p(M;\mathbb{C}^3) := \left\{ \boldsymbol{F} : M \longrightarrow \mathbb{C}^3 \left| \|\boldsymbol{F}\|_{L^p(M;\mathbb{C}^3)} := \left(\iiint_M |\boldsymbol{F}(\boldsymbol{r})|^p \, \mathrm{d}^3\boldsymbol{r} \right)^{1/p} < +\infty \right.\right\}.$$

The space $L^p(M;\mathbb{C}^3)$ becomes a Banach space equipped with the L^p-norm $\|\cdot\|_{L^p(M;\mathbb{C}^3)}$ for $p \geq 1$. In the specific case $p = 2$, the Banach space $L^2(M;\mathbb{C}^3)$ becomes a Hilbert space, on which the L^2-norm satisfies $\|\boldsymbol{F}\|^2_{L^2(M;\mathbb{C}^3)} = \langle \boldsymbol{F}, \boldsymbol{F}\rangle_M$ with

$$\langle \boldsymbol{F}, \boldsymbol{G}\rangle_M := \iiint_M \boldsymbol{F}^*(\boldsymbol{r}) \cdot \boldsymbol{G}(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r}$$

being a scalar product satisfying the Cauchy-Schwarz inequality $|\langle \boldsymbol{F}, \boldsymbol{G}\rangle_M| \leq \|\boldsymbol{F}\|_{L^2(M;\mathbb{C}^3)}\|\boldsymbol{G}\|_{L^2(M;\mathbb{C}^3)}$. (Here, it is important to append a subscript for the scalar product $\langle \boldsymbol{F}, \boldsymbol{G}\rangle_{\mathfrak{H}}$ to highlight the related Hilbert space $\mathfrak{H}$. When we consider the Hilbert space $\mathfrak{H} = L^2(M;\mathbb{C}^3)$, we use the short-hand notation $\langle \boldsymbol{F}, \boldsymbol{G}\rangle_{L^2(M;\mathbb{C}^3)} = \langle \boldsymbol{F}, \boldsymbol{G}\rangle_M$ hereafter.)

For a generic $\boldsymbol{F} \in L^p(M;\mathbb{C}^3)$, where M is bounded and open, we may define its derivatives $D^{\boldsymbol{\mu}}\boldsymbol{F}, |\boldsymbol{\mu}| = m \geq 0$ (in the distributional sense) by the relation $\iiint_M \psi(\boldsymbol{r})D^{\boldsymbol{\mu}}\boldsymbol{F}(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r} = (-1)^{|\boldsymbol{\mu}|} \iiint_M \boldsymbol{F}(\boldsymbol{r})D^{\boldsymbol{\mu}}\psi(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r}, \forall\psi \in C_0^\infty(M;\mathbb{C})$. We define the family of *vector fields of Sobolev type*-(m, p) on M as

$$W^{m,p}(M;\mathbb{C}^3) := \left\{ \boldsymbol{F} \in L^p(M;\mathbb{C}^3) \left| \|\boldsymbol{F}\|_{W^{m,p}(M;\mathbb{C}^3)} := \left(\sum_{|\boldsymbol{\nu}|\leq m} \iiint_M |k^{-|\boldsymbol{\nu}|}D^{\boldsymbol{\nu}}\boldsymbol{F}(\boldsymbol{r})|^p \, \mathrm{d}^3\boldsymbol{r} \right)^{1/p} < +\infty \right.\right\},$$

where $m = 0, 1, 2, \dots$, and $1 \leq p < +\infty$. Naturally, we have $C^m(M';\mathbb{C}^3) \subset W^{m,p}(M;\mathbb{C}^3)$ if $M \Subset M'$. ▮

With the above definitions, we may proceed with specifying the topological and analytic conditions for the light scattering problem on a dielectric volume V.

Definition (Regular Shapes and Energetically Admissible Fields) We call the non-empty dielectric volume V "regular" if it is a bounded open set. (In non-technical terms, the volume V represents a three-dimensional bulk material that has finite spatial extent.) We call the internal electric field $\boldsymbol{E}(\boldsymbol{r}), \boldsymbol{r} \in V$ "energetically admissible" if it is square-integrable $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$ (i.e., with finite "electric energy" $\iiint_V \boldsymbol{E}^*(\boldsymbol{r}) \cdot \boldsymbol{E}(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r} < +\infty$). ▮

We note that our requirements on shape regularity are much more lenient than the conventions given by Colton and Kress' book on scattering theory (Ref. [18], Chap. 2), or Müller's book on electromagnetic waves (Ref. [8], Chap. VI). Noticeably, we have not required that V can be decomposed into finitely many connected components $V_j, j = 1, \dots, N_V < +\infty$, nor have we required that the boundary ∂V be a smooth surface. During most of our discussions later in S1, we neither require that the exterior volume $\mathbb{R}^3 \setminus (V \cup \partial V)$ be a connected set as in Ref. [8]. Meanwhile, we note that our requirements on admissible "electric fields" are also relatively lenient. We have not imposed the Helmholtz equation on $\boldsymbol{E}$ as in Ref. [8], nor have we mandated that the derivatives of $\boldsymbol{E}$ satisfy certain Lipschitz-Hölder inequalities as in Ref. [18]. Moreover, our energy admissibility conditions only concern the bounded volume $V \Subset \mathbb{R}^3$, rather than the entire space $\mathbb{R}^3$.

---

[12] Unless otherwise noted in certain circumstances (e.g. in S1.1.1 and S2.1.1), every integral appearing in this supplementary material is interpreted as a Lebesgue integral (see Ref. [31], Chap. V), and the domain of integration M is understood as a Lebesgue-measurable set (see Ref. [31], p. 62).

The rationale behind our definition of shape regularity and energy admissibility is the need to understand the solvability of the Born equation in a general context. Namely, not only do we want to deal with dielectrics whose boundary surface may or may not be smooth, but also to treat the Born equation as a differo-integral equation in its own right, without the assumption that $\boldsymbol{E}$ must be an electric field that solves the Maxwell equations.

S1.0.2. Born Equation for Energetically Admissible Electric Fields

Despite the apparent lack of boundary smoothness and the apparent absence of the Helmholtz equation and the Maxwell equations in our definition of "regular shapes" and "energetically admissible electric fields", we will see that the Born equation is still well-defined in our formulation of the regularity and admissibility conditions, as demonstrated in the following proposition.[13]

Proposition 1.1 *The Born operator* $\hat{\mathcal{B}} : C_0^\infty(V;\mathbb{C}^3) \longrightarrow C^\infty(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3)$

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) := (1+\chi)\boldsymbol{E}(\boldsymbol{r}) - \chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad \boldsymbol{r}\in V \tag{1.1}$$

is well-defined where the partial derivatives on the right-hand side of Eq. 1.1 are interpreted in the classical sense. Furthermore, Eq. 1.1 continuously extends the definition of the operator to $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$*, where the partial derivatives are interpreted in the distributional sense.*

Proof For $\boldsymbol{E} \in C_0^\infty(V;\mathbb{C}^3) \subset L^2(V;\mathbb{C}^3)$, we may define

$$(\hat{\mathcal{C}}\boldsymbol{E})(\boldsymbol{r}) := \iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \tag{1.2}$$

and use integration by parts to verify that $\hat{\mathcal{C}}\boldsymbol{E} \in C^\infty(\mathbb{R}^3;\mathbb{C}^3) \subset C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$. Therefore, we have $\hat{\mathcal{B}}\boldsymbol{E} \in C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$ for every $\boldsymbol{E}\in C_0^\infty(V;\mathbb{C}^3)$.

As we decompose the Born operator $\hat{\mathcal{B}} : C_0^\infty(V;\mathbb{C}^3) \longrightarrow C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$ in the following form

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) = (1+\chi)\boldsymbol{E}(\boldsymbol{r}) - \chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' - \chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')(e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad \boldsymbol{r}\in V, \tag{1.3}$$

we may recognize that the above formula is also meaningful as a mapping $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$, so long as the partial derivatives are interpreted in the distributional sense. There are two underlying reasons for this. On one hand, for any $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, its Newton potential

$$(\hat{\mathcal{N}}\boldsymbol{E})(\boldsymbol{r}) := -\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$$

belongs to Sobolev type-$(2,2)$, i.e. $\hat{\mathcal{N}}\boldsymbol{E} \in W^{2,2}(V;\mathbb{C}^3)$ as long as V is open and bounded (see Theorem 9.9 in Ref. [26], also Theorem 10.1.1 in Ref. [30]), so we must have $D^{\boldsymbol{\mu}}\hat{\mathcal{N}}\boldsymbol{E}\in L^2(V;\mathbb{C}^3), \forall|\boldsymbol{\mu}|=2$. In particular, we have $\nabla\times\nabla\times\hat{\mathcal{N}}\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$. Moreover, we also have the identity $\sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}\hat{\mathcal{N}}\boldsymbol{E}\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = \|\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}$ (see Theorem 9.9 in Ref. [26], also Theorem 10.1.1 in Ref. [30]). On the other hand, we may consider the following Hessian matrix

$$\begin{aligned}
&\nabla\nabla\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\\
:=&\begin{pmatrix}\frac{\partial^2}{\partial x^2} & \frac{\partial^2}{\partial x\partial y} & \frac{\partial^2}{\partial x\partial z}\\ \frac{\partial^2}{\partial y\partial x} & \frac{\partial^2}{\partial y^2} & \frac{\partial^2}{\partial y\partial z}\\ \frac{\partial^2}{\partial z\partial x} & \frac{\partial^2}{\partial z\partial y} & \frac{\partial^2}{\partial z^2}\end{pmatrix}\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\\
=&\frac{k^2}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\left\{\left[\frac{1-e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}(1+ik|\boldsymbol{r}-\boldsymbol{r}'|)}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2}\right]\nVdash + \left[3\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}(1+ik|\boldsymbol{r}-\boldsymbol{r}'|)-1}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2} - e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}\right]\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right\}
\end{aligned} \tag{1.4}$$

---

[13] From now on, the volume $V\subset\mathbb{R}^3$ is understood as a generic bounded open set, unless further geometric and/or topological constraints are imposed thereupon.

where $\mathbb{1}$ stands for the 3×3 identity matrix, and the superscript "${}^{\mathrm{T}}$" transposes a column vector to a row vector in the expression

$$\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}=\frac{\boldsymbol{r}-\boldsymbol{r}'}{|\boldsymbol{r}-\boldsymbol{r}'|}\left(\frac{\boldsymbol{r}-\boldsymbol{r}'}{|\boldsymbol{r}-\boldsymbol{r}'|}\right)^{\mathrm{T}}.$$

One may recognize that all the elements in the Hessian matrix become of order $O(|\boldsymbol{r}-\boldsymbol{r}'|^{-1})$ as $|\boldsymbol{r}-\boldsymbol{r}'|\to 0$, which is a square-integrable singularity. Therefore, we may differentiate under the integral sign

$$\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')(e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'=\iiint_V\left[(-\nabla^2+\nabla\nabla)\frac{(e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'$$

and construct a linear operator[14] $\hat{\gamma}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ by

$$(\hat{\gamma}\boldsymbol{E})(\boldsymbol{r}):=\iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'=\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')(e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$$

where the 3×3 matrix

$$\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')=\frac{k^2e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\mathbb{1}+\nabla\nabla\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}$$

satisfies the Hilbert-Schmidt bound

$$\gamma[k;V]:=\sqrt{\iiint_V\left\{\iiint_V\operatorname{Tr}\left[\hat{\Gamma}^*(\boldsymbol{r},\boldsymbol{r}')\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\right]\mathrm{d}^3\boldsymbol{r}'\right\}\mathrm{d}^3\boldsymbol{r}}<+\infty. \tag{1.5}$$

Using the Cauchy-Schwarz inequality, we may verify the bound estimate

$$\begin{aligned}\|\hat{\gamma}\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}&=\iiint_V\left|\iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'\right|^2\mathrm{d}^3\boldsymbol{r}\\&\le\iiint_V\left(\iiint_V\sqrt{\lambda_{\max}\left[\hat{\Gamma}^*(\boldsymbol{r},\boldsymbol{r}')\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\right]}|\boldsymbol{E}(\boldsymbol{r}')|\,\mathrm{d}^3\boldsymbol{r}'\right)^2\mathrm{d}^3\boldsymbol{r}\\&\le\|\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}\iiint_V\left[\iiint_V\lambda_{\max}\left[\hat{\Gamma}^*(\boldsymbol{r},\boldsymbol{r}')\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\right]\mathrm{d}^3\boldsymbol{r}'\right]\mathrm{d}^3\boldsymbol{r}\\&\le\|\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}\iiint_V\left[\iiint_V\operatorname{Tr}\left[\hat{\Gamma}^*(\boldsymbol{r},\boldsymbol{r}')\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\right]\mathrm{d}^3\boldsymbol{r}'\right]\mathrm{d}^3\boldsymbol{r}=(\gamma[k;V]\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)})^2,\end{aligned}$$

where $\lambda_{\max}$ extracts the largest eigenvalue of the 3×3 positive-semidefinite matrix $\hat{\Gamma}^*(\boldsymbol{r},\boldsymbol{r}')\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')$. Therefore, so long as $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, we must have $\hat{\mathscr{B}}\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$ as well.

Now we need to explain that there is a continuous extension from $\hat{\mathscr{B}}:C_0^\infty(V;\mathbb{C}^3)\longrightarrow C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$ to $\hat{\mathscr{B}}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$. According to a standard result (see Lemma 2.19 in Ref. [29]), the function space $C_0^\infty(V;\mathbb{C}^3)$ is dense in $L^2(V;\mathbb{C}^3)$, in the sense that every member $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$ can be approximated by a sequence $\{\boldsymbol{E}_s\in C_0^\infty(V;\mathbb{C}^3)|s=1,2,\dots\}$ in L^2-norm: $\lim_{s\to\infty}\|\boldsymbol{E}_s-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$. In the meantime, we have $\sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}\hat{\mathscr{N}}(\boldsymbol{E}_s-\boldsymbol{E})\|^2_{L^2(V;\mathbb{C}^3)}\le\sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}\hat{\mathscr{N}}(\boldsymbol{E}_s-\boldsymbol{E})\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)}=\|\boldsymbol{E}_s-\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}\to 0$ and $\|\hat{\gamma}(\boldsymbol{E}_s-\boldsymbol{E})\|_{L^2(V;\mathbb{C}^3)}\le\gamma[k;V]\|\boldsymbol{E}_s-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\to 0$, as well. Thus, we have $\lim_{s\to\infty}\|\hat{\mathscr{B}}\boldsymbol{E}_s-\hat{\mathscr{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$. In the light of this, to define $\hat{\mathscr{B}}\boldsymbol{E}$ for an arbitrary $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$ which might involve distributional derivatives, it would suffice to pick a convergent sequence $\{\boldsymbol{E}_s\in C_0^\infty(V;\mathbb{C}^3)|s=1,2,\dots\}$ that tends to $\boldsymbol{E}$ in L^2-norm, and consider the L^2-limit of the sequence $\{\hat{\mathscr{B}}\boldsymbol{E}_s\in C_0^\infty(V;\mathbb{C}^3)|s=1,2,\dots\}$, where only classical derivatives are involved in the definition of each $\hat{\mathscr{B}}\boldsymbol{E}_s$. ▮

---

[14] Actually, this operator extends to $\hat{\gamma}:L^2(V;\mathbb{C}^3)\longrightarrow C^0(\mathbb{R}^3;\mathbb{C}^3)$, because convolution of a square-integrable function with a square-integrable convolution kernel yields a continuous function, as can be verified by the Cauchy-Schwarz inequality (see Ref. [25], p. 271). It is the continuity of the vector field $\hat{\gamma}\boldsymbol{E}\in C^0(\mathbb{R}^3;\mathbb{C}^3)$ that justifies the differentiation under the integral sign, by a standard argument (sometimes known as the Leibniz integral rule) based on the Fubini theorem (see Ref. [31], p. 148 or Ref. [29], p. 25) and the Newton-Leibniz formula.

Remark Using integration by parts, one may verify that for an "electric field" $\boldsymbol{E} \in C^1(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3)$ that is both continuously differentiable and square integrable, we still have $\hat{\mathcal{N}}\boldsymbol{E} \in C^2(V;\mathbb{C}^3)$, so the distributional and classical derivatives of $\hat{\mathcal{N}}\boldsymbol{E}$ coincide up to second order. In other words, the continuous extension along the chain $C_0^\infty(V;\mathbb{C}^3) \subset C^1(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3) \subset L^2(V;\mathbb{C}^3)$ preserves the the term $\nabla\times\nabla\times\hat{\mathcal{N}}\boldsymbol{E}$ in the Born equation as classical derivatives for "physically reasonable" $\boldsymbol{E}$ fields (say, of type $C^1(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3)$) that do not necessarily vanish in a neighborhood of ∂V. So our definition of the extended Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ consistently leads to a differo-integral equation whenever $\boldsymbol{E}$ is sufficiently well-behaved. Moreover, our definition of $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is also consistent with the discrete dipole approximation method [15, 16, 32, 33, 34, 35, 36, 37] (also known as the digitized Green's function method) in the literature on light scattering (see S1.3.1 for details). ▮

S1.1. Uniqueness Theorem for Electromagnetic Scattering

In Proposition 1.1 of S1.0, we have established the fact that for regularly shaped volume V, the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is an endomorphism that sends every square-integrable field to another square integrable field, that is, $\hat{\mathcal{B}}L^2(V;\mathbb{C}^3) \subset L^2(V;\mathbb{C}^3)$.

This subsection is devoted to the proof of the fact that the solution to the Born equation $\hat{\mathcal{B}}\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ is unique under certain conditions. Namely, we will prove that the conditions $\|\boldsymbol{E}_{\text{inc}}\|_{L^2(V;\mathbb{C}^3)} = \|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$ and $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$ imply $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$, when some constraints on the susceptibility χ and/or dielectric volume V are imposed. Before this can be done, we need to establish a generalized form of optical theorem for the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$, with the help of Fourier transforms.

S1.1.1. Fourier Transform of the Born Operator

Definition (Fourier Transforms) For any $\boldsymbol{F} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$, we use the following notation

$$\tilde{\boldsymbol{F}}(\boldsymbol{q}) := \iiint\hspace{-1.2em}-\hspace{0.5em}_{\mathbb{R}^3} \boldsymbol{F}(\boldsymbol{r}) e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\, \mathrm{d}^3\boldsymbol{r} \tag{1.6a}$$

to represent its Fourier transform, and accordingly,

$$\boldsymbol{F}(\boldsymbol{r}) := \frac{1}{(2\pi)^3} \iiint\hspace{-1.2em}-\hspace{0.5em}_{\mathbb{R}^3} \tilde{\boldsymbol{F}}(\boldsymbol{q}) e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\, \mathrm{d}^3\boldsymbol{q} \tag{1.6b}$$

for the inverse Fourier transform. The pair of vectors fields $\boldsymbol{F} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$ and $\tilde{\boldsymbol{F}} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$ satisfy the Parseval-Plancherel identity (see Ref. [38], p. 104)

$$\iiint_{\mathbb{R}^3} |\boldsymbol{F}(\boldsymbol{r})|^2\, \mathrm{d}^3\boldsymbol{r} = \frac{1}{(2\pi)^3} \iiint_{\mathbb{R}^3} |\tilde{\boldsymbol{F}}(\boldsymbol{q})|^2\, \mathrm{d}^3\boldsymbol{q}, \tag{1.7}$$

which can recast into an equivalent form for $\boldsymbol{F}, \boldsymbol{G} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$ and $\tilde{\boldsymbol{F}}, \tilde{\boldsymbol{G}} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$ as

$$\langle \boldsymbol{F}, \boldsymbol{G}\rangle_{\mathbb{R}^3} = \iiint_{\mathbb{R}^3} \boldsymbol{F}^*(\boldsymbol{r}) \cdot \boldsymbol{G}(\boldsymbol{r})\, \mathrm{d}^3\boldsymbol{r} = \frac{1}{(2\pi)^3} \iiint_{\mathbb{R}^3} \tilde{\boldsymbol{F}}^*(\boldsymbol{q}) \cdot \tilde{\boldsymbol{G}}(\boldsymbol{q})\, \mathrm{d}^3\boldsymbol{q} = \frac{1}{(2\pi)^3} \langle \tilde{\boldsymbol{F}}, \tilde{\boldsymbol{G}}\rangle_{\mathbb{R}^3}. \tag{1.8}$$

Here, if we further have $\boldsymbol{F} \in L^1(\mathbb{R}^3;\mathbb{C}^3) \cap L^2(\mathbb{R}^3;\mathbb{C}^3)$, then Eq. 1.6a coincides with a usual integral $\tilde{\boldsymbol{F}}(\boldsymbol{q}) = \iiint_{\mathbb{R}^3} \boldsymbol{F}(\boldsymbol{r}) e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\, \mathrm{d}^3\boldsymbol{r}$ that converges absolutely. Likewise, if $\tilde{\boldsymbol{F}} \in L^1(\mathbb{R}^3;\mathbb{C}^3) \cap L^2(\mathbb{R}^3;\mathbb{C}^3)$, then Eq. 1.6b becomes a usual integral $\boldsymbol{F}(\boldsymbol{r}) = \frac{1}{(2\pi)^3} \iiint_{\mathbb{R}^3} \tilde{\boldsymbol{F}}(\boldsymbol{q}) e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\, \mathrm{d}^3\boldsymbol{q}$. For generic cases $\boldsymbol{F}, \tilde{\boldsymbol{F}} \in L^2(\mathbb{R}^3;\mathbb{C}^3)$, Eqs. 1.6a and 1.6b still have precise meanings as continuous extensions of the Fourier transform from $L^1(\mathbb{R}^3;\mathbb{C}^3) \cap L^2(\mathbb{R}^3;\mathbb{C}^3)$ to $L^2(\mathbb{R}^3;\mathbb{C}^3)$. (See Ref. [38], p. 104 for further technical explanations of the operation "$\iiint\hspace{-1.2em}-\hspace{0.5em}$".) In general, when "$\iiint\hspace{-1.2em}-\hspace{0.5em}$" appears in an equality, it means that the left- and right-hand sides agree with possible exceptions on a subset $V_0 \subset \mathbb{R}^3$ of zero volume (i.e. $\iiint_{V_0} \mathrm{d}^3\boldsymbol{r} = 0$ or $\iiint_{V_0} \mathrm{d}^3\boldsymbol{q} = 0$ depending on context). ▮

Proposition 1.2 *For a bounded and open dielectric volume V and a finitely large susceptibility $|\chi| < +\infty$, the Born operator can be recast into $\hat{\mathcal{B}} = (1+\chi)\hat{I} - \chi\hat{\mathcal{D}} - \chi\hat{\gamma} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$. Here, $\hat{I} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is* *the identity operator;* $\hat{\mathcal{D}}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ *is the "dipole radiation" operator*[15] *defined by*

$$(\hat{\mathcal{D}}\boldsymbol{E})(\boldsymbol{r}) := -\frac{1}{(2\pi)^3}\fint\!\!\!\fint\!\!\!\fint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad \textit{with } \tilde{\boldsymbol{E}}(\boldsymbol{q}) = \iiint_V \boldsymbol{E}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r},\quad \forall\boldsymbol{q}\in\mathbb{R}^3; \tag{1.9}$$

and $\hat{\gamma}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ *is given by*

$$(\hat{\gamma}\boldsymbol{E})(\boldsymbol{r}) := \iiint_V \hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' = \iiint_V\left(\frac{k^2e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\nVdash + \nabla\nabla\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'.$$

Furthermore, we have the bound estimates $\|\hat{\mathcal{D}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$*, and* $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq (|1+\chi| + |\chi| + |\chi|k^2R_V^2)\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$*, where* $R_V := \min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ *is the minimal radius of all circumscribed spheres.*[16]

Proof We point out that for any $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, we have the following Fourier inversion formulae for second order partial derivatives in the distributional sense

$$(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\fint\!\!\!\fint\!\!\!\fint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad \forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}. \tag{1.10}$$

We defer the rigorous derivation of Eq. 1.10 to S1.3.2.

As a specific case of Eq. 1.10, we may confirm the represenation of the "dipole radiation" operator $\hat{\mathcal{D}}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ as given in Eq. 1.9. Consequently, we have the following estimate that derives from the Parseval-Plancherel identity (Eq. 1.7)

$$\|\hat{\mathcal{D}}\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)} \leq \|\hat{\mathcal{D}}\boldsymbol{E}\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\left|\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}\right|^2\mathrm{d}^3\boldsymbol{q} \leq \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}|\tilde{\boldsymbol{E}}(\boldsymbol{q})|^2\,\mathrm{d}^3\boldsymbol{q} = \|\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}.$$

Meanwhile, we have the estimate $\|\hat{\gamma}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq k^2R_V^2\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$ (see S1.3.3 for a detailed proof). It is then evident that

$$\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq |1+\chi|\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} + |\chi|\|\hat{\mathcal{D}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} + |\chi|\|\hat{\gamma}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq \left(|1+\chi| + |\chi| + |\chi|k^2R_V^2\right)\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}.$$

In other words, let the operator norm of the Born operator be defined as $\|\hat{\mathcal{B}}\|_{L^2(V;\mathbb{C}^3)} := \min\{b>0 | \|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq b\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}, \forall\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\}$, then we have $\|\hat{\mathcal{B}}\|_{L^2(V;\mathbb{C}^3)} \leq |1+\chi| + |\chi|(1+k^2R_V^2) < +\infty$. The mapping $\hat{\mathcal{B}} = (1+\chi)\hat{I} - \chi\hat{\mathcal{D}} - \chi\hat{\gamma}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is thus a bounded linear operator, hence a continuous linear transformation. ∎

Remark Later on, we will also write $\hat{\mathcal{G}} = \hat{\gamma} + \hat{\mathcal{D}} - \hat{I}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$, where

$$(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r}) := \iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' - \frac{1}{(2\pi)^3}\fint\!\!\!\fint\!\!\!\fint_{\mathbb{R}^3}\frac{\boldsymbol{q}[\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}$$

so that the Born operator can be cast into $\hat{\mathcal{B}} = \hat{I} - \chi\hat{\mathcal{G}}: L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$.

---

[15] Recall that the electric radiation at point $\boldsymbol{r}$ generated by an oscillating dipole located at point $\boldsymbol{r}'$ reads $\boldsymbol{E}(\boldsymbol{r},t) = (4\pi\epsilon_0)^{-1}\nabla\times\nabla\times([\boldsymbol{p}]/|\boldsymbol{r}-\boldsymbol{r}'|)$, where $[\boldsymbol{p}] = \boldsymbol{p}(t')$ is the dipole moment at retarded time $t' = t - c^{-1}|\boldsymbol{r}-\boldsymbol{r}'|$. This resembles the formula $\chi(\hat{\mathcal{D}}\boldsymbol{E})(\boldsymbol{r}) = (4\pi\epsilon_0)^{-1}\nabla\times\nabla\times\iiint_V\boldsymbol{P}(\boldsymbol{r}')|\boldsymbol{r}-\boldsymbol{r}'|^{-1}\,\mathrm{d}^3\boldsymbol{r}'$, hence the name "dipole radiation" operator.

[16] It can be verified that R_V is a well-defined finite number, for two reasons: (1) for every fixed $\boldsymbol{r}$, the function $|\boldsymbol{r}-\boldsymbol{r}'|$ is continuous in $\boldsymbol{r}'$, and thus can attain its maximum in a compact set $V\cup\partial V \Subset \mathbb{R}^3$, so that $\sup_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}| = \max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}| < +\infty$; (2) As the supremum of a family of continuous functions parameterized by $\boldsymbol{r}'$, the function $\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ is lower semi-continuous with respect to $\boldsymbol{r}$, and can attain its minimum when $\boldsymbol{r}$ ranges in any compact subset of $\mathbb{R}^3$, so $\inf_{|\boldsymbol{r}|\leq 2\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'|}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}| = \min_{|\boldsymbol{r}|\leq 2\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'|}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ is a well-defined finite number that is not greater than $\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ for any $\boldsymbol{r}\in\mathbb{R}^3$. In other words, the minimal radius of all circumscribed spheres $R_V := \min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}| = \inf_{|\boldsymbol{r}|\leq 2\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'|}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ is well-defined, and there exists at least one point $\boldsymbol{r}_0\in\mathbb{R}^3$ such that $V\subset O(\boldsymbol{r}_0,R_V)$. The concept of R_V is different from the circumradius R_{circ} of a polyhedron, which is defined as the radius of the circumsphere that touches all the vertices of the polyhedron. If V has the shape of a polyhedron, we have $R_V \leq R_{\text{circ}}$.

Based on some detailed derivations in S1.3.3 and S1.3.4, we may obtain a refined estimate of operator norms as

$$\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)} \leq 1+\min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, 3\sqrt{2}kR_V\right\}+\min\left\{\frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{5}{8}kR_V\right\}, \tag{1.11}$$

and

$$\|\hat{\mathcal{G}}+\hat{I}\|_{L^2(V;\mathbb{C}^3)} \geq \frac{\sqrt{2}}{3}kr_V = \frac{\sqrt{2}}{3}k\sup\{R\in\mathbb{R}|O(\boldsymbol{r}_0,R)\subset V, \boldsymbol{r}_0\in\mathbb{R}^3\}. \tag{1.12}$$

Here, the quantity r_V represents the maximal radius of all inscribed spheres. ▮

Corollary 1.1 *The Born operator* $\hat{\mathcal{B}} = (1+\chi)\hat{I}-\chi\hat{\mathcal{D}}-\chi\hat{\gamma} = \hat{I}-\chi\hat{\mathcal{G}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ *has a limit representation:*

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) = (1+\chi)\boldsymbol{E}(\boldsymbol{r})+\chi\lim_{\varepsilon\to 0^+}\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}. \tag{1.13}$$

Proof For every $\varepsilon > 0$, and $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, we point out that we have the Fourier inversion formula (see S1.3.2 for detailed proof):

$$(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}, \quad \forall \boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}. \tag{1.14}$$

For any $\varepsilon > 0$, we can define the approximate Born operator $\hat{\mathcal{B}}^{(\varepsilon)} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ as

$$\begin{aligned}(\hat{\mathcal{B}}^{(\varepsilon)}\boldsymbol{E})(\boldsymbol{r}) &:= (1+\chi)\boldsymbol{E}(\boldsymbol{r})+\chi\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}\\ &= (1+\chi)\boldsymbol{E}(\boldsymbol{r})-\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\\ &= (1+\chi)\boldsymbol{E}(\boldsymbol{r})+\chi\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}\\ &\quad -\chi\nabla\times\nabla\times\iiint_V\frac{[e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}-1+i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'\end{aligned}$$

where the equalities hold because of Eq. 1.14. It is thus clear that $\hat{\mathcal{B}}^{(\varepsilon)} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a bounded linear operator for every $\varepsilon > 0$, and

$$(\hat{\mathcal{B}}^{(\varepsilon)}\boldsymbol{E})(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) = -\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')[e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}-e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}+\varepsilon|\boldsymbol{r}-\boldsymbol{r}'|]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$$

satisfies $\|\hat{\mathcal{B}}^{(\varepsilon)}-\hat{\mathcal{B}}\|_{L^2(V;\mathbb{C}^3)} \leq C_{k;V}\varepsilon^2$ for some constant $C_{k;V}$ only dependent on the wave number k and the shape of the scattering medium V, which confirms the representation as claimed. ▮

Remark In Ref. [34], A. R. Holt *et al.* have recast the Born equation with the dyadic Green function $(1+k^{-2}\nabla\nabla)[e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}/(4\pi|\boldsymbol{r}-\boldsymbol{r}'|)]$, and have obtained the Fourier transform of the Born equation in dyadic form (Eq. 16 in Ref. [34]), which is mathematically equivalent to Eq. 1.13. ▮

S1.1.2. A Generalized Optical Theorem

Here, we will state a critical identity (generalized optical theorem) satisfied by the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ and give two independent proofs of it.

Theorem 1.1 (Generalized Optical Theorem) *The Born operator* $\hat{\mathcal{B}} = \hat{I}-\chi\hat{\mathcal{G}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ *satisfies the following functional relation:*

$$\begin{aligned}&\frac{|\chi|^2k^5}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left[\boldsymbol{n}\times\iiint_V \boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right]\cdot\left[\boldsymbol{n}\times\iiint_V \boldsymbol{E}^*(\boldsymbol{r}')e^{-ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right]\mathrm{d}\,\Omega\\ &= \operatorname{Im}\left[\chi k^2\iiint_V|\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}\right]-\operatorname{Im}\left[\chi k^2\iiint_V(\hat{\mathcal{B}}\boldsymbol{E})^*(\boldsymbol{r})\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right].\end{aligned} \tag{1.15}$$

Proof (Configuration Space Method) We first give a proof of Eq. 1.15 by computations in the configuration space ($\boldsymbol{r}$-space).

We may momentarily consider $\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3)$. Using integration by parts, we can verify that

$$\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\in C^\infty(\mathbb{R}^3;\mathbb{C}^3). \tag{1.16}$$

For $\boldsymbol{r}\in V$, we have

$$\begin{aligned}(\hat{\mathcal{B}}\boldsymbol{F})(\boldsymbol{r}) &= (1+\chi)\boldsymbol{F}(\boldsymbol{r})-\chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\\ &= (1+\chi)\boldsymbol{F}(\boldsymbol{r})+\chi\nabla^2\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'-\chi\nabla\left[\nabla\cdot\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\\ &= \boldsymbol{F}(\boldsymbol{r})-\chi k^2\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'-\chi\nabla\left[\nabla\cdot\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\end{aligned}$$

which satisfies

$$\chi\nabla\times\nabla\times(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})=\nabla\times\nabla\times\Big[\boldsymbol{F}(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{F})(\boldsymbol{r})\Big]=k^2[(1+\chi)\boldsymbol{F}(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{F})(\boldsymbol{r})]=\chi k^2\boldsymbol{F}(\boldsymbol{r})+\chi k^2(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r});$$

For $\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$, we have

$$\begin{aligned}\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r}) &= \chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\\ &= -\chi\nabla^2\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'+\chi\nabla\left[\nabla\cdot\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\\ &= \chi k^2\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'+\chi\nabla\left[\nabla\cdot\iiint_V \frac{\boldsymbol{F}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\end{aligned}$$

which satisfies

$$\chi\nabla\times\nabla\times(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})=\chi k^2(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r}).$$

As $\hat{\mathcal{G}}\boldsymbol{F}\in C^2(\mathbb{R}^3;\mathbb{C}^3)$ for every $\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3)$, the following computations are valid:

$$\begin{aligned}&\iiint_{O(\boldsymbol{0},R_M)}\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]^*\cdot\nabla\times\nabla\times\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]\,\mathrm{d}^3\boldsymbol{r}-\iiint_{O(\boldsymbol{0},R_M)}\nabla\times\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]^*\cdot\nabla\times\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]\,\mathrm{d}^3\boldsymbol{r}\\ =&-\iiint_{O(\boldsymbol{0},R_M)}\nabla\cdot\Big\{\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]^*\times\Big[\chi\nabla\times(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]\Big\}\,\mathrm{d}^3\boldsymbol{r}=\oiint_{\partial O(\boldsymbol{0},R_M)}\Big[\chi\nabla\times(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]\cdot\Big\{\Big[\chi(\hat{\mathcal{G}}\boldsymbol{F})(\boldsymbol{r})\Big]^*\times\boldsymbol{n}\Big\}\end{aligned}\tag{1.17}$$

Here, the symbol $\boldsymbol{n}$ stands for the unit outward normal vector on the spherical surface $\partial O(\boldsymbol{0},R_M)$. Now, we may take the imaginary part of both sides of Eq. 1.17 to arrive at

$$\begin{aligned}&\operatorname{Im}\langle\boldsymbol{F}-\hat{\mathcal{B}}\boldsymbol{F},\chi k^2\boldsymbol{F}\rangle=\operatorname{Im}\left\{\chi k^2\iiint_V\Big[\boldsymbol{F}(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{F})(\boldsymbol{r})\Big]^*\cdot\boldsymbol{F}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right\}\\ =&\ \frac{|\chi|^2k^5}{16\pi^2}\oiint_{|\boldsymbol{r}|=R_M}\left[\frac{\boldsymbol{r}}{|\boldsymbol{r}|}\times\iiint_V\boldsymbol{F}(\boldsymbol{r}')e^{\frac{ik\boldsymbol{r}\cdot\boldsymbol{r}'}{|\boldsymbol{r}|}}\,\mathrm{d}^3\boldsymbol{r}'\right]\cdot\left[\frac{\boldsymbol{r}}{|\boldsymbol{r}|}\times\iiint_V\boldsymbol{F}^*(\boldsymbol{r}')e^{-\frac{ik\boldsymbol{r}\cdot\boldsymbol{r}'}{|\boldsymbol{r}|}}\,\mathrm{d}^3\boldsymbol{r}'\right]\frac{\mathrm{d}S}{|\boldsymbol{r}|^2}+O\left(\frac{1}{R_M}\right),\end{aligned}$$

where we have made use of the well-known fact that

$$\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}=\frac{e^{-ik|\boldsymbol{r}|}}{4\pi|\boldsymbol{r}|}e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\left[1+O\left(\frac{1}{k|\boldsymbol{r}|}\right)\right],\quad\text{where }\boldsymbol{n}=\frac{\boldsymbol{r}}{|\boldsymbol{r}|}.$$

Eventually, taking the limit $R_M\to+\infty$, we have verified Eq. 1.15 under the assumption that $\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3)\subset L^2(V;\mathbb{C}^3)$.

Now, for a generic square-integrable vector field $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$, we may pick a sequence $\{\boldsymbol{E}_s \in C_0^\infty(V;\mathbb{C}^3) | s = 1,2,\dots\}$ that converges to $\boldsymbol{E}$ in L^2-norm: $\lim_{s\to\infty} \|\boldsymbol{E}_s - \boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$. Now, we may take limits for the sequence of identities

$$\operatorname{Im}\langle \boldsymbol{E}_s - \hat{\mathcal{B}}\boldsymbol{E}_s, \chi k^2 \boldsymbol{E}_s\rangle_V = \frac{|\chi|^2 k^5}{16\pi^2} \oiint_{|\boldsymbol{n}|=1} \left| \boldsymbol{n} \times \iiint_V \boldsymbol{E}_s(\boldsymbol{r}') e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \mathrm{d}\,\Omega, \quad s = 1,2,\dots$$

On the left-hand side, we have $\langle \boldsymbol{E}_s, \boldsymbol{E}_s\rangle_V \to \langle \boldsymbol{E}, \boldsymbol{E}\rangle_V$, as $s \to \infty$. This is because a convergent sequence is uniformly bounded $\|\boldsymbol{E}_s\|_{L^2(V;\mathbb{C}^3)} \le C < +\infty, s = 1,2,\dots$, and the Cauchy-Schwarz inequality can be applied to the scalar product:

$$|\langle \boldsymbol{E}_s, \boldsymbol{E}_s\rangle_V - \langle \boldsymbol{E}, \boldsymbol{E}\rangle_V| = |\langle \boldsymbol{E}_s - \boldsymbol{E}, \boldsymbol{E}_s\rangle_V - \langle \boldsymbol{E}, \boldsymbol{E} - \boldsymbol{E}_s\rangle_V| \le \left(C + \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\right) \|\boldsymbol{E}_s - \boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \to 0.$$

In a similar vein, we have

$$\begin{aligned} \left|\langle \hat{\mathcal{B}}\boldsymbol{E}_s, \boldsymbol{E}_s\rangle_V - \langle \hat{\mathcal{B}}\boldsymbol{E}, \boldsymbol{E}\rangle_V\right| &= \left|\langle \hat{\mathcal{B}}\boldsymbol{E}_s - \hat{\mathcal{B}}\boldsymbol{E}, \boldsymbol{E}_s\rangle_V - \langle \hat{\mathcal{B}}\boldsymbol{E}, \boldsymbol{E} - \boldsymbol{E}_s\rangle_V\right| \\ &\le \|\hat{\mathcal{B}}\|_{L^2(V;\mathbb{C}^3)} \left(C + \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\right) \|\boldsymbol{E}_s - \boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \to 0. \end{aligned}$$

These lead to $\operatorname{Im}\langle \boldsymbol{E}_s - \hat{\mathcal{B}}\boldsymbol{E}_s, \chi k^2 \boldsymbol{E}_s\rangle_V \to \operatorname{Im}\langle \boldsymbol{E} - \hat{\mathcal{B}}\boldsymbol{E}, \chi k^2 \boldsymbol{E}\rangle_V$, as $s \to \infty$. Meanwhile, for a bounded volume V, we may use the Cauchy-Schwarz inequality to conclude that the sequence

$$\left| \iiint_V \boldsymbol{E}_s(\boldsymbol{r}') e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \le \|\boldsymbol{E}_s\|^2_{L^2(V;\mathbb{C}^3)} \iiint_V \mathrm{d}^3\boldsymbol{r} \le C \iiint_V \mathrm{d}^3\boldsymbol{r}$$

is also uniformly bounded, and the pointwise convergence

$$\left| \iiint_V \left[\boldsymbol{E}_s(\boldsymbol{r}') - \boldsymbol{E}(\boldsymbol{r}')\right] e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \le \|\boldsymbol{E}_s - \boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)} \iiint_V \mathrm{d}^3\boldsymbol{r} \to 0$$

is ensured for every unit vector $\boldsymbol{n}$ satisfying $|\boldsymbol{n}| = 1$. Therefore, according to the dominated convergence theorem, we have

$$\oiint_{|\boldsymbol{n}|=1} \left| \boldsymbol{n} \times \iiint_V \boldsymbol{E}_s(\boldsymbol{r}') e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \mathrm{d}\,\Omega \to \oiint_{|\boldsymbol{n}|=1} \left| \boldsymbol{n} \times \iiint_V \boldsymbol{E}(\boldsymbol{r}') e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \mathrm{d}\,\Omega, \quad \text{as } s \to \infty.$$

Thus, we see that the generalized optical theorem can be continuously extended from $C_0^\infty(V;\mathbb{C}^3)$ to $L^2(V;\mathbb{C}^3)$, and we have

$$\operatorname{Im}\langle \boldsymbol{E} - \hat{\mathcal{B}}\boldsymbol{E}, \chi k^2 \boldsymbol{E}\rangle_V = \frac{|\chi|^2 k^5}{16\pi^2} \oiint_{|\boldsymbol{n}|=1} \left| \boldsymbol{n} \times \iiint_V \boldsymbol{E}(\boldsymbol{r}') e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'} \,\mathrm{d}^3\boldsymbol{r}' \right|^2 \mathrm{d}\,\Omega, \quad \forall \boldsymbol{E} \in L^2(V;\mathbb{C}^3)$$

just as claimed. ▮

Proof (Momentum Space Method) We give another proof of Eq. 1.15 by computations in the momentum space ($\boldsymbol{q}$-space), which is considerably more succinct.

We directly aim to evaluate the scalar product by using Eq. 1.13 in Corollary 1.1:

$$\begin{aligned} \operatorname{Im}\langle \boldsymbol{E} - \hat{\mathcal{B}}\boldsymbol{E}, \chi k^2 \boldsymbol{E}\rangle_V &= \lim_{\varepsilon\to 0^+} \operatorname{Im}\langle \boldsymbol{E} - \hat{\mathcal{B}}_\varepsilon\boldsymbol{E}, \chi k^2 \boldsymbol{E}\rangle_V \\ &= -\operatorname{Im}\left\{ \chi k^2 \iiint_V \left[\chi \lim_{\varepsilon\to 0^+} \frac{1}{(2\pi)^3} \iiint_{\mathbb{R}^3} \frac{\boldsymbol{q} \times [\boldsymbol{q} \times \tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2 - (k - i\varepsilon)^2} e^{-i\boldsymbol{q}\cdot\boldsymbol{r}} \,\mathrm{d}^3\boldsymbol{q} \right]^* \cdot \boldsymbol{E}(\boldsymbol{r}) \,\mathrm{d}^3\boldsymbol{r} \right\}. \end{aligned}$$

Now, using the Parseval-Plancherel identity (Eq. 1.8), we obtain

$$\operatorname{Im}\langle \boldsymbol{E} - \hat{\mathcal{B}}\boldsymbol{E}, \chi k^2 \boldsymbol{E}\rangle_V = -\frac{|\chi|^2 k^2}{(2\pi)^3} \operatorname{Im} \lim_{\varepsilon\to 0^+} \iiint_{\mathbb{R}^3} \frac{\boldsymbol{q} \times [\boldsymbol{q} \times \tilde{\boldsymbol{E}}^*(\boldsymbol{q})]}{(|\boldsymbol{q}| - k - i\varepsilon)(|\boldsymbol{q}| + k + i\varepsilon)} \cdot \tilde{\boldsymbol{E}}(\boldsymbol{q}) \,\mathrm{d}^3\boldsymbol{q}.$$

With the vector identity $\tilde{\boldsymbol{E}}(\boldsymbol{q}) \cdot \{\boldsymbol{q} \times [\boldsymbol{q} \times \tilde{\boldsymbol{E}}^*(\boldsymbol{q})]\} = -[\boldsymbol{q} \times \tilde{\boldsymbol{E}}(\boldsymbol{q})] \cdot [\boldsymbol{q} \times \tilde{\boldsymbol{E}}^*(\boldsymbol{q})] = -|\boldsymbol{q} \times \tilde{\boldsymbol{E}}(\boldsymbol{q})|^2$, and the Plemelj jump relation

$$\operatorname{Im} \lim_{\varepsilon\to 0^+} \int_0^{+\infty} \frac{f(q) \,\mathrm{d}\,q}{q - k - i\varepsilon} = \pi f(k),$$

we arrive at the generalized optical theorem

$$\mathrm{Im}\,\langle \boldsymbol{E}-\hat{\mathcal{B}}\boldsymbol{E},\chi k^2\boldsymbol{E}\rangle_V=\frac{|\chi|^2k^2}{(2\pi)^3}\mathrm{Im}\lim_{\varepsilon\to0^+}\int_0^{+\infty}|\boldsymbol{q}|^2\,\mathrm{d}\,|\boldsymbol{q}|\oiint\mathrm{d}\,\Omega\frac{|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})|^2}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}=\frac{|\chi|^2k^5}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\tilde{\boldsymbol{E}}(k\boldsymbol{n})\right|^2\mathrm{d}\,\Omega.$$

In other words, we have

$$\iiint_V\left[\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}^*(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}=\mathrm{Im}\,\langle\hat{\mathcal{G}}\boldsymbol{E},\boldsymbol{E}\rangle_V=\frac{k^3}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\tilde{\boldsymbol{E}}(k\boldsymbol{n})\right|^2\mathrm{d}\,\Omega \tag{1.18}$$

for every $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$. ∎

Remark We note that the proof based on configuration space method has a direct physical interpretation. For $\boldsymbol{E}\in C^1(V\cup\partial V;\mathbb{C}^3)$, the integrand $\nabla\cdot\{[\chi(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]^*\times[\chi\nabla\times(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]\}$ may be likened to the divergence of the Poynting vector in electrodynamics. If the boundary ∂V is a smooth surface with outward normal vector $\boldsymbol{n}$, one may also directly confirm that $[\chi\nabla\times(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]\cdot\{[\chi(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]^*\times\boldsymbol{n}\}=-\boldsymbol{n}\cdot\{[\chi(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]^*\times[\chi\nabla\times(\hat{\mathcal{G}}\boldsymbol{E})(\boldsymbol{r})]\}$ is continuous across the boundary, for all vector fields $\boldsymbol{E}\in C^1(V\cup\partial V;\mathbb{C}^3)$ that do not necessarily vanish in a neighborhood of ∂V. In other words, at the dielectric boundary ∂V, the influx of the "electromagnetic energy" always cancels the efflux, because the normal component of the "Poynting vector"[17] is continuous. As a net result, the dielectric boundary is neither a source of nor a drain on "electromagnetic energy". Therefore, the total "electromagnetic energy" input from the incident beam ($\propto\mathrm{Im}\,\langle-\hat{\mathcal{B}}\boldsymbol{E},\chi k^2\boldsymbol{E}\rangle$) will be dispatched into "Joule heating" inside the dielectric medium ($\propto\mathrm{Im}\,\langle-\boldsymbol{E},\chi k^2\boldsymbol{E}\rangle$) and the "scattered light" ($\propto\oiint_{|\boldsymbol{n}|=1}|\boldsymbol{n}\times\iiint_V\boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'|^2\,\mathrm{d}\,\Omega$) that radiates to infinite distances $|\boldsymbol{r}|\to\infty$.

Meanwhile, the generalized optical theorem is not limited to light scattered by homogeneous media, as the proof based on the momentum space method allows some immediate generalizations to inhomogeneous and anisotropic media (see S1.3.6 for details). ∎

S1.1.3. The Uniqueness Theorem for Light Scattering

For physically meaningful light scattering problems related to a homogeneous medium, we require that the susceptibility χ satisfy $\mathrm{Im}\,\chi\leq0$ in the Born equation (Eq. 1.13 in Corollary 1.1). When $\chi=0$, the Born operator trivializes to the identity operator. In the following, we will establish the uniqueness theorem for the physically meaningful and non-trivial susceptibilities $\mathrm{Im}\,\chi\leq0,\chi\neq0$.

Theorem 1.2 (Uniqueness Theorem for Light Scattering) *For the Born operator* $\hat{\mathcal{B}}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ *(as defined in Eq. 1.13 in Corollary 1.1), the homogeneous equation* $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$ *admits only a trivial solution* $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$ *if one of the following two conditions holds:*

(A) The dielectric medium is dissipative with $\mathrm{Im}\,\chi<0$*;*

(B) The dielectric medium is non-dissipative with $\mathrm{Im}\,\chi=0,\chi\neq0,-1$*, its boundary* ∂V *occupies zero volume* $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$,[18] *and the exterior volume* $\mathbb{R}^3\setminus(V\cup\partial V)$ *is connected.*

Proof For situation (A), we substitute the condition $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$ into the generalized optical theorem (Theorem 1.1), and obtain

$$\frac{|\chi|^2k^5}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\iiint_V\boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right|^2\mathrm{d}\,\Omega=\mathrm{Im}\left[\chi k^2\iiint_V|\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}\right]. \tag{1.19}$$

---

[17] We put quotation marks on "electromagnetic energy" and "Poynting vector", because $\boldsymbol{E}\in C^1(V\cup\partial V;\mathbb{C}^3)$ may or may not satisfy the Helmholtz equation, thus is not necessarily a true electric field permitted by the Maxwell equations.

[18] Generally speaking, the condition $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$ is not redundant, because there exist fractal structures that are bounded open sets whose boundaries have positive "volume" (in terms of Lebesgue measure), while $\mathbb{R}^3\setminus(V\cup\partial V)$ remains connected. Let $U=\bigcup_{m=1}^{\infty}U_m\subset[0,1]$ be an open set constructed inductively via the Smith-Volterra-Cantor procedure: start with an open interval $U_1=(3/8,5/8)$ of length 1/4 in the middle of the closed interval $[0,1]$; for $m=2,3,\dots$, pick U_m as the union of 2^{m-1} open intervals, each of which has length 4^{-m} and lies in the middle of a connected component of $[0,1]\setminus\bigcup_{j=1}^{m-1}U_j$. Then we have $\int_{\partial U}\mathrm{d}\,x=\int_{[0,1]\setminus U}\mathrm{d}\,x=1/2$, and $\mathrm{Cl}\,(U)=[0,1]$. The bounded open set $V=(0,1)\times(0,1)\times U$ satisfies $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=1/2$ and $\mathbb{R}^3\setminus(V\cup\partial V)=\mathbb{R}^3\setminus[0,1]^3$ is connected. Even a bounded "regularly open" set (which equals the interior of its own topological closure) may not be immune from such pathology — see Ref. [39] for a construction of a bounded "regularly open" set that has a boundary with positive "volume".

Now that the left-hand side of Eq. 1.19 is non-negative, and the right-hand side of Eq. 1.19 is non-positive, the only possibility is that both sides vanish. As $\operatorname{Im}\chi \neq 0$, the vanishing right-hand side of Eq. 1.19 implies $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$. Actually, this argument extends to be valid inside the following circular disk

$$|\chi|^2 \min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3 \iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\} < \operatorname{Im}\chi, \quad R_V := \min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$$

which resides in the upper half plane ($\operatorname{Im}\chi > 0$). This is because the left-hand side of Eq. 1.19 is always bounded by $(4\pi)^{-1}|\chi|^2k^5 \iiint_V |\boldsymbol{E}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r} \iiint_V \mathrm{d}^3\boldsymbol{r}$ according to the Cauchy-Schwarz inequality, and the bound estimates proportional to kR_V and $(k^3\iiint_V \mathrm{d}^3\boldsymbol{r})^{2/3}$ are derived in S1.3.4.

Moreover, we see that for $\operatorname{Im}\chi < 0$, the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow \hat{\mathcal{B}}L^2(V;\mathbb{C}^3)$ is coercive by the following lower bound estimate arising from the application of the Cauchy-Schwarz inequality to the generalized optical theorem (Theorem 1.1):

$$|\chi|\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \geq |\operatorname{Im}[\chi\langle\hat{\mathcal{B}}\boldsymbol{E},\boldsymbol{E}\rangle_V]| \geq -\operatorname{Im}\chi\|\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)}. \tag{1.20}$$

The coercivity bound $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \geq (-\operatorname{Im}\chi/|\chi|)\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$ indicates the invertibility of the Born operator on its range $\hat{\mathcal{B}}L^2(V;\mathbb{C}^3)$, and the inverse mapping $\hat{\mathcal{B}}^{-1} : \hat{\mathcal{B}}L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ has bounded operator norm $\|\hat{\mathcal{B}}^{-1}\|_{\hat{\mathcal{B}}L^2(V;\mathbb{C}^3)} \leq -|\chi|/\operatorname{Im}\chi < +\infty$.

For situation (B), we point out that $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$ and Theorem 1.1 together imply $\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})] = \boldsymbol{0}$ for all $|\boldsymbol{q}| = k$. As we further note that $\tilde{\boldsymbol{E}} \in C^1(\mathbb{R}^3;\mathbb{C}^3)$ arises from the boundedness of second-order moments of $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$, we see that $\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]/(|\boldsymbol{q}|^2-k^2)$ is bounded and continuous for all $\boldsymbol{q}\in\mathbb{R}^3$. Consequently, we can use the dominated convergence theorem to prove that

$$\lim_{\varepsilon\to0^+}\iiint_{\mathbb{R}^3}\left|\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2} - \frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2}\right|^2\mathrm{d}^3\boldsymbol{q} = 0,$$

so

$$\chi\frac{1}{(2\pi)^3}{-\!\!\!\!\!\!\iiint}_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}$$

is a valid Fourier inversion formula for the corresponding term in the Born equation. Inside the dielectric volume V, we have

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) := (1+\chi)\boldsymbol{E}(\boldsymbol{r}) + \chi\frac{1}{(2\pi)^3}{-\!\!\!\!\!\!\iiint}_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} = \boldsymbol{0}, \quad \forall\boldsymbol{r}\in V\setminus V_0,$$

for some subset $V_0 \subset V$ with zero volume $\iiint_{V_0}\mathrm{d}^3\boldsymbol{r} = 0$. Outside the dielectric volume V, we have

$$\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}{-\!\!\!\!\!\!\iiint}_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} = \boldsymbol{0}, \quad \forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V\cup V_0')$$

for some subset $V_0' \subset \mathbb{R}^3\setminus(V\cup\partial V)$ with zero volume $\iiint_{V_0'}\mathrm{d}^3\boldsymbol{r} = 0$, provided that $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected. This is because the vanishing scattering amplitudes in all directions

$$\boldsymbol{f}(\boldsymbol{n}) := -\frac{\chi k^2}{4\pi}\boldsymbol{n}\times\left[\boldsymbol{n}\times\iiint_V\boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right] \equiv \boldsymbol{0}, |\boldsymbol{n}| = 1$$

imply a vanishing scattering field in the connected open set $\mathbb{R}^3\setminus(V\cup\partial V)$

$$\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r}) \ := \ \chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\mathrm{d}^3\boldsymbol{r}' \equiv \boldsymbol{0}, \quad \boldsymbol{r}\notin V\cup\partial V,$$

according to an analytic continuation procedure (details deferred to Lemma 1.8 in S1.3.5).

Now, patching the identities inside and outside the volume V, we obtain

$$(1+\chi)\frac{1}{(2\pi)^3}{-\!\!\!\!\!\!\iiint}_{\mathbb{R}^3}\tilde{\boldsymbol{E}}(\boldsymbol{q})e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} + \chi\frac{1}{(2\pi)^3}{-\!\!\!\!\!\!\iiint}_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} = \boldsymbol{0}, \quad \forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V_0\cup\partial V\cup V_0').$$

Recalling the condition $\iiint_{\partial V} \mathrm{d}^3\boldsymbol{r} = 0$, we have $\iiint_{V_0\cup\partial V\cup V_0} \mathrm{d}^3\boldsymbol{r} = 0$ as a consequence. Therefore, by performing Fourier transform on an identity that holds in $\boldsymbol{r}$-space with the possible exception of a subset $V_0 \cup \partial V \cup V_0$ with zero volume, we may deduce

$$(1+\chi)\tilde{\boldsymbol{E}}(\boldsymbol{q}) + \chi\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2} = \boldsymbol{0}, \quad \forall\boldsymbol{q}\in\mathbb{R}^3\setminus V_0'' \tag{1.21}$$

where $\iiint_{V_0''} \mathrm{d}^3\boldsymbol{q} = 0$. Now, after multiplying Eq. 1.21 by $\boldsymbol{q}\times$, we obtain

$$(1+\chi)\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q}) - \chi|\boldsymbol{q}|^2\frac{\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-k^2} = \boldsymbol{0}, \quad \forall\boldsymbol{q}\in\mathbb{R}^3\setminus V_0'',$$

which implies $\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q}) = \boldsymbol{0}, \forall\boldsymbol{q}\in\mathbb{R}^3\setminus(V_0''\cup\partial O(\boldsymbol{0},|\sqrt{1+\chi}|k))$. Referring back to Eq. 1.21, and using the condition that $\chi\neq-1$, we then arrive at

$$\tilde{\boldsymbol{E}}(\boldsymbol{q}) = \boldsymbol{0}, \quad \forall\boldsymbol{q}\in\mathbb{R}^3\setminus(V_0''\cup\partial O(\boldsymbol{0},|\sqrt{1+\chi}|k))$$

which implies $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = (2\pi)^{-3/2}\|\tilde{\boldsymbol{E}}\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = 0$, just as claimed. $\blacksquare$

Remark The uniqueness theorem in the $\chi=-1$ case is much more complicated, yet still has important implications. This exceptional situation will be handled by Proposition 1.5, as well as Propositions 2.3, 2.8 and 2.4. $\blacksquare$

S1.2. Existence Theorem for Electromagnetic Scattering

In S1.1, we established the uniqueness theorem that the solution to the homogeneous linear equation $\|\hat{\mathcal{B}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$ is trivial under certain geometric and physical conditions. Now, in S1.2, we will rely on this information to conclude that the solution to the Born equation $\hat{\mathcal{B}}\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ must exist for any square-integrable incident field $\boldsymbol{E}_{\text{inc}}\in L^2(V;\mathbb{C}^3)$, when certain geometric and physical requirements are fulfilled.

Had the Born operator $\hat{\mathcal{B}}$ been a finite-dimensional square matrix, the uniqueness theorem would imply the existence theorem immediately. Now that $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ is a linear operator acting on an infinite-dimensional Hilbert space, the relation between uniqueness and existence could become subtle. In S1.2.1, we will point out that in most cases, the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ is a special type of Fredholm operator that indeed has similar behavior as that of a finite-dimensional square matrix. With this information, we can proceed with proving the existence theorem of the Born equation in S1.2.2. Later, in S1.2.3, we will also show that the solution to the Born equation, whenever existent, is stable against small perturbations in experimental parameters (refractive index and shape of the dielectric, wavelength of the light, etc.).

S1.2.1. Fredholm Decomposition of the Born operator

Recall that in Proposition 1.2, we decomposed the Born operator into $\hat{\mathcal{B}} = (1+\chi)\hat{I} - \chi\hat{\mathcal{D}} - \chi\hat{\gamma} : L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$, where the "dipole radiation" operator $\hat{\mathcal{D}} : L^2(V;\mathbb{C}^3)\longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ was defined by

$$(\hat{\mathcal{D}}\boldsymbol{E})(\boldsymbol{r}) := -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}, \quad \text{with } \tilde{\boldsymbol{E}}(\boldsymbol{q}) = \iiint_V \boldsymbol{E}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r}, \forall\boldsymbol{q}\in\mathbb{R}^3.$$

From this, we naturally have $\hat{I}-\hat{\mathcal{D}} : L^2(V;\mathbb{C}^3)\longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ as

$$((\hat{I}-\hat{\mathcal{D}})\boldsymbol{E})(\boldsymbol{r}) = \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}[\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}.$$

Now, we will study the operator $\hat{I}-\hat{\mathcal{D}}$ in further detail.

Lemma 1.1 *For every $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, we have the following relation:*

$$0\le\langle(\hat{I}-\hat{\mathcal{D}})\boldsymbol{E},\boldsymbol{E}\rangle_V\le\langle\boldsymbol{E},\boldsymbol{E}\rangle_V.$$

For $\chi\notin(-\infty,-1]$, the operator $[(1+\chi)\hat I-\chi\hat{\mathcal D}]^{-1}:L^2(V;\mathbb C^3)\longrightarrow L^2(V;\mathbb C^3)$ is well-defined, with an estimate for the operator norm as:

$$\|[(1+\chi)\hat I-\chi\hat{\mathcal D}]^{-1}\|_{L^2(V;\mathbb C^3)}=\frac{1}{|\chi|}\|[\chi^{-1}\hat I+(\hat I-\hat{\mathcal D})]^{-1}\|_{L^2(V;\mathbb C^3)}\le\frac{1}{|\chi|\operatorname{dist}(-\chi^{-1},[0,1])}.$$

Here, the distance from a complex number $z\in\mathbb C$ to a compact set $W\subset\mathbb C$ is given by $\operatorname{dist}(z,W):=\min_{w\in W}|z-w|$.

Proof Using the Parseval-Plancherel identity (Eq. 1.8), we may compute that

$$\langle(\hat I-\hat{\mathcal D})\boldsymbol E,\boldsymbol E\rangle_V=\frac{1}{(2\pi)^3}\iiint_{\mathbb R^3}\frac{|\boldsymbol q\cdot\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q,$$

which immediately leads to the inequality

$$0\le\langle(\hat I-\hat{\mathcal D})\boldsymbol E,\boldsymbol E\rangle_V=\frac{1}{(2\pi)^3}\iiint_{\mathbb R^3}\frac{|\boldsymbol q\cdot\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q\le\frac{1}{(2\pi)^3}\iiint_{\mathbb R^3}|\tilde{\boldsymbol E}(\boldsymbol q)|^2\,\mathrm d^3\boldsymbol q=\langle\boldsymbol E,\boldsymbol E\rangle.$$

Therefore, we see that $\hat I-\hat{\mathcal D}:L^2(V;\mathbb C^3)\longrightarrow L^2(\mathbb R^3;\mathbb C^3)$ is a positive-semidefinite operator with operator norm $\|\hat I-\hat{\mathcal D}\|_{L^2(\mathbb R^3;\mathbb C^3)}\le1$, giving rise to an estimate of its spectrum as $\sigma(\hat I-\hat{\mathcal D})\subset[0,1]$. Here, for an arbitrary bounded linear operator $\hat{\mathcal A}:\mathfrak H\longrightarrow\mathfrak H$ acting on a Hilbert space $\mathfrak H$, its spectrum $\sigma(\hat{\mathcal A})$ (see Ref. [40], Chap. 8) is a compact set specified by

$$\sigma(\hat{\mathcal A}):=\{\lambda\in\mathbb C|(\lambda\hat I-\hat{\mathcal A}):\mathfrak H\longrightarrow\mathfrak H\text{ has no bounded inverse}\}.$$

Now, if $\chi\notin(-\infty,-1]$, then $-\chi^{-1}\notin\sigma(\hat I-\hat{\mathcal D})$, and the inverse of $\chi^{-1}\hat I+(\hat I-\hat{\mathcal D})$ must be accordingly a bounded linear operator. We know that if the bounded linear mapping $\hat{\mathcal A}:\mathfrak H\longrightarrow\mathfrak H$ is also a Hermitian operator (satisfying $\langle\hat{\mathcal A}\boldsymbol E,\boldsymbol E\rangle\in\mathbb R,\forall\boldsymbol E\in\mathfrak H$), then for every $\lambda\notin\sigma(\hat{\mathcal A})$, the identity $\|(\lambda\hat I-\hat{\mathcal A})^{-1}\|_{\mathfrak H}=[\operatorname{dist}(\lambda,\sigma(\hat{\mathcal A}))]^{-1}$ holds (see Ref. [40], p. 149, also Ref. [41], p. 272). In the current context, we have $\operatorname{dist}(-\chi^{-1},\sigma(\hat I-\hat{\mathcal D}))\ge\operatorname{dist}(-\chi^{-1},[0,1])$, thus leading to the estimate

$$\|[\chi^{-1}\hat I+(\hat I-\hat{\mathcal D})]^{-1}\|_{L^2(V;\mathbb C^3)}=\frac{1}{\operatorname{dist}(-\chi^{-1},\sigma(\hat I-\hat{\mathcal D}))}\le\frac{1}{\operatorname{dist}(-\chi^{-1},[0,1])}.\quad\blacksquare$$

To bridge the gap between the uniqueness theorem and the existence theorem, we need to consider Fredholm operators (see Ref. [42], Chap. XI) on Hilbert spaces, which can be deemed as "well-behaved matrices of infinite dimension".

Definition Formally, a bounded linear operator $\hat{\mathcal O}:\mathfrak H\longrightarrow\mathfrak H$ acting on a Hilbert space is called a Fredholm operator if it satisfies three criteria:

(1) The range $\operatorname{ran}(\hat{\mathcal O}):=\{\hat{\mathcal O}\boldsymbol F|\boldsymbol F\in\mathfrak H\}$ is a closed (with respect to the L^2-limit on $\mathfrak H$) subspace of $\mathfrak H$, satisfying $\operatorname{ran}(\hat{\mathcal O})=\operatorname{Cl}(\operatorname{ran}(\hat{\mathcal O}))$;[19]

(2) The kernel space $\ker(\hat{\mathcal O}):=\{\boldsymbol F\in\mathfrak H|\hat{\mathcal O}\boldsymbol F=\boldsymbol 0\}$ is a finite-dimensional linear space $\dim\ker(\hat{\mathcal O})<+\infty$;

(3) The orthogonal complement of the range $[\operatorname{ran}(\hat{\mathcal O})]^\perp:=\{\boldsymbol G\in\mathfrak H|\langle\boldsymbol G,\boldsymbol F\rangle_{\mathfrak H}=0,\forall\boldsymbol F\in\hat{\mathcal O}\mathfrak H\}$ is a finite-dimensional linear space $\dim[\operatorname{ran}(\hat{\mathcal O})]^\perp<+\infty$.

For a Fredholm operator $\hat{\mathcal O}:\mathfrak H\longrightarrow\mathfrak H$, its index is given by $\operatorname{ind}(\hat{\mathcal O})=\dim\ker(\hat{\mathcal O})-\dim[\operatorname{ran}(\hat{\mathcal O})]^\perp$. ∎

Proposition 1.3 *If $\chi\notin(-\infty,-1]$, the Born operator $\hat{\mathcal B}=(1+\chi)\hat I-\chi\hat{\mathcal D}-\chi\hat\gamma:L^2(V;\mathbb C^3)\longrightarrow L^2(V;\mathbb C^3)$ is a Fredholm operator with index zero* $\operatorname{ind}(\hat{\mathcal B})=0$.

---

[19] Here, we use "Cl" to denote the topological closure in the Hilbert space $\mathfrak H$ with respect to the L^2-limit, the same notation as the topological closure with respect to the usual limit in $\mathbb R^3$ as defined in S1.0. The exact meaning of "Cl" is contextually clear.

Proof According to Lemma 1.1, when $\chi \notin (-\infty,-1]$, the operator $(1+\chi)\hat{I}-\chi\hat{\mathcal{D}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is invertible, with $\ker[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}] = \{\mathbf{0}\}$ and $\operatorname{ran}[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}] = L^2(V;\mathbb{C}^3)$. Thus, we have $\operatorname{ind}[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}] = 0$.

Meanwhile, the square-integrability $\gamma[k;V] < +\infty$ (as indicated in Eq. 1.5) of the convolution kernel $\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')$ qualifies $\hat{\gamma} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ as a compact operator that transforms bounded subsets of $L^2(V;\mathbb{C}^3)$ to relatively compact subsets of $L^2(V;\mathbb{C}^3)$.

Thus, for $\chi \notin (-\infty,-1]$, we have established the Born operator as a zero-index Fredholm operator perturbed by a compact operator. It is a fundamental result (see Ref. [27], p. 508 or Ref. [42], p. 366) that compact perturbation turns a Fredholm operator into a Fredholm operator with the index intact. Hence, we have $\operatorname{ind}(\hat{\mathcal{B}}) = \operatorname{ind}[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}-\chi\hat{\gamma}] = \operatorname{ind}[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}] = 0$, whenever $\chi \notin (-\infty,-1]$. ▮

Remark Unlike $\hat{\gamma} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$, which is a compact operator so long as the volume $V \Subset \mathbb{R}^3$ is open and bounded, the linear transformation $\hat{I}-\hat{\mathcal{D}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is not compact, even for spherical geometry $V = O(\mathbf{0},R), 0 < R < +\infty$. To prove this, we note that $\boldsymbol{F}_{\ell m}(\boldsymbol{r}) = \boldsymbol{F}_{\ell m}(|\boldsymbol{r}|,\theta,\phi) = \nabla[|\boldsymbol{r}|^\ell Y_{\ell m}(\theta,\phi)]$ satisfies the the following eigenequation

$$\begin{aligned}((\hat{I}-\hat{\mathcal{D}})\boldsymbol{F}_{\ell m})(\boldsymbol{r}) &= \nabla \oiint_{\partial O(\mathbf{0},R)} \frac{\boldsymbol{n}'\cdot\boldsymbol{F}_{\ell m}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,S' \\ &= \nabla \sum_{\ell'=0}^{\infty}\sum_{m'=-\ell'}^{\ell'} \oiint_{\partial O(\mathbf{0},R)} \frac{\boldsymbol{n}'\cdot\boldsymbol{F}_{\ell m}(\boldsymbol{r}')|\boldsymbol{r}|^\ell Y^*_{\ell'm'}(\theta',\phi')Y_{\ell'm'}(\theta,\phi)}{(2\ell+1)|\boldsymbol{r}'|^{\ell+1}}\,\mathrm{d}\,S' = \frac{\ell\boldsymbol{F}_{\ell m}(\boldsymbol{r})}{2\ell+1},\end{aligned}$$

thus providing a sequence of eigenvalues $\{\ell/(2\ell+1)|\ell = 1,2,\dots\}$ that accumulates at the point $\{1/2\}$. Meanwhile, the Riesz-Schauder theory of compact operators (see Ref. [43] or Ref. [44], pp. 283-284) dictates that the eigenvalues of a compact operator may only accumulate at the point $\{0\}$. In the light of this, for bounded and open volume $V \Subset \mathbb{R}^3$, the operator $\hat{I}-\hat{\mathcal{D}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is not (necessarily) compact, neither is the operator $\hat{I}-\hat{\mathcal{B}} = \chi(\hat{\mathcal{D}}-\hat{I}+\hat{\gamma}) : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$.

Thus, as pointed out in Ref. [16], we usually cannot reduce the Born equation to the prototypic Fredholm integral equation $(\hat{I}+\hat{\mathcal{K}})\boldsymbol{F} = \boldsymbol{G}$ where $\hat{\mathcal{K}}$ is a compact operator, and the theory for prototypic Fredholm integral equations are not directly applicable. Nonetheless, we have seen that the non-compact operator $\hat{I}-\hat{\mathcal{B}} = \chi(\hat{\mathcal{D}}-\hat{I}+\hat{\gamma})$ does not hamper further analytic understanding: the knowledge of the spectrum $\sigma(\hat{I}-\hat{\mathcal{D}}) \subset [0,1]$ makes it possible to use the Riesz-Fredholm theory for a detailed spectral analysis of the Born equation. ▮

S1.2.2. The Existence Theorem for Light Scattering

Now, we may consider the Born equation $\hat{\mathcal{B}}\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ for a square-integrable incident field $\boldsymbol{E}_{\text{inc}} \in L^2(V;\mathbb{C}^3)$, and assume that the dielectric susceptibility does not vanish $\chi \neq 0$. We regard the solution to the Born equation as existent and well-behaved if the number $1/\chi$ belongs to the resolvent set $\rho(\hat{\mathcal{G}}) = \mathbb{C}\setminus\sigma(\hat{\mathcal{G}})$ of the bounded linear operator $\hat{\mathcal{G}}$. Here, the resolvent set has two equivalent definitions (see Ref. [40], Chap. 8, or Ref. [44], Chap. VIII)

$$\rho(\hat{\mathcal{G}}) := \{\lambda \in \mathbb{C}|(\lambda\hat{I}-\hat{\mathcal{G}}) : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3) \text{ has a continuous inverse}\},$$

and

$$\rho(\hat{\mathcal{G}}) := \{\lambda \in \mathbb{C}|\ker(\lambda\hat{I}-\hat{\mathcal{G}}) = \{\mathbf{0}\}, \operatorname{ran}(\lambda\hat{I}-\hat{\mathcal{G}}) = L^2(V;\mathbb{C}^3)\},$$

where $\ker(\lambda\hat{I}-\hat{\mathcal{G}})$ also represents the totality of $\boldsymbol{E}$ that solves the eigenequation $\hat{\mathcal{G}}\boldsymbol{E} = \lambda\boldsymbol{E}$.

According to the spectral theory of bounded linear operators [40, 44], the spectrum $\sigma(\hat{\mathcal{G}}) = \mathbb{C}\setminus\rho(\hat{\mathcal{G}})$ can be decomposed into three disjoint subsets $\sigma(\hat{\mathcal{G}}) = \sigma_p(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})\cup\sigma_r(\hat{\mathcal{G}})$ as follows:

(i) The point spectrum (i.e. the set of eigenvalues)

$$\sigma_p(\hat{\mathcal{G}}) := \{\lambda \in \mathbb{C}|\ker(\lambda\hat{I}-\hat{\mathcal{G}}) \neq \{\mathbf{0}\}\};$$

(ii) The continuous spectrum[20]

$$\sigma_c(\hat{\mathcal{G}}) := \{\lambda \in \mathbb{C} |\, \ker(\lambda\hat{I} - \hat{\mathcal{G}}) = \{\mathbf{0}\}, \operatorname{ran}(\lambda\hat{I} - \hat{\mathcal{G}}) \neq L^2(V;\mathbb{C}^3), \operatorname{Cl}(\operatorname{ran}(\lambda\hat{I} - \hat{\mathcal{G}})) = L^2(V;\mathbb{C}^3)\};$$

(iii) The residual spectrum

$$\sigma_r(\hat{\mathcal{G}}) := \{\lambda \in \mathbb{C} |\, \ker(\lambda\hat{I} - \hat{\mathcal{G}}) = \{\mathbf{0}\}, \operatorname{Cl}(\operatorname{ran}(\lambda\hat{I} - \hat{\mathcal{G}})) \neq L^2(V;\mathbb{C}^3)\}.$$

Now, we will state the existence theorem that outlines the range of the resolvent set $\rho(\hat{\mathcal{G}})$, and provide bound estimates for $\|\hat{\mathcal{B}}^{-1}\|_{L^2(V;\mathbb{C}^3)} = \|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)}, 1/\chi \in \rho(\hat{\mathcal{G}})$, as far as we can. Here, the upper bounds of the operator norms for $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)}$ help provide an estimate of the condition number $\kappa(\hat{I} - \chi\hat{\mathcal{G}}) := \|\hat{I} - \chi\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)}$, the latter of which is useful for the error analysis in the finite element approach to light scattering [16].

Theorem 1.3 (Existence Theorem for Light Scattering) *For the Born equation* $\hat{\mathcal{B}}\boldsymbol{E} = (\hat{I} - \chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$, *where the dielectric susceptibility does not vanish* $\chi \neq 0$, *the solution exists in the following sense:*

(A) If the dielectric medium is dissipative with $\operatorname{Im}\chi < 0$, *then* $1/\chi \in \rho(\hat{\mathcal{G}})$, *and* $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq -|\chi|/\operatorname{Im}\chi$*;*

(B) If the dielectric medium is non-dissipative with $\chi > -1$, *its boundary* ∂V *occupies zero volume* $\iiint_{\partial V} \mathrm{d}^3\boldsymbol{r} = 0$, *and* $\mathbb{R}^3 \setminus (V \cup \partial V)$ *is connected, then* $1/\chi \in \rho(\hat{\mathcal{G}})$*;*

(C) If the dielectric medium satisfies $|\chi| < 1/\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}$, *then* $1/\chi \in \rho(\hat{\mathcal{G}})$ *and* $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq (1 - |\chi|\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)})^{-1}$*;*

(D) If the dielectric medium satisfies $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} < |1 + \chi^{-1}| - 1$, *then* $1/\chi \in \rho(\hat{\mathcal{G}})$ *and* $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq (|1 + \chi| - |\chi| - |\chi|\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)})^{-1}$*;*

(E) If the dielectric medium satisfies $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} < \operatorname{dist}(-\chi^{-1}, [0,1])$, *then* $1/\chi \in \rho(\hat{\mathcal{G}})$ *and* $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq |\chi|^{-1}[\operatorname{dist}(-\chi^{-1}, [0,1]) - \|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}]^{-1}$.

Proof First of all, we point out that a direct consequence of Proposition 1.3 is the set inclusion relation $\mathbb{C} \setminus [-1,0] \subset \rho(\hat{\mathcal{G}}) \cup \sigma_p(\hat{\mathcal{G}})$. Here, if $1/\chi \in \mathbb{C} \setminus [-1,0]$, then $\hat{\mathcal{B}} = \hat{I} - \chi\hat{\mathcal{G}}$ is a Fredholm operator with index zero $\operatorname{ind}(\hat{\mathcal{B}}) = \operatorname{ind}(\hat{I} - \chi\hat{\mathcal{G}}) = 0$. On one hand, if we have $\dim\ker(\hat{I} - \chi\hat{\mathcal{G}}) = 0$ (i.e. if the uniqueness theorem holds), then we may deduce $\operatorname{ran}(\hat{I} - \chi\hat{\mathcal{G}}) = L^2(V;\mathbb{C}^3)$ from the index equation $\dim[\operatorname{ran}(\hat{I} - \chi\hat{\mathcal{G}})]^{\perp} = \dim\ker(\hat{I} - \chi\hat{\mathcal{G}}) - \operatorname{ind}(\hat{I} - \chi\hat{\mathcal{G}}) = 0$ and the fact that the range $\operatorname{ran}(\hat{I} - \chi\hat{\mathcal{G}})$ of the Fredholm operator $\hat{I} - \chi\hat{\mathcal{G}}$ is closed. In other words, when $1/\chi \in \mathbb{C} \setminus [-1,0]$, the uniqueness theorem $\dim\ker(\hat{I} - \chi\hat{\mathcal{G}}) = 0$ implies the existence theorem $1/\chi \in \rho(\hat{\mathcal{G}})$. On the other hand, if we have $0 < \dim\ker(\hat{I} - \chi\hat{\mathcal{G}}) < +\infty$ instead, then obviously $1/\chi \in \sigma_p(\hat{\mathcal{G}})$ is an eigenvalue of the operator $\hat{\mathcal{G}}$ that has finite multiplicity, i.e. the eigenvalue has a finite-dimensional eigenspace subordinate to it.

In situation (A), we quote from the uniqueness theorem (Theorem 1.2) that $\dim\ker(\hat{I} - \chi\hat{\mathcal{G}}) = 0$ for whatever bounded and open volume V with $\operatorname{Im}\chi < 0$. Therefore, we have $1/\chi \in \rho(\hat{\mathcal{G}})$ for $\operatorname{Im}\chi < 0$. Accordingly, the coercivity bound (Eq. 1.20) leads us to $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq -|\chi|/\operatorname{Im}\chi$. Furthermore, according to Corollaries 1.2 and 1.3 in S1.3.4, we may conclude that if χ satisfies the relation

$$|\chi|^2 \min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\} < \operatorname{Im}\chi,$$

then we have $1/\chi \in \rho(\hat{\mathcal{G}})$ as well as the norm estimate

$$\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq \frac{|\chi|}{\operatorname{Im}\chi}\left(1 - \frac{|\chi|^2}{\operatorname{Im}\chi}\min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\}\right)^{-1};$$

---

[20] Here, the "point spectrum" σ_p and "continuous spectrum" σ_c do not necessarily correspond to any geometric "discreteness" and "continuity" in the language of physical/chemical spectroscopy. The point spectrum σ_p of a bounded operator may form a geometric continuum (see Ref. [45], p. 194 or Ref. [46], p. 83), whereas the continuous spectrum σ_c of a bounded operator may well be isolated points.

if χ satisfies the inequality

$$|\chi|^2\left(1+\min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3},3\sqrt{2}kR_V\right\}\right)<|\operatorname{Re}\chi|,$$

then we have $1/\chi\in\rho(\hat{\mathcal{G}})$ as well as the norm estimate

$$\|(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)}\leq\frac{|\chi|}{|\operatorname{Re}\chi|}\left[1-\frac{|\chi|^2}{|\operatorname{Re}\chi|}\left(1+\min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3},3\sqrt{2}kR_V\right\}\right)\right]^{-1}.$$

In situation (B), we quote from the uniqueness theorem (Theorem 1.2) that $\dim\ker(\hat{I}-\chi\hat{\mathcal{G}})=0$ if the exterior volume $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected and the boundary occupies zero volume $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$, and thus we have $1/\chi\in\rho(\hat{\mathcal{G}})$ according to the theory of Fredholm operators.

In situation (C), the inverse of the Born operator

$$\hat{\mathcal{B}}^{-1}=(\hat{I}-\chi\hat{\mathcal{G}})^{-1}=\sum_{s=0}^{\infty}(\chi\hat{\mathcal{G}})^s=\hat{I}+\sum_{s=1}^{\infty}(\chi\hat{\mathcal{G}})^s$$

is a well-defined linear operator with finite operator norm $\|\hat{\mathcal{B}}^{-1}\|_{L^2(V;\mathbb{C}^3)}\leq 1+\sum_{s=1}^{\infty}\|(\chi\hat{\mathcal{G}})^s\|_{L^2(V;\mathbb{C}^3)}\leq(1-|\chi|\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)})^{-1}$. We have $1/\chi\in\rho(\hat{\mathcal{G}})$ without further topological constraints on the shape V. This shows that for optically soft particles satisfying $|\chi|<1/\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}$, the light scattering problem is always well-conditioned. Here, the relations in Eqs. 1.11 and 1.12 can provide a practical estimate for the range of $|\chi|$ fulfilling the optical softness criterion. Meanwhile, this also gives an error estimate for the first-order[21] Born approximation $\boldsymbol{E}\approx\boldsymbol{E}_{\text{inc}}$ as

$$\|\boldsymbol{E}-\boldsymbol{E}_{\text{inc}}\|_{L^2(V;\mathbb{C}^3)}\leq\frac{|\chi|\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{E}_{\text{inc}}\|_{L^2(V;\mathbb{C}^3)}}{1-|\chi|\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}}.$$

More generally, we may truncate the Born series $\hat{\mathcal{B}}^{-1}=\hat{I}+\sum_{s=1}^{\infty}(\chi\hat{\mathcal{G}})^s$ at the m-th power to obtain the Born approximation at $(m+1)$-st order, with an error estimate

$$\left\|\boldsymbol{E}-\left[\hat{I}+\sum_{s=1}^{m}(\chi\hat{\mathcal{G}})^s\right]\boldsymbol{E}_{\text{inc}}\right\|_{L^2(V;\mathbb{C}^3)}\leq\frac{|\chi|^m\|\hat{\mathcal{G}}\|^m_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{E}_{\text{inc}}\|_{L^2(V;\mathbb{C}^3)}}{1-|\chi|\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}}.$$

In situation (D), we may deduce from $|1+\chi|-|\chi|>|\chi|\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}\geq 0$ and $\|\hat{\mathcal{D}}\|_{L^2(V;\mathbb{C}^3)}\leq 1$ the following estimate

$$\left\|\left(\hat{I}-\frac{\chi}{1+\chi}\hat{\mathcal{D}}\right)^{-1}\right\|_{L^2(V;\mathbb{C}^3)}\leq\left(1-\left|\frac{\chi}{1+\chi}\right|\right)^{-1},$$

and

$$\left(\hat{I}-\frac{\chi}{1+\chi}\hat{\mathcal{D}}\right)^{-1}\frac{\hat{\mathcal{B}}}{1+\chi}=\hat{I}-\left(\hat{I}-\frac{\chi}{1+\chi}\hat{\mathcal{D}}\right)^{-1}\frac{\chi}{1+\chi}\hat{\gamma}.$$

Therefore, we have

$$\begin{aligned}\|\hat{\mathcal{B}}^{-1}\|_{L^2(V;\mathbb{C}^3)}&\leq\frac{1}{|1+\chi|}\left\|\left(\hat{I}-\frac{\chi}{1+\chi}\hat{\mathcal{D}}\right)^{-1}\right\|_{L^2(V;\mathbb{C}^3)}\frac{1}{1-\left\|\left(\hat{I}-\frac{\chi}{1+\chi}\hat{\mathcal{D}}\right)^{-1}\right\|_{L^2(V;\mathbb{C}^3)}\left|\frac{\chi}{1+\chi}\right|\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}}\\&\leq\frac{\left(1-\left|\frac{\chi}{1+\chi}\right|\right)^{-1}}{|1+\chi|\left[1-\left(1-\left|\frac{\chi}{1+\chi}\right|\right)^{-1}\left|\frac{\chi}{1+\chi}\right|\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}\right]}=\frac{1}{|1+\chi|-|\chi|-|\chi|\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}}.\end{aligned}$$

---

[21] The nomenclature for the order of Born approximation is consistent with the convention in quantum scattering problems. (See Ref. [47]). We also note that the convergence condition for Born series has been described as "difficult to establish" in Ref. [10], p. 710. Our effort here has been devoted to providing a sufficient condition for the convergence of Born series with a computable bound on χ.

Thus, for fixed χ, so long as the size of the particle is sufficiently small, the condition $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} < |1+\chi^{-1}|-1$ can always be satisfied, and the light scattering problem becomes well-conditioned irrespective of the topology of V.

In situation (E), we note that when $\|[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}]^{-1}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} < 1$, we may invert

$$\hat{I}-\chi\hat{\mathcal{G}} = [(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}]\{\hat{I}-[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}]^{-1}\chi\hat{\gamma}\}$$

to obtain the estimate

$$\|(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq \frac{\|[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}]^{-1}\|_{L^2(V;\mathbb{C}^3)}}{1-\|[(1+\chi)\hat{I}-\chi\hat{\mathcal{D}}]^{-1}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}} \leq \frac{1}{|\chi|[\operatorname{dist}(-\chi^{-1},[0,1])-\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}]}.$$

In sum, we see that for any bounded and open volume V, the Born equation $\hat{\boldsymbol{\mathcal{B}}}\boldsymbol{E} = \boldsymbol{E}_{\text{inc}} \in L^2(V;\mathbb{C}^3)$ is well-conditioned at least for the following range of susceptibilities:

- $\operatorname{Im}\chi < 0$; or
- $\|\hat{\mathcal{G}}\|_{L^2(V;\mathbb{C}^3)}|\chi| < 1$; or
- $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} < \max\left\{\left|1+\frac{1}{\chi}\right|-1, \operatorname{dist}\left(-\frac{1}{\chi},[0,1]\right)\right\}$; or
- $\operatorname{Im}\chi > |\chi|^2 \min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\}$; or
- $|\operatorname{Re}\chi| > |\chi|^2\left(1+\min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, 3\sqrt{2}kR_V\right\}\right)$.

If we further have $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r} = 0$ and $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected, then the solvable range also includes $\chi \in (-1,+\infty)$. ▮

The properties of Fredholm operators ensure that $\mathbb{C}\setminus[-1,0]\subset\rho(\hat{\mathcal{G}})\cup\sigma_p(\hat{\mathcal{G}})$, it would then be natural to ask the status of a number $\lambda\in[-1,0]$ in the decomposition $\mathbb{C}=\sigma(\hat{\mathcal{G}})\cup\rho(\hat{\mathcal{G}})=\sigma_p(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})\cup\sigma_r(\hat{\mathcal{G}})\cup\rho(\hat{\mathcal{G}})$.

Proposition 1.4 (Absence of Residual Spectrum) *So long as the volume V is bounded and open, the residual spectrum of $\hat{\mathcal{G}}$ is empty, i.e. $\sigma_r(\hat{\mathcal{G}})=\emptyset$.*

Proof We have the following transpose identity

$$\begin{aligned}
&\langle\boldsymbol{F},(\lambda\hat{I}-\hat{\mathcal{G}})\boldsymbol{E}\rangle_V\\
={}&\langle\boldsymbol{F},\lambda\boldsymbol{E}\rangle_V+\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{[\boldsymbol{q}\cdot\tilde{\boldsymbol{F}}^*(\boldsymbol{q})][\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}\,\mathrm{d}^3\boldsymbol{q}-\iiint_V\boldsymbol{F}^*(\boldsymbol{r})\cdot\left[\iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'\right]\mathrm{d}^3\boldsymbol{r}\\
={}&\langle\boldsymbol{E}^*,\lambda\boldsymbol{F}^*\rangle_V+\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{[\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})][\boldsymbol{q}\cdot\tilde{\boldsymbol{F}}^*(\boldsymbol{q})]}{|\boldsymbol{q}|^2}\,\mathrm{d}^3\boldsymbol{q}-\iiint_V\boldsymbol{E}(\boldsymbol{r}')\cdot\left[\iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{F}^*(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right]\mathrm{d}^3\boldsymbol{r}'\\
={}&\langle\boldsymbol{E}^*,(\lambda\hat{I}-\hat{\mathcal{G}})\boldsymbol{F}^*\rangle_V.
\end{aligned}\tag{1.22}$$

In the above derivation, we have exploited the reciprocal symmetry $\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')=\hat{\Gamma}(\boldsymbol{r}',\boldsymbol{r})$ and interchanged the integrals with respect to $\mathrm{d}^3\boldsymbol{r}$ and $\mathrm{d}^3\boldsymbol{r}'$. The interchange of the two volume integrals can be justified by the absolute integrability of the iterated integrals (which derives from the Hilbert-Schmidt bound in Eq. 1.5) and an application of the the Fubini theorem (see Ref. [31], p. 148 or Ref. [29], p. 25).

Now, suppose that the uniqueness theorem holds, i.e. $\ker(\lambda\hat{I}-\hat{\mathcal{G}})=\{\boldsymbol{0}\}$. We point out that with the transpose identity (Eq. 1.22), we may deduce that $\operatorname{Cl}(\operatorname{ran}(\lambda\hat{I}-\hat{\mathcal{G}}))=\operatorname{Cl}((\lambda\hat{I}-\hat{\mathcal{G}})L^2(V;\mathbb{C}^3))=L^2(V;\mathbb{C}^3)$, which shows that $\lambda\notin\sigma_r(\hat{\mathcal{G}})$. If the opposite is true, we will be able to find a non-zero member of $L^2(V;\mathbb{C}^3)$ that is orthogonal to all the elements of $(\lambda\hat{I}-\hat{\mathcal{G}})L^2(V;\mathbb{C}^3)$, which means that for some $\boldsymbol{F}\neq\boldsymbol{0}$, we have $\langle\boldsymbol{F},(\lambda\hat{I}-\hat{\mathcal{G}})\boldsymbol{E}\rangle=\langle\boldsymbol{E}^*,(\lambda\hat{I}-\hat{\mathcal{G}})\boldsymbol{F}^*\rangle=0,\forall\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$. This immediately entails $(\lambda\hat{I}-\hat{\mathcal{G}})\boldsymbol{F}^*=\boldsymbol{0}$. According to the assumption that $\ker(\lambda\hat{I}-\hat{\mathcal{G}})=\{\boldsymbol{0}\}$, we have $\boldsymbol{F}^*=\boldsymbol{0}$ as well. The contradiction means that we must have $\operatorname{Cl}(\operatorname{ran}(\lambda\hat{I}-\hat{\mathcal{G}}))=L^2(V;\mathbb{C}^3)$. ▮

Proposition 1.5 *So long as the volume V is bounded and open, we have $-1\in\sigma_p(\hat{\mathcal{G}})$ as an eigenvalue of infinite multiplicity* $\dim\ker(\hat{I}+\hat{\mathcal{G}})=+\infty$*, and $0\in\sigma_p(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})$ being outside the resolvent set $\rho(\hat{\mathcal{G}})$.*

Proof Take any $f \in C_0^\infty(V;\mathbb{C})$, then it follows that

$$((\hat{I}+\hat{\mathcal{G}})\nabla f)(\boldsymbol{r}) = \nabla\times\nabla\times\iiint_V \frac{\nabla' f(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\nabla\times\iiint_V \nabla'\times\left[\frac{\nabla' f(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\mathrm{d}^3\boldsymbol{r}' = \boldsymbol{0},$$

where one can use the Gauss theorem to convert the last volume integral into a surface integral with vanishing integrand. This proves that $\dim\ker(\hat{I}+\hat{\mathcal{G}}) = +\infty$ and $-1 \in \sigma_p(\hat{\mathcal{G}})$.

If we assume that $0 \in \rho(\hat{\mathcal{G}})$, then $\hat{\mathcal{D}} - \hat{I} = \hat{\mathcal{G}} - \hat{\gamma}$ will become a Fredholm operator, satisfying $\dim\ker(\hat{\mathcal{D}}-\hat{I}) < +\infty$, which contradicts the fact that

$$((\hat{\mathcal{D}}-\hat{I})\nabla\times\boldsymbol{F})(\boldsymbol{r}) = \nabla\left[\nabla\cdot\iiint_V \frac{\nabla'\times\boldsymbol{F}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right] = -\nabla\left[\iiint_V \nabla'\cdot\left(\frac{\nabla'\times\boldsymbol{F}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)\mathrm{d}^3\boldsymbol{r}'\right] = \boldsymbol{0},\quad \forall\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3),$$

i.e. $\dim\ker(\hat{\mathcal{D}}-\hat{I}) = +\infty$. As the residual spectrum is empty $\sigma_r(\hat{\mathcal{G}}) = \emptyset$, we then must have $0 \in \sigma_p(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})\cup\sigma_r(\hat{\mathcal{G}}) = \sigma_p(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})$. $\blacksquare$

Remark While the eigenvalue $-1 \in \sigma_p(\hat{\mathcal{G}})$ of infinite multiplicity $\dim\ker(\hat{I}+\hat{\mathcal{G}}) = +\infty$ seems to suggest that the light scattering problem becomes seriously ill-conditioned for $\chi = -1$, we will show, in Propositions 2.8 and 2.4, that such a concern about the case of $\chi = -1$ is physically irrelevant. $\blacksquare$

Proposition 1.6 *If $\mathbb{R}^3 \setminus (V \cup \partial V)$ is connected and $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r} = 0$, then $(-1,0] \subset \rho(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})$.*

Proof For $-1 < \lambda < 0$, the uniqueness theorem (Theorem 1.2, situation (B)) implies that λ is not an eigenvalue, so it must belong to the set $\rho(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})\cup\sigma_r(\hat{\mathcal{G}}) = \rho(\hat{\mathcal{G}})\cup\sigma_c(\hat{\mathcal{G}})$.

For $\lambda = 0$, we can still show that $\|\hat{\mathcal{G}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$ implies $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$, so $0 \notin \sigma_p(\hat{\mathcal{G}})$. The underlying rationale is that Eq. 1.18 in Theorem 1.1 ensures that the relation $\|\hat{\mathcal{G}}\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$ leads us to $\tilde{\boldsymbol{E}}(\boldsymbol{q}) = \boldsymbol{0}, \forall|\boldsymbol{q}| = k$. Accordingly, we may adapt the proof of the uniqueness theorem (Theorem 1.2, situation (B)) to show that both

$$\tilde{\boldsymbol{E}}(\boldsymbol{q}) + \frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-k^2} = \boldsymbol{0},\quad \forall\boldsymbol{q}\in\mathbb{R}^3\setminus V_0''$$

and

$$\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q}) - |\boldsymbol{q}|^2\frac{\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-k^2} = \boldsymbol{0},\quad \forall\boldsymbol{q}\in\mathbb{R}^3\setminus V_0''$$

hold, where $\iiint_{V_0''}\mathrm{d}^3\boldsymbol{q} = 0$. Therefore, we may proceed to conclude that $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = (2\pi)^{-3/2}\|\tilde{\boldsymbol{E}}\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = 0$. $\blacksquare$

S1.2.3. Stable Dependence of the Solution on Experimental Parameters

In Theorem 1.3 of S1.2.2, we specified the conditions that ensure the existence of a unique solution to the Born equation $\hat{\mathcal{B}}\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$, for a square-integrable input $\boldsymbol{E}_{\text{inc}} \in L^2(V;\mathbb{C}^3)$. In the meantime, we have already provided a proof of stable response of the solution with respect to small changes in the incident field $\boldsymbol{E}_{\text{inc}}$ (recall the definition of the resolvent set $\rho(\hat{\mathcal{G}}) := \{\lambda\in\mathbb{C}|(\hat{\mathcal{G}}-\lambda\hat{I})^{-1} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a continuous linear mapping$\}$). Now we will proceed to specify some further stability results for the solution to the Born equation.

Proposition 1.7 (Complex-Analytic Dependence on Susceptibility) *The solution $\boldsymbol{E}$ to the Born equation $(\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ is complex-analytic with respect to $\chi = \operatorname{Re}\chi + i\operatorname{Im}\chi$ for $\chi \in \{0\}\cup\{1/\lambda|\lambda\in\rho(\hat{\mathcal{G}})\}$.*

Proof In a neighborhood of a non-vanishing susceptibility $\chi_0 \in \{1/\lambda|\lambda\in\rho(\hat{\mathcal{G}})\}\setminus\{0\}$, the statement in the proposition is a direct consequence of the analytic properties of the resolvent $(\hat{\mathcal{G}}-\lambda\hat{I})^{-1}$ for $\lambda = 1/\chi \in \rho(\hat{\mathcal{G}})$. Explicitly, so long as $\chi_0 \neq 0$ and

$$\left|\frac{1}{\chi}-\frac{1}{\chi_0}\right|\left\|\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right\|_{L^2(V;\mathbb{C}^3)} < 1, \tag{1.23}$$

the following operator-valued power series

$$\begin{aligned}\frac{1}{\chi}\left(\frac{1}{\chi}\hat{I}-\hat{\mathcal{G}}\right)^{-1} &= \frac{1}{\chi}\left[\left(\frac{1}{\chi}-\frac{1}{\chi_0}\right)\hat{I}+\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)\right]^{-1}=\frac{1}{\chi}\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\left[\left(\frac{1}{\chi}-\frac{1}{\chi_0}\right)\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}+\hat{I}\right]^{-1}\\ &= \frac{1}{\chi}\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\sum_{m=0}^{\infty}\left[\left(\frac{1}{\chi_0}-\frac{1}{\chi}\right)\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right]^{m}\end{aligned}$$

is meaningful, because of the following bound estimate

$$\begin{aligned}\left\|\sum_{m=0}^{\infty}\left[\left(\frac{1}{\chi_0}-\frac{1}{\chi}\right)\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right]^{m}\right\|_{L^2(V;\mathbb{C}^3)} &\le \sum_{m=0}^{\infty}\left\|\left[\left(\frac{1}{\chi_0}-\frac{1}{\chi}\right)\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right]^{m}\right\|_{L^2(V;\mathbb{C}^3)}\\ &\le \sum_{m=0}^{\infty}\left\|\left[\left(\frac{1}{\chi_0}-\frac{1}{\chi}\right)\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right]\right\|^{m}_{L^2(V;\mathbb{C}^3)}=\frac{1}{1-\left|\frac{1}{\chi}-\frac{1}{\chi_0}\right|\left\|\left(\frac{1}{\chi_0}\hat{I}-\hat{\mathcal{G}}\right)^{-1}\right\|_{L^2(V}}\end{aligned}$$

Here, we have used the convention that $\hat{\mathcal{A}}^0=\hat{I}$ for an arbitrary operator $\hat{\mathcal{A}}$. From this power series expansion, we see that the solution $\boldsymbol{E}=(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}}$ is complex-analytic with respect to χ in a neighborhood of $\chi_0\neq 0$ specified by Eq. 1.23. In a neighborhood of $\chi_0=0$, the operator-valued Taylor expansion follows directly from the argument for situation (C) in Theorem 1.3. ▮

Proposition 1.8 (Continuous Dependence on Wave Length) *The solution* $\boldsymbol{E}$ *to the Born equation as defined in Eq. 1.13 is a continuous function with respect to the wave number* $k=2\pi/\lambda_{wave}\in(0,+\infty)$.

Proof As $|k-k'|\to 0$, the difference between the corresponding Born operators

$$(\hat{\mathcal{B}}_{[k']}\boldsymbol{E})(\boldsymbol{r})-(\hat{\mathcal{B}}_{[k]}\boldsymbol{E})(\boldsymbol{r}) = -\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')[e^{-ik'|\boldsymbol{r}-\boldsymbol{r}'|}-e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}-i(k-k')|\boldsymbol{r}-\boldsymbol{r}'|]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$$

satisfies $\|\hat{\mathcal{B}}_{[k']}-\hat{\mathcal{B}}_{[k]}\|_{L^2(V;\mathbb{C}^3)}=O(|k'-k|^2)\to 0$. We accordingly have the estimate $\|\hat{\mathcal{B}}^{-1}_{[k']}-\hat{\mathcal{B}}^{-1}_{[k]}\|_{L^2(V;\mathbb{C}^3)}\le\|\hat{\mathcal{B}}^{-1}_{[k']}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\mathcal{B}}_{[k']}-\hat{\mathcal{B}}_{[k]}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\mathcal{B}}^{-1}_{[k]}\|_{L^2(V;\mathbb{C}^3)}\to 0$ as $|k'-k|\to 0$. ▮

To discuss the shape-dependence of the Born equation, we will explicitly append a subscript to denote the volume under consideration. For instance, we will use $\hat{\mathcal{B}}_V\boldsymbol{E}=(\hat{I}-\chi\hat{\mathcal{G}}_V)\boldsymbol{E}=\boldsymbol{E}_{\text{inc}}$ to denote the Born equation

$$(1+\chi)\boldsymbol{E}(\boldsymbol{r})-\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'=\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}),\quad \boldsymbol{r}\in V.$$

Proposition 1.9 (Continuous Dependence on Shape Distortions) *Let* χ *be a fixed susceptibility satisfying* $\operatorname{Im}\chi<0$, *the volume* V *be a bounded open subset of* $\mathbb{R}^3$, $\{V_{(s)}\subset V|\iiint_{\partial V_{(s)}}\mathrm{d}^3\boldsymbol{r}=0,s=1,2,\dots\}$ *be a sequence of bounded open subsets satisfying* $\lim_{s\to\infty}\iiint_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\mathrm{d}^3\boldsymbol{r}=0$, *and* $\boldsymbol{E};\boldsymbol{E}_{(s)},s=1,2,\dots$ *be the solutions to the respective Born equations* $(\hat{\mathcal{B}}_V\boldsymbol{E})(\boldsymbol{r})=\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}),\boldsymbol{r}\in V$; $(\hat{\mathcal{B}}_{V_{(s)}}\boldsymbol{E}_{(s)})(\boldsymbol{r})=\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}),\boldsymbol{r}\in V_{(s)}\subset V,s=1,2,\dots$, *then we have the error estimate for the effects of shape distortion*

$$\|\boldsymbol{E}-\boldsymbol{E}_{(s)}\|_{L^2(V_{(s)};\mathbb{C}^3)} \le \|\hat{\mathcal{B}}^{-1}_{V_{(s)}}\|_{L^2(V_{(s)};\mathbb{C}^3)}|\chi|\|\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\boldsymbol{E}\|_{L^2(V_{(s)};\mathbb{C}^3)}\le-\frac{|\chi|^2}{\operatorname{Im}\chi}\|\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\boldsymbol{E}\|_{L^2(V_{(s)};\mathbb{C}^3)} \tag{1.24}$$

In particular, we have

$$\lim_{s\to\infty}\|\boldsymbol{E}-\boldsymbol{E}_{(s)}\|_{L^2(V_{(s)};\mathbb{C}^3)}=0. \tag{1.25}$$

Proof We may compare the two equations

$$\begin{aligned}(\hat{\mathcal{B}}_{V_{(s)}}\boldsymbol{E}_{(s)})(\boldsymbol{r})=\boldsymbol{E}_{(s)}(\boldsymbol{r})-\chi(\hat{\mathcal{G}}_{V_{(s)}}\boldsymbol{E}_{(s)})(\boldsymbol{r}) &= \boldsymbol{E}_{\text{inc}}(\boldsymbol{r}),\quad \boldsymbol{r}\in V_{(s)}\\ (\hat{\mathcal{B}}_{V_{(s)}}\boldsymbol{E})(\boldsymbol{r})=\boldsymbol{E}(\boldsymbol{r})-\chi(\hat{\mathcal{G}}_{V_{(s)}}\boldsymbol{E})(\boldsymbol{r}) &= [\boldsymbol{E}_{\text{inc}}(\boldsymbol{r})+\chi(\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\boldsymbol{E})(\boldsymbol{r})],\quad \boldsymbol{r}\in V_{(s)}\end{aligned}$$

to arrive at Eq. 1.24.

By restricting a given square-integrable vector field $\boldsymbol{E}=(\hat{I}-\chi\hat{\mathcal{G}}_V)^{-1}\boldsymbol{E}_{\text{inc}}\in L^2(V;\mathbb{C}^3)$ to a sequence of subsets $\{V_{(s)}\subset V|\iiint_{\partial V_{(s)}}\mathrm{d}^3\boldsymbol{r}=0,s=1,2,\dots\}$ satisfying $\lim_{s\to\infty}\iiint_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\mathrm{d}^3\boldsymbol{r}=0$, we may use the absolute integrability to deduce that $\lim_{s\to\infty}\|\boldsymbol{E}\|_{L^2(V\setminus(V_{(s)}\cup\partial V_{(s)});\mathbb{C}^3)}=0$. As the sequence of operators $\{\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}:L^2(V\setminus(V_{(s)}\cup\partial V_{(s)});\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)|s=1,2,\dots\}$ is uniformly bounded in operator norm: $\sup_{s\in\{1,2,\dots\}}\|\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\|_{L^2(V\setminus(V_{(s)}\cup\partial V_{(s)});\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)}<+\infty$, we may thus conclude that $\lim_{s\to\infty}\|\hat{\mathcal{G}}_{V\setminus(V_{(s)}\cup\partial V_{(s)})}\boldsymbol{E}\|_{L^2(V_{(s)};\mathbb{C}^3)}=0$, which confirms Eq. 1.25. ▮

Remark The continuous dependence of the scattering field on the shape distortions of the scattering medium has both experimental and theoretical implications.

Experimentally speaking, the shape of the dielectric scattering medium may not be known in exact detail, due to manufacturing imperfections, surface defects, thermodynamic fluctuations, among other factors. Here, Proposition 1.9 ensures that incomplete knowledge of the dielectric shape does not practically undermine the reliability of light scattering models based on idealized geometry.

Theoretically speaking, the effectiveness of an algorithm for the solution of light scattering typically hinges on the smoothness of the dielectric boundary. The stability result proved in Proposition 1.9 allows us to treat the problem of light scattering on certain shapes with non-smooth boundaries (say, polyhedral shaped dielectrics) as the limiting case for light scattering on dielectrics with smooth boundary surfaces, by a procedure of "shape approximation from within". ▮

S1.3. Additional Comments and Details

In this subsection, we supply detailed quantitative arguments for certain technical claims in the foregoing discussions. We gather these comments and details here, as they may require highly specialized techniques, may involve extensive computations, or may form a topical digression from our major interest in homogeneous dielectric media.

S1.3.1. Born Equation in the Discrete Dipole Approximation Method for Light Scattering

In S1.0.2, we defined the Born operator $\hat{\mathcal{B}}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ by interpreting the partial derivatives on the right-hand side of the following equation

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}):=(1+\chi)\boldsymbol{E}(\boldsymbol{r})-\chi\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}',\quad \boldsymbol{r}\in V$$

in the distributional sense.

In the literature concerning numerical solutions of the light scattering problem, the Born equation is usually formulated via the digitized Green function approach (also known as the discrete dipole approximation) as follows [15, 16, 32, 33, 34, 35, 36, 37]:

$$(\hat{\mathcal{B}}'\boldsymbol{E})(\boldsymbol{r}):=\boldsymbol{E}(\boldsymbol{r})-\chi\left[-\frac{\boldsymbol{E}(\boldsymbol{r})}{3}+k^2\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'+\lim_{\varepsilon\to0^+}\iiint_{V\setminus O(\boldsymbol{r},\varepsilon)}\nabla\nabla\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\cdot\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'\right]\tag{1.26}$$

where the Hessian matrix reads

$$\nabla\nabla\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}=\frac{k^2e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\left[-\left(\frac{i}{k|\boldsymbol{r}-\boldsymbol{r}'|}+\frac{1}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2}\right)\mathbb{1}+\left(-1+\frac{3i}{k|\boldsymbol{r}-\boldsymbol{r}'|}+\frac{3}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2}\right)\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right],$$

with $\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}=(\boldsymbol{r}-\boldsymbol{r}')(\boldsymbol{r}-\boldsymbol{r}')^{\mathrm{T}}/|\boldsymbol{r}-\boldsymbol{r}'|^2$ (cf. Eq. 1.4 in S1.0.2).

It can be verified that the above two definitions of the Born operator agree with each other when $\boldsymbol{E}\in C_0^\infty(V;\mathbb{C}^3)$, that is, $\hat{\mathcal{B}}=\hat{\mathcal{B}}':C_0^\infty(V;\mathbb{C}^3)\longrightarrow C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$. To show that these two definitions agree for all square-integrable inputs $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, that is, to show that $\hat{\mathcal{B}}=\hat{\mathcal{B}}':L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$, it would suffice to give a direct proof that $\hat{\mathcal{B}}':L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ defines a continuous linear mapping, as demonstrated below.

Proposition 1.10 *The linear mapping* $\hat{\mathcal{B}}' = \hat{I} - \chi(\hat{\mathcal{G}}_0 + \hat{\mathcal{G}}_1 + \hat{\mathcal{G}}_2 + \hat{\mathcal{G}}_3) : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ *is continuous, where*

$$\begin{aligned}
(\hat{\mathcal{G}}_0\boldsymbol{E})(\boldsymbol{r}) &:= -\frac{\boldsymbol{E}(\boldsymbol{r})}{3},\\
(\hat{\mathcal{G}}_1\boldsymbol{E})(\boldsymbol{r}) &:= \iiint_V \frac{k^2 e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\left(\overleftrightarrow{I} - \hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right)\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' = \iiint_V \hat{\Gamma}_1(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}',\\
(\hat{\mathcal{G}}_2\boldsymbol{E})(\boldsymbol{r}) &:= -i\iiint_V \frac{k e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|^2}\left(\overleftrightarrow{I} - 3\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right)\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' = \iiint_V \hat{\Gamma}_2(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}',\\
(\hat{\mathcal{G}}_3\boldsymbol{E})(\boldsymbol{r}) &:= -\lim_{\varepsilon\to0^+}\iiint_{V\setminus O(\boldsymbol{r},\varepsilon)} \frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|^3}\left(\overleftrightarrow{I} - 3\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right)\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' = \lim_{\varepsilon\to0^+}\iiint_{V\setminus O(\boldsymbol{r},\varepsilon)} \hat{\Gamma}_3(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'.
\end{aligned}$$

Proof We claim that there exist four finite numbers $G_m < +\infty, m = 0,1,2,3$ such that $\|\hat{\mathcal{G}}_m\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq G_m\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}, m = 0,1,2,3$ for all $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$. Once this can be shown, we can further arrive at $\|\hat{\mathcal{B}}'\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq [1 + |\chi|(G_0 + G_1 + G_2 + G_3)]\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}, \forall\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$.

Here, it is obvious that one can choose $G_0 = 1/3 < +\infty$.

To show that the constant $G_1 + G_2 < +\infty$, we need the Young inequality (see Ref. [25], p. 271) for convolutions $h(\boldsymbol{r}) = \iiint_{\mathbb{R}^3} f(\boldsymbol{r}')g(\boldsymbol{r}-\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'$:

$$\iiint_{\mathbb{R}^3} |h(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r} \leq \left(\iiint_{\mathbb{R}^3} |g(\boldsymbol{r}')|\,\mathrm{d}^3\boldsymbol{r}'\right)^2 \iiint_{\mathbb{R}^3} |f(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}.$$

In the current context, we may pick any $\boldsymbol{u} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$, and apply the Young inequality to

$$f(\boldsymbol{r}) = \begin{cases} \boldsymbol{u}\cdot\boldsymbol{E}(\boldsymbol{r}), & \boldsymbol{r} \in V \\ 0, & \boldsymbol{r} \notin V \end{cases}, \quad g(\boldsymbol{r}) = \begin{cases} e^{-ik|\boldsymbol{r}|}|\boldsymbol{r}|^{-m}, & |\boldsymbol{r}| < 2\max_{\boldsymbol{r}',\boldsymbol{r}''\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}''| \\ 0, & |\boldsymbol{r}| \geq 2\max_{\boldsymbol{r}',\boldsymbol{r}''\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}''| \end{cases},$$

so as to confirm that all the matrix elements of $\hat{\Gamma}_m(\boldsymbol{r},\boldsymbol{r}'), m = 1,2$ contribute to a continuous linear mapping.

To show that $G_3 < +\infty$, we need the Calderón-Zygmund theory in harmonic analysis that deals with singular integral operators [48]. Specifically, the convolution kernel $K(\boldsymbol{r},\boldsymbol{r}')$ in question (i.e. any matrix element of $\hat{\Gamma}_3(\boldsymbol{r},\boldsymbol{r}')$) qualifies as a singular integral operator of Calderón-Zygmund type (see Ref. [49], p. 293, also Ref. [25], p. 39) — for some finite positive constant $A < +\infty$, the convolution kernel $K(\boldsymbol{r},\boldsymbol{r}'), \boldsymbol{r} \neq \boldsymbol{r}'$ satisfies the following inequalities:

$$\begin{aligned}
|K(\boldsymbol{r},\boldsymbol{r}')| &\leq \frac{A}{|\boldsymbol{r}-\boldsymbol{r}'|^d},\\
|K(\boldsymbol{r}_1,\boldsymbol{r}') - K(\boldsymbol{r}_2,\boldsymbol{r}')| &\leq \frac{A|\boldsymbol{r}_1-\boldsymbol{r}_2|}{|\boldsymbol{r}_1-\boldsymbol{r}'|^{d+1}} \quad \text{if } |\boldsymbol{r}_1-\boldsymbol{r}_2| \leq \frac{|\boldsymbol{r}_1-\boldsymbol{r}'|}{2},\\
|K(\boldsymbol{r},\boldsymbol{r}'_1) - K(\boldsymbol{r},\boldsymbol{r}'_2)| &\leq \frac{A|\boldsymbol{r}'_1-\boldsymbol{r}'_2|}{|\boldsymbol{r}'_1-\boldsymbol{r}|^{d+1}} \quad \text{if } |\boldsymbol{r}'_1-\boldsymbol{r}'_2| \leq \frac{|\boldsymbol{r}'_1-\boldsymbol{r}|}{2},
\end{aligned} \tag{1.27}$$

where $d = 3$ is the dimensionality of the space under consideration. Furthermore, the Calderón-Zygmund kernels $K(\boldsymbol{r},\boldsymbol{r}')$ that arise from any one of the matrix elements of $\hat{\Gamma}_3(\boldsymbol{r},\boldsymbol{r}')$ also satisfy the "cancellation condition" (see Ref. [49], p. 291 and pp. 305-306, also Ref. [25], p. 39)

$$\iiint_{\varepsilon<|\boldsymbol{r}-\boldsymbol{r}'|<N} K(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' = 0, \quad \text{whenever } 0 < \varepsilon < N < +\infty.$$

Here, for off-diagonal elements of $\hat{\Gamma}_3(\boldsymbol{r},\boldsymbol{r}')$, the cancellation results from reflection symmetry, which leads to

$$\iiint_{\varepsilon^2<X^2+Y^2+Z^2<N^2} \frac{g(|\boldsymbol{R}|)}{|\boldsymbol{R}|^2} XY\,\mathrm{d}X\,\mathrm{d}Y\,\mathrm{d}Z = 0, \text{ etc.} \tag{1.28}$$

where $g(|\boldsymbol{R}|), |\boldsymbol{R}| = |\boldsymbol{r}-\boldsymbol{r}'| = \sqrt{X^2+Y^2+Z^2}$ is a generic spherically symmetric function; for the diagonal elements of $\hat{\Gamma}_3(\boldsymbol{r},\boldsymbol{r}')$, the cancellation results from rotation symmetry, i.e. we have

$$\iiint_{\varepsilon^2<X^2+Y^2+Z^2<N^2} \frac{g(|\boldsymbol{R}|)}{|\boldsymbol{R}|^2} X^2\,\mathrm{d}X\,\mathrm{d}Y\,\mathrm{d}Z = \frac{1}{3}\iiint_{\varepsilon^2<X^2+Y^2+Z^2<N^2} g(|\boldsymbol{R}|)\,\mathrm{d}X\,\mathrm{d}Y\,\mathrm{d}Z, \text{ etc.} \tag{1.29}$$

It is a standard result in harmonic analysis that a Calderón-Zygmund kernel satisfying the cancellation condition induces a bounded linear operator from $L^2(\mathbb{R}^3;\mathbb{C}^3)$ to $L^2(\mathbb{R}^3;\mathbb{C}^3)$, which necessarily specializes to a continuous mapping from $L^2(V;\mathbb{C}^3)$ to $L^2(V;\mathbb{C}^3)$. This shows that there exists a finite constant $G_3 < +\infty$, such that $\|\hat{\mathcal{G}}_3\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \leq G_3\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$ for all $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$. ▮

In the light of this, we can say that the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ defined by distributional derivatives in S1.0.2 (and accordingly the Born operator $\hat{\mathcal{B}} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ defined by Fourier transforms in S1.1.1) is mathematically equivalent to the formulation of discrete dipole approximation (i.e. digitized Green function approach) $\hat{\mathcal{B}}' : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ in the literature on light scattering. In S1.1 and S1.2, we were not using the matrix representation of the convolution kernel $\hat{\Gamma}_m(\boldsymbol{r},\boldsymbol{r}'), m = 1,2,3$ as in $\hat{\mathcal{B}}' : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ for further analysis, because bound estimates based on $G_m, m = 1,2,3$ are quantitatively less transparent than the bound estimates derived from the distributional derivatives in S1.0.2 (and accordingly the Fourier transforms in S1.1.1).

Nonetheless, the Calderón-Zygmund theory has the capability of handling non-convolution kernels $K(\boldsymbol{r},\boldsymbol{r}')$ that do not depend solely on the displacement $\boldsymbol{r}-\boldsymbol{r}'$ (see Ref. [49], Chap. VII). We mention, in advance, that the general Calderón-Zygmund theory plays an indispensable rôle in the proof of the "optical resonance theorem" — a critical result concerning light scattering in dielectric media with smooth boundary (Theorem 2.1 in S2.2).

S1.3.2. Rigorous Derivations of Some Fourier Inversion Formulae

The major arguments in S1.1.1 rely on some Fourier inversion formulae, which we now justify in detail.

Lemma 1.2 *The following Fourier inversion formulae hold for all* $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$*:*

$$(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad \forall \boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\},$$

where the partial derivatives are interpreted in the distributional sense.

Proof First, we point out that a convolution with the Newtonian potential $(4\pi|\boldsymbol{r}-\boldsymbol{r}'|)^{-1}$ in $\boldsymbol{r}$-space can be rendered as a multiplication of $|\boldsymbol{q}|^{-2}$ in $\boldsymbol{q}$-space

$$(\hat{\mathcal{N}}\boldsymbol{F})(\boldsymbol{r}) = \iiint_{\mathbb{R}^3}\frac{\boldsymbol{F}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}, \boldsymbol{r}\in\mathbb{R}^3\quad\text{for } \tilde{\boldsymbol{F}}(\boldsymbol{q}) := \iiint_{\mathbb{R}^3}\boldsymbol{F}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r}$$

so long as $\boldsymbol{F}\in\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$ (see Lemma 1(a) on p. 117 of Ref. [25]). Here, the Schwartz space is defined by

$$\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3) := \left\{\boldsymbol{F}\in C^\infty(\mathbb{R}^3;\mathbb{C}^3)\,\middle|\,\sup_{\boldsymbol{r}\in\mathbb{R}^3}|\boldsymbol{r}^{\boldsymbol{\mu}}D^{\boldsymbol{\nu}}\boldsymbol{F}(\boldsymbol{r})| < +\infty, \forall\boldsymbol{\mu},\boldsymbol{\nu}\right\}$$

where $\boldsymbol{r}^{\boldsymbol{\mu}}$ is a monomial $x^{\mu_x}y^{\mu_y}z^{\mu_z}$ of degree $\mu_x+\mu_y+\mu_z = |\boldsymbol{\mu}|$. In simple terms, the Schwartz space $\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$ forms the totality of infinitely differentiable vector fields with rapidly decaying amplitudes at infinity. If $\boldsymbol{F}\in\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$, then $\tilde{\boldsymbol{F}}\in\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$, and vice versa (Corollary 2.2.15 in Ref. [38]). It is clear from the definition that $C_0^\infty(V;\mathbb{C}^3)\subset\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$.

Then, we note that using integration by parts, one can justify that $\hat{\mathcal{N}}\boldsymbol{F}\in C^2(\mathbb{R}^3;\mathbb{C}^3)$ for all $\boldsymbol{F}\in\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$. Meanwhile, $\tilde{\boldsymbol{F}}\in\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$ implies the absolute convergence of the following integrals:

$$\iiint_{\mathbb{R}^3}\frac{|(\boldsymbol{u}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})|}{|\boldsymbol{q}|^2}\,\mathrm{d}^3\boldsymbol{q} < +\infty,\quad \iiint_{\mathbb{R}^3}\frac{|(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})|}{|\boldsymbol{q}|^2}\,\mathrm{d}^3\boldsymbol{q} < +\infty,\quad \forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}$$

and hence the continuity (with respect to $\boldsymbol{r}\in\mathbb{R}^3$) of the corresponding Fourier integrals

$$\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}\in C^0(\mathbb{R}^3;\mathbb{C}^3),\quad \iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}\in C^0(\mathbb{R}^3;\mathbb{C}^3),\quad \forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}.$$

All this information combined allows us to take derivatives under the integral sign, according to the Fubini theorem and the Newton-Leibniz formula, thus leading to the following result for all $\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3)\subset\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$

$$(\hat{\mathcal{D}}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{F})(\boldsymbol{r}) := (\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_{\mathbb{R}^3}\frac{\boldsymbol{F}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad\forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}. \tag{1.30}$$

We recall that $C_0^\infty(V;\mathbb{C}^3)$ is dense in $L^2(V;\mathbb{C}^3)$. For a generic square-integrable vector field $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$, let $\{\boldsymbol{E}_s \in C_0^\infty(V;\mathbb{C}^3) | s = 1, 2, \dots\}$ be a sequence that converges to $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$ in L^2-norm, we then have the L^2-convergence (mean square convergence) of the Fourier inversion formulae:

$$\Delta_s(\boldsymbol{r}) := -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}_s(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} + \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q} \xrightarrow{L^2} \boldsymbol{0}$$

as is evident from the bound estimate of the residual error based on Parseval-Plancherel identity (Eq. 1.7):

$$\|\Delta_s\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\left|\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})[\tilde{\boldsymbol{E}}_s(\boldsymbol{q})-\tilde{\boldsymbol{E}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2}\right|^2\mathrm{d}^3\boldsymbol{r} \le \frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}|\tilde{\boldsymbol{E}}_s(\boldsymbol{q})-\tilde{\boldsymbol{E}}(\boldsymbol{q})|^2\,\mathrm{d}^3\boldsymbol{r} = \|\boldsymbol{E}_s-\boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)} \to 0.$$

Meanwhile, in the $\boldsymbol{r}$-space, we also have $\|\hat{\mathcal{D}}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E}_s - \hat{\mathcal{D}}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E}\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)} \le \sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}\hat{\mathcal{N}}(\boldsymbol{E}_s-\boldsymbol{E})\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)} = \|\boldsymbol{E}_s - \boldsymbol{E}\|^2_{L^2(V;\mathbb{C}^3)} \to 0$. Therefore, the Fourier inversion formula (Eq. 1.30) is preserved in the procedure of taking L^2-limits, and thus extends to all $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$, just as claimed. ▮

Lemma 1.3 *We have the Fourier inversion formula:*

$$(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad \forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}, \tag{1.31}$$

for every $\varepsilon > 0$ and $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$.

Proof To prove Eq. 1.31, we may first consider $\boldsymbol{F} \in C_0^\infty(V;\mathbb{C}^3)$. In this case, owing to the exponential decay factor $e^{-\varepsilon|\boldsymbol{r}-\boldsymbol{r}'|}$ in the convolution kernel and the vanishing modulus $|\boldsymbol{F}(\boldsymbol{r})|$ at the boundary ∂V, the convolution $\iiint_V(4\pi|\boldsymbol{r}-\boldsymbol{r}'|)^{-1}\boldsymbol{F}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$ yields a vector field of type $L^1(\mathbb{R}^3;\mathbb{C}^3)\cap C^2(\mathbb{R}^3;\mathbb{C}^3)$. Thus, we may evaluate the Fourier transform of the convolution via an absolutely convergent integral and apply the Fubini theorem to interchange the order of integrations:

$$\iiint_{\mathbb{R}^3}\left[\iiint_V\frac{\boldsymbol{F}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r} := \iiint_V\boldsymbol{F}(\boldsymbol{r}')e^{i\boldsymbol{q}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\iiint_{\mathbb{R}^3}\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}''|}}{4\pi|\boldsymbol{r}''|}e^{i\boldsymbol{q}\cdot\boldsymbol{r}''}\,\mathrm{d}^3\boldsymbol{r}'' = \frac{\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2},$$

and obtain the Fourier inversion formula in the form of a usual integral:

$$\iiint_V\frac{\boldsymbol{F}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q},\quad\forall\boldsymbol{F}\in C_0^\infty(V;\mathbb{C}^3).$$

Now that we have the absolute convergence of the following integrals:

$$\iiint_{\mathbb{R}^3}\left|\frac{(\boldsymbol{u}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}\right|\mathrm{d}^3\boldsymbol{q} < +\infty,\qquad\iiint_{\mathbb{R}^3}\left|\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}\right|\mathrm{d}^3\boldsymbol{q} < +\infty,\quad\forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}$$

which implies the continuity of corresponding Fourier integrals, we can proceed to confirm that for every $\boldsymbol{F} \in C_0^\infty(V;\mathbb{C}^3)$, the following identity holds for all $\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}$:

$$(\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{F})(\boldsymbol{r}) := (\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V\frac{\boldsymbol{F}(\boldsymbol{r}')e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = -\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{F}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}.$$

To reach Eq. 1.31 from the equation above, it would suffice to prove that both sides of Eq. 1.31 indeed define a continuous mapping from $L^2(V;\mathbb{C}^3)$ to $L^2(\mathbb{R}^3;\mathbb{C}^3)$.

Here, it is clear that the right-hand side of Eq. 1.31 represents a continuous mapping from $L^2(V;\mathbb{C}^3)$ to $L^2(\mathbb{R}^3;\mathbb{C}^3)$, as the following estimate holds for every $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$:

$$\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\left|\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}\right|^2\mathrm{d}^3\boldsymbol{q} \le \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\max_{\boldsymbol{q}\in\mathbb{R}^3}\left|\frac{(\boldsymbol{u}\cdot\boldsymbol{q})(\boldsymbol{v}\cdot\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}\right|^2 = A_\varepsilon\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$$

where $A_\varepsilon < +\infty$.

The left-hand side of Eq. 1.31 represents a continuous mapping from $L^2(V;\mathbb{C}^3)$ to $L^2(\mathbb{R}^3;\mathbb{C}^3)$ as well, for two reasons.

First, for every $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, one naturally has

$$(\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E})(\boldsymbol{r})=(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_{\mathbb{R}^3}\frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'+(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')[e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}-1]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'.$$

Pick $R_V:=\min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ to be the radius of the circumscribing sphere, and we may assume, without loss of generality, that $V\subset O(\mathbf{0},2R_V)$. Thus, we accordingly have the following bound estimate on the open ball $O(\mathbf{0},2R_V)$

$$\begin{aligned}&\|\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E}\|_{L^2(O(\mathbf{0},2R_V),\mathbb{C}^3)}\\ \le{}&\left(1+\sqrt{\iiint_{O(\mathbf{0},2R_V)}\left\{\iiint_V\operatorname{Tr}\left[\left(\nabla\nabla\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)^*\left(\nabla\nabla\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}-1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)\right]\mathrm{d}^3\boldsymbol{r}'\right\}\mathrm{d}^3\boldsymbol{r}}\right)\|\boldsymbol{E}\|_{L^2(V,\mathbb{C}^3)}.\end{aligned}$$

Here, the multiple integral under the square root is finite, owing to the square integrability of the weak singularity of order $O(|\boldsymbol{r}-\boldsymbol{r}'|^{-1})$.

Second, we note that so long as $\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$, we have the square-integrable convolution kernel:

$$\iiint_V\left|(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right|^2\mathrm{d}^3\boldsymbol{r}<+\infty,$$

so we may differentiate under the integral sign for $\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E}\in C^0(\mathbb{R}^3\setminus(V\cup\partial V);\mathbb{C}^3)$ to deduce

$$(\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E})(\boldsymbol{r}):=\iiint_V\boldsymbol{E}(\boldsymbol{r}')(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}',\quad\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$$

and perform the estimate

$$\begin{aligned}&\|\hat{\mathcal{D}}^{(\varepsilon)}_{\boldsymbol{u},\boldsymbol{v}}\boldsymbol{E}\|_{L^2(\mathbb{R}^3\setminus O(\mathbf{0},2R_V),\mathbb{C}^3)}\\ \le{}&\sqrt{\iiint_{\mathbb{R}^3\setminus O(\mathbf{0},2R_V)}\left\{\iiint_V\operatorname{Tr}\left[\left(\nabla\nabla\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)^*\left(\nabla\nabla\frac{e^{-i(k-i\varepsilon)|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right)\right]\mathrm{d}^3\boldsymbol{r}'\right\}\mathrm{d}^3\boldsymbol{r}}\,\|\boldsymbol{E}\|_{L^2(V,\mathbb{C}^3)}.\end{aligned}$$

Here, the multiple integral under the square root is finite, owing to the exponential decay factor $e^{-\varepsilon|\boldsymbol{r}-\boldsymbol{r}'|}$.

Combining the above efforts, we have verified the Fourier inversion formula as claimed in Eq. 1.31. ∎

S1.3.3. Norm Estimates of the Born Operator

In the quantitative analysis of the Born operator, we need to bound the operator norm $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}$ for

$$(\hat{\gamma}\boldsymbol{E})(\boldsymbol{r}):=\iiint_V\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'$$

where the 3×3 matrix $\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')$ is given by[22]

$$\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')=\frac{k^2}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\left\{\left[e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}+\frac{1-e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}(1+ik|\boldsymbol{r}-\boldsymbol{r}'|)}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2}\right]\nVdash+\left[3\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}(1+ik|\boldsymbol{r}-\boldsymbol{r}'|)-1}{k^2|\boldsymbol{r}-\boldsymbol{r}'|^2}-e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}\right]\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\right\}.$$

Here, the 3×3 matrix

$$\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}=\frac{\boldsymbol{r}-\boldsymbol{r}'}{|\boldsymbol{r}-\boldsymbol{r}'|}\left(\frac{\boldsymbol{r}-\boldsymbol{r}'}{|\boldsymbol{r}-\boldsymbol{r}'|}\right)^{\mathrm{T}}$$

---

[22] Hereafter, without affecting the evaluation of the integral, we may always assume that $\boldsymbol{r}\neq\boldsymbol{r}'$, so that the matrix is never ill-defined.

satisfies $\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}\hat{\boldsymbol{r}} = \hat{\boldsymbol{r}}$, so it has an eigenvalue $\lambda_1 = 1$, with eigenvector $\hat{\boldsymbol{r}}$. The three eigenvalues of $\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}$ are actually $\lambda_1 = 1, \lambda_2 = \lambda_3 = 0$, as evident from the computations $\operatorname{Tr}(\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}) = \lambda_1 + \lambda_2 + \lambda_3 = 1$ and $\det(\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}) = \lambda_1\lambda_2\lambda_3 = 0$.

To facilitate further analysis, we may use the decomposition $\hat{\gamma} = \hat{\gamma}_C - i\hat{\gamma}_S : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ where

$$(\hat{\gamma}_C \boldsymbol{E})(\boldsymbol{r}) = \nabla\times\nabla\times \iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')[\cos(k|\boldsymbol{r}-\boldsymbol{r}'|)-1]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad (\hat{\gamma}_S \boldsymbol{E})(\boldsymbol{r}) = \nabla\times\nabla\times \iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'.$$

We may accordingly define $\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}') = \operatorname{Re}\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')$ and $\hat{\Gamma}_S(\boldsymbol{r},\boldsymbol{r}') = -\operatorname{Im}\hat{\Gamma}(\boldsymbol{r},\boldsymbol{r}')$, so that $(\hat{\gamma}_C\boldsymbol{E})(\boldsymbol{r}) = \iiint_V \hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'$ and $(\hat{\gamma}_S\boldsymbol{E})(\boldsymbol{r}) = \iiint_V \hat{\Gamma}_S(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}'$. The bound estimate for the operator norm of $\hat{\gamma} = \hat{\gamma}_C - i\hat{\gamma}_S$ can be then obtained from $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} \leq \|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)} + \|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}$.

Lemma 1.4 *We have the following upper bound estimate*

$$\max\{\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)}, \|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}\} \leq \frac{2}{3\pi}\sqrt[3]{\frac{2}{\pi}}\left(\frac{1}{2}+\frac{\pi}{6}\right)\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3} \leq \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}.$$

Proof For brevity, we will narrate the proof for $\hat{\gamma}_C$ only, as the same procedure, *mutatis mutandis*, will also apply to $\hat{\gamma}_S$.

It is easy to recognize that $\hat{\gamma}_C$ is a compact Hermitian operator. The corresponding absolute value of the Rayleigh quotient $|\langle\boldsymbol{E},\hat{\gamma}_C\boldsymbol{E}\rangle_V/\langle\boldsymbol{E},\boldsymbol{E}\rangle_V|$ (defined for $\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} \neq 0$) attains its maximum when $\boldsymbol{E}$ is an eigenvector subordinate to the eigenvalue of $\hat{\gamma}_C$ with the largest modulus, and we have the equality $\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)} = \max_{\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\neq 0}|\langle\boldsymbol{E},\hat{\gamma}_C\boldsymbol{E}\rangle_V/\langle\boldsymbol{E},\boldsymbol{E}\rangle_V| = \max_{\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=1}|\langle\boldsymbol{E},\hat{\gamma}_C\boldsymbol{E}\rangle_V|$ as well. Consequently, we just need to bound the absolute value of the quadratic form $\langle\boldsymbol{E},\hat{\gamma}_C\boldsymbol{E}\rangle_V$ from above for an upper bound estimate of $\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)}$.

Now, we have the inequality

$$\begin{aligned}
|\langle\boldsymbol{E},\hat{\gamma}_C\boldsymbol{E}\rangle_V| &= \left|\iiint_V\iiint_V \boldsymbol{E}^*(\boldsymbol{r})\cdot[\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')]\,\mathrm{d}^3\boldsymbol{r}'\,\mathrm{d}^3\boldsymbol{r}\right| \leq \iiint_V\iiint_V |\boldsymbol{E}^*(\boldsymbol{r})||\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')|\,\mathrm{d}^3\boldsymbol{r}'\,\mathrm{d}^3\boldsymbol{r}\\
&\leq \iiint_V\iiint_V |\boldsymbol{E}^*(\boldsymbol{r})|\left\{|\lambda|_{\max}\left[\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\right]\right\}|\boldsymbol{E}(\boldsymbol{r}')|\,\mathrm{d}^3\boldsymbol{r}'\,\mathrm{d}^3\boldsymbol{r}
\end{aligned}$$

where $|\lambda|_{\max}$ extracts the eigenvalue with the largest modulus. As the 3×3 symmetric matrix $\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}') = a\mathbb{I} + b\hat{\boldsymbol{r}}\hat{\boldsymbol{r}}^{\mathrm{T}}$ has three eigenvalues $\lambda_1 = a+b, \lambda_2 = \lambda_3 = a$, we may deduce that

$$|\lambda|_{\max}\left[\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\right] = \frac{k^2}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\max\left\{\left|\operatorname{Re}\left(e^{-i\xi}+\frac{1-e^{-i\xi}(1+i\xi)}{\xi^2}\right)\right|, 2\left|\operatorname{Re}\left(\frac{1-e^{-i\xi}(1+i\xi)}{\xi^2}\right)\right|\right\}, \quad \xi = k|\boldsymbol{r}-\boldsymbol{r}'|.$$

We claim that the above formula implies that

$$|\lambda|_{\max}\left[\hat{\Gamma}_C(\boldsymbol{r},\boldsymbol{r}')\right] \leq \frac{k^2}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\left(\frac{1}{2}+\frac{\pi}{6}\right). \tag{1.32}$$

Here, for $\xi > 0$, we have $|e^{i\xi}-1-i\xi| = |\int_0^\xi(e^{i\eta}-1)\,\mathrm{d}\eta| \leq \int_0^\xi|(e^{i\eta}-1)|\,\mathrm{d}\eta = \int_0^\xi|2\sin(\eta/2)|\,\mathrm{d}\eta \leq \xi^2/2$, thus we obtain $2\left|[1-e^{-i\xi}(1+i\xi)]/\xi^2\right| \leq 1$. Meanwhile, the inequality $|e^{i\xi}-1-i\xi+\xi^2/2| = |\int_0^\xi(e^{i\eta}-1-i\eta)\,\mathrm{d}\eta| \leq \int_0^\xi(\eta^2/2)\,\mathrm{d}\eta = \xi^3/6$ allows us to estimate

$$\Lambda(\xi) := \left|e^{-i\xi}+\frac{1-e^{-i\xi}(1+i\xi)}{\xi^2}\right| \leq \frac{1}{2}+\frac{|e^{i\xi}-1-i\xi+\xi^2/2|}{\xi^2} \leq \frac{1}{2}+\frac{\xi}{6} \leq \frac{1}{2}+\frac{\pi}{6} = 1.0236+, \quad 0 \leq \xi \leq \pi,$$

whereas in the regime $\pi < \xi \leq 3\pi/2$, we have

$$\begin{aligned}
\Lambda(\xi) &= \sqrt{\left(1-\frac{1-\cos\xi}{\xi^2}\right)^2+\frac{(\xi-\sin\xi)^2}{\xi^4}} \leq \sqrt{\left(1-\frac{1-\cos\frac{3\pi}{2}}{\left(\frac{3\pi}{2}\right)^2}\right)^2+\left[\frac{\xi+\sin(\xi-\pi)}{\xi^2}\right]^2}\\
&\leq \sqrt{\left(1-\frac{4}{9\pi^2}\right)^2+\frac{1}{\pi^2}\left[\frac{\pi}{\xi}\left(2-\frac{\pi}{\xi}\right)\right]^2} \leq \sqrt{\left(1-\frac{4}{9\pi^2}\right)^2+\frac{1}{\pi^2}} = 1.0066+.
\end{aligned}$$

Here, we note that the term $(\pi/\xi)[2-(\pi/\xi)]$ is a quadratic form in $1/\xi$ that attains its maximum for $\xi=\pi$. Moreover, we have

$$\Lambda(\xi)\le\sqrt{\left(1-\frac{1-\cos\frac{7\pi}{4}}{\left(\frac{7\pi}{4}\right)^2}\right)^2+\left(\frac{\xi+1}{\xi^2}\right)^2}\le\sqrt{\left(1-\frac{1-\frac{1}{\sqrt{2}}}{\left(\frac{7\pi}{4}\right)^2}\right)^2+\left(\frac{\frac{3\pi}{2}+1}{\left(\frac{3\pi}{2}\right)^2}\right)^2}=1.0231+,\quad 3\pi/2<\xi\le 7\pi/4;$$

$$\Lambda(\xi)\le\sqrt{1+\left(\frac{\xi-\sin\xi}{\xi^2}\right)^2}\le\sqrt{1+\left(\frac{\xi+1}{\xi^2}\right)^2}\le\sqrt{1+\left(\frac{\frac{7\pi}{4}+1}{\left(\frac{7\pi}{4}\right)^2}\right)^2}=1.0228+,\quad \xi>7\pi/4.$$

This establishes Eq. 1.32 as claimed. (One may replace the constant $1/2+\pi/6=1.0236+$ by the numerical value of the maximum $\max_{\xi>0}\Lambda(\xi)=1.014+$, but the improvement is not significant.)

Using the Hardy-Littlewood-Sobolev inequality [50, 51, 52] with Lieb's sharp constant (see Ref. [53], as well as Theorem 4.3 in Ref. [29]):

$$\left|\int_{\mathbb{R}^d}\int_{\mathbb{R}^d}\frac{f(\boldsymbol r')h(\boldsymbol r)}{|\boldsymbol r-\boldsymbol r'|^m}\,\mathrm{d}^d\boldsymbol r'\,\mathrm{d}^d\boldsymbol r\right|\le\pi^{m/2}\frac{\Gamma(\frac{d-m}{2})}{\Gamma(d-\frac{m}{2})}\left[\frac{\Gamma(\frac{d}{2})}{\Gamma(d)}\right]^{-1+\frac{m}{d}}\|f\|_{L^{2d/(2d-m)}(\mathbb{R}^d;\mathbb{C})}\|h\|_{L^{2d/(2d-m)}(\mathbb{R}^d;\mathbb{C})},$$

$$\text{where } d=3, m=1,\frac{2d}{2d-m}=\frac{6}{5};\quad f(\boldsymbol r)=h(\boldsymbol r)=\begin{cases}|\boldsymbol E(\boldsymbol r)|, & \boldsymbol r\in V\\ 0, & \boldsymbol r\notin V\end{cases},$$

we may deduce from Eq. 1.32 the following estimate

$$|\langle\boldsymbol E,\hat\gamma_C\boldsymbol E\rangle_V|\le\frac{2k^2}{3\pi}\sqrt[3]{\frac{2}{\pi}}\left(\frac12+\frac{\pi}{6}\right)\|\boldsymbol E\|^2_{L^{6/5}(\mathbb{R}^d;\mathbb{C}^3)}\le\frac{2\langle\boldsymbol E,\boldsymbol E\rangle_V}{3\pi}\sqrt[3]{\frac{2}{\pi}}\left(\frac12+\frac{\pi}{6}\right)\left(k^3\iiint_V\mathrm{d}^3\boldsymbol r\right)^{2/3}\le\frac{3}{5\pi}\left(k^3\iiint_V\mathrm{d}^3\boldsymbol r\right)^{2/3}.$$

Here, the Hölder inequality (Theorem 2.3 in Ref. [29]) has been used to derive the estimate $\|\boldsymbol E\|_{L^{6/5}(V;\mathbb{C}^3)}\le\|\boldsymbol E\|_{L^2(V;\mathbb{C}^3)}(\iiint_V\mathrm{d}^3\boldsymbol r)^{1/3}$, and the ratio $\frac23(\frac{2}{\pi})^{1/3}(\frac12+\frac{\pi}{6}):\frac35$ lies between 97% and 98%. ▮

Remark When $V=\{(x,y,z)\in\mathbb{R}^3|x^2+y^2<R^2,0<z<h\}$ takes the shape of a cylindrical "antenna", we may compute that $\max\{\|\hat\gamma_C\|_{L^2(V;\mathbb{C}^3)},\|\hat\gamma_S\|_{L^2(V;\mathbb{C}^3)}\}\le\frac25(\frac12+\frac{\pi}{6})(k^3R^2h)^{2/3}$, and the upper bound $\|\hat\gamma\|_{L^2(V;\mathbb{C}^3)}\le\frac45(\frac12+\frac{\pi}{6})(k^3R^2h)^{2/3}$ scales as $(kh)^{2/3}$ for large heights $h\gg 2\pi/k$. When $V=O(\mathbf{0},R)$ is an open ball with radius R, the above reasoning leads to an estimate $\max\{\|\hat\gamma_C\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)},\|\hat\gamma_S\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}\}\le\frac12(kR)^2$. Thus, we may also conclude that $\|\hat\gamma\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}\le(kR)^2$. It is worth pointing out that for the spherical geometry, the quadratic growth rate $(kR)^2$ overestimates the operator norm for large radii $R\gg 2\pi/k$, as indicated by the lemma below. ▮

Lemma 1.5 *We have the following upper bound estimate for the operator norm of* $\hat\gamma_C:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$

$$\|\hat\gamma_C\|_{L^2(V;\mathbb{C}^3)}\ \le\ k\left[\sqrt2\sqrt[3]{\frac{3}{(2\sqrt3+12)\pi}\iiint_V\mathrm{d}^3\boldsymbol r'}+\sqrt[3]{\frac{27}{4}+\frac{9\sqrt3}{8}}\sqrt{R_V}\sqrt[6]{\frac{1}{2\pi}\iiint_V\mathrm{d}^3\boldsymbol r'}\right]+\sqrt2kR_V\le3\sqrt2kR_V,$$

where $R_V:=\min_{\boldsymbol r\in\mathbb{R}^3}\max_{\boldsymbol r'\in V\cup\partial V}|\boldsymbol r'-\boldsymbol r|$ *is the minimal radius of a circumscribed sphere.*

Proof Without compounding the validity of the proof, we may regard $\boldsymbol E\in C_0^\infty(V;\mathbb{C}^3)$ as a vector field with compact support, and accordingly $\hat\gamma_C\boldsymbol E\in C^\infty(\mathbb{R}^3;\mathbb{C}^3)$ as a smooth vector field defined on the entire space. This is because we have $\|\hat\gamma_C\|_{L^2(V;\mathbb{C}^3)}=\max_{\|\boldsymbol E\|_{L^2(V;\mathbb{C}^3)}=1}|\langle\boldsymbol E,\hat\gamma_C\boldsymbol E\rangle_V|=\sup_{\boldsymbol E\in C_0^\infty(V;\mathbb{C}^3)\setminus\{\mathbf{0}\}}|\langle\boldsymbol E,\hat\gamma_C\boldsymbol E\rangle_V/\langle\boldsymbol E,\boldsymbol E\rangle_V|$. For convenience, we may also assume that the center of the circumscribed sphere with radius R_V coincides with the origin $\boldsymbol r=\mathbf{0}$ of the coordinate system, so we have the set inclusion relation $V\subset O(\mathbf{0},R_V)$. Furthermore, we may confine our discussion to the regime where $kR_V>2\sqrt2$, because the conclusion in Lemma 1.4 already implies $\|\hat\gamma_C\|_{L^2(V;\mathbb{C}^3)}\le\frac12(kR_V)^2\le\sqrt2kR_V$ for $kR_V\le2\sqrt2$.

Now that $\boldsymbol E\in C_0^\infty(V;\mathbb{C}^3)\subset\mathcal{S}(\mathbb{R}^3;\mathbb{C}^3)$ is a member of the Schwartz space, the Fourier transform $\tilde{\boldsymbol E}(\boldsymbol q)$ will decay sufficiently fast as $|\boldsymbol q|\to+\infty$. This allows us to interchange limit with integrations for the Fourier representation of the quadratic form in question

$$\begin{aligned}\langle \boldsymbol E,\hat\gamma_C\boldsymbol E\rangle_V &= k\lim_{\varepsilon\to0^+}\frac{1}{(2\pi)^3}\iiint_{\mathbb R^3}\left[\frac{1}{2k}\left(\frac{1}{|\boldsymbol q|^2-(k-i\varepsilon)^2}+\frac{1}{|\boldsymbol q|^2-(k+i\varepsilon)^2}\right)-\frac{1}{|\boldsymbol q|^2k}\right]|\boldsymbol q\times\tilde{\boldsymbol E}(\boldsymbol q)|^2\,\mathrm d^3\boldsymbol q\\ &= -\frac{k}{(2\pi)^3}\lim_{\varepsilon\to0^+}\int_0^{+\infty}\left[\iiint_{\mathbb R^3}\cos(|\boldsymbol q|\Xi)\frac{|\boldsymbol q\times\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q\right]e^{-\varepsilon\Xi}\sin(k\Xi)\,\mathrm d\,\Xi\\ &= -\frac{k}{(2\pi)^3}\int_0^{+\infty}\left[\iiint_{\mathbb R^3}\cos(|\boldsymbol q|\Xi)\frac{|\boldsymbol q\times\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q\right]\sin(k\Xi)\,\mathrm d\,\Xi.\end{aligned}$$

By splitting the integration $\int_0^{+\infty}\mathrm d\,\Xi=\int_0^{R_*}\mathrm d\,\Xi+\int_{R_*}^{+\infty}\mathrm d\,\Xi$ for $R_*>2R_V=2\min_{\boldsymbol r\in\mathbb R^3}\max_{\boldsymbol r'\in V\cup\partial V}|\boldsymbol r'-\boldsymbol r|$, we obtain the representation $\langle \boldsymbol E,\hat\gamma_C\boldsymbol E\rangle_V=I_1[k;V]+I_2[k;V]$, where

$$I_1[k;V]=-\frac{k}{(2\pi)^3}\int_0^{R_*}\left[\iiint_{\mathbb R^3}\cos(|\boldsymbol q|\Xi)\frac{|\boldsymbol q\times\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q\right]\sin(k\Xi)\,\mathrm d\,\Xi$$

satisfies

$$\begin{aligned}&|I_1[k;V]|\le\frac{k}{(2\pi)^3}\iiint_{\mathbb R^3}\frac{|\boldsymbol q\times\tilde{\boldsymbol E}(\boldsymbol q)|^2}{|\boldsymbol q|^2}\,\mathrm d^3\boldsymbol q\int_0^{R_*}|\sin(k\Xi)|\,\mathrm d\,\Xi\le\langle\boldsymbol E,\boldsymbol E\rangle_V\int_0^{kR_*}|\sin\xi|\,\mathrm d\,\xi,\quad\text{with } kR_*>4\sqrt2,\\ &\text{and }\frac{1}{kR_*}\int_0^{kR_*}|\sin\xi|\,\mathrm d\,\xi=\frac{2}{\pi}-\frac{4}{\pi}\sum_{m=1}^{\infty}\frac{\sin(2mkR_*)}{2mkR_*(4m^2-1)}\le\frac{2}{\pi}+\frac{1}{\sqrt2\pi}\sum_{m=1}^{\infty}\frac{1}{2m(4m^2-1)}=\frac{2}{\pi}+\frac{\ln4-1}{2\sqrt2\pi}\le\frac{1}{\sqrt2};\end{aligned}$$

while $I_2[k;V]$ can be recast into the configuration space ($\boldsymbol r$-space) as

$$I_2[k;V]=k\int_{R_*}^{+\infty}\left\{\iiint_V\boldsymbol E^*(\boldsymbol r'')\cdot\frac{\partial}{\partial\Xi}\left[\oiint_{|\boldsymbol r-\boldsymbol r''|=\Xi,\boldsymbol r''\in V}\left(\nabla\times\nabla\times\iiint_V\frac{\boldsymbol E(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d^3\boldsymbol r'\right)\frac{\mathrm d\,S}{4\pi\Xi}\right]\mathrm d^3\boldsymbol r''\right\}\sin(k\Xi)\,\mathrm d\,\Xi.$$

For $|\boldsymbol r-\boldsymbol r''|=\Xi\ge R_*>2R_V,\boldsymbol r''\in V$, a point $\boldsymbol r\in V$ will not contribute to the integral representation of $I_2[k;V]$. In this case, we only need to consider $\boldsymbol r\notin V$, and

$$\nabla\times\nabla\times\iiint_V\frac{\boldsymbol E(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d^3\boldsymbol r'=\iiint_V\left(\nabla\nabla\frac{1}{4\pi|\boldsymbol r-\boldsymbol r'|}\right)\boldsymbol E(\boldsymbol r')\,\mathrm d^3\boldsymbol r'=\iiint_V\left(\frac{3\hat{\boldsymbol r}\hat{\boldsymbol r}^{\mathrm T}-\nVdash}{4\pi|\boldsymbol r-\boldsymbol r'|^3}\right)\boldsymbol E(\boldsymbol r')\,\mathrm d^3\boldsymbol r'$$

satisfies

$$\left|\nabla\times\nabla\times\iiint_V\frac{\boldsymbol E(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d^3\boldsymbol r'\right|\le\iiint_V\frac{|\boldsymbol E(\boldsymbol r')|}{2\pi|\boldsymbol r-\boldsymbol r'|^3}\,\mathrm d^3\boldsymbol r'\le\frac{\iiint_V|\boldsymbol E(\boldsymbol r')|\,\mathrm d^3\boldsymbol r'}{2\pi(\Xi-2R_V)^3}\le\frac{\|\boldsymbol E\|_{L^2(V;\mathbb C^3)}\sqrt{\iiint_V\mathrm d^3\boldsymbol r'}}{2\pi(\Xi-2R_V)^3}$$

when $|\boldsymbol r-\boldsymbol r''|=\Xi$ holds for some $\boldsymbol r''\in V$. Here, the triangle inequality $|\boldsymbol r-\boldsymbol r'|+2R_V\ge|\boldsymbol r-\boldsymbol r'|+|\boldsymbol r'-\boldsymbol r''|\ge|\boldsymbol r-\boldsymbol r''|=\Xi$ has been exploited.

Using the fact that $|\cos(|\boldsymbol q|\Xi)|\le1$ and $|\sin(k\Xi)|\le1$, we may proceed with estimating $|I_2[k;V]|$ by

$$\begin{aligned}|I_2[k;V]|&\le k\|\boldsymbol E\|_{L^2(V;\mathbb C^3)}\int_{R_*}^{+\infty}\sqrt{\iiint_{\partial O(\boldsymbol r,\Xi)\cap V\neq\emptyset}\left|\nabla\times\nabla\times\iiint_V\frac{\boldsymbol E(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d^3\boldsymbol r'\right|^2\mathrm d^3\boldsymbol r}\,\mathrm d\,\Xi\\ &\le\frac{k\langle\boldsymbol E,\boldsymbol E\rangle_V}{2\pi}\sqrt{\iiint_V\mathrm d^3\boldsymbol r'}\int_{R_*}^{+\infty}\frac{\sqrt{\iiint_{\Xi-R_V\le|\boldsymbol r|\le\Xi+R_V}\mathrm d^3\boldsymbol r}}{(\Xi-2R_V)^3}\,\mathrm d\,\Xi.\end{aligned}$$

Here, we note that if $|\boldsymbol r-\boldsymbol r''|=\Xi$ holds for some $\boldsymbol r''\in V$, then we must have $\Xi=|\boldsymbol r-\boldsymbol r''|\le|\boldsymbol r|+|\boldsymbol r''|\le|\boldsymbol r|+R_V$, as well as $\Xi+R_V\ge\Xi+|\boldsymbol r''|=|\boldsymbol r-\boldsymbol r''|+|\boldsymbol r''|\ge|\boldsymbol r|$, which justifies the domain of integration $\Xi-R_V\le|\boldsymbol r|\le\Xi+R_V$. The computation

$$\sqrt{\iiint_{\Xi-R_V\le|\boldsymbol r|\le\Xi+R_V}\mathrm d^3\boldsymbol r}=\sqrt{\frac{8\pi R_V(R_V^2+3\Xi^2)}{3}}\le\sqrt{\frac{8\pi R_V}{3}}(R_V+\sqrt3\Xi)$$

brings us the estimate

$$|I_2[k;V]| \;\leq\; \frac{k\langle \boldsymbol{E},\boldsymbol{E}\rangle_V}{2\pi}\sqrt{\frac{8\pi R_V}{3}\iiint_V \mathrm{d}^3\boldsymbol{r}'}\left[\frac{\sqrt{3}}{R_*-2R_V}+\frac{(2\sqrt{3}+1)R_V}{2(R_*-2R_V)^2}\right].$$

Now, we may choose an appropriate radius $R_* > 2R_V$ to make the right-hand side of $\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)} \leq |I_1[k;V]| + |I_2[k;V]|$ as small as possible. By choosing R_* to minimize the sum

$$\frac{R_*-2R_V}{\sqrt{2}}+\frac{(2\sqrt{3}+1)\sqrt{\frac{8\pi}{3}R_V^3\iiint_V \mathrm{d}^3\boldsymbol{r}'}}{2\pi(R_*-2R_V)^2}$$

we obtain the upper bound

$$\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)} \leq k\left[\sqrt{2}\sqrt[3]{\frac{3}{(2\sqrt{3}+12)\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}'}+\sqrt[3]{\frac{27}{4}+\frac{9\sqrt{3}}{8}}\sqrt{R_V}\sqrt[6]{\frac{1}{2\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}'}\right]+\sqrt{2}kR_V. \tag{1.33}$$

Inserting the inequality $\iiint_V \mathrm{d}^3\boldsymbol{r} \leq 4\pi R_V^3/3$ into the right-hand side of Eq. 1.33, we may confirm that $\|\hat{\gamma}_C\|_{L^2(V;\mathbb{C}^3)} \leq 3\sqrt{2}kR_V$. ▮

Remark In case $V = O(\mathbf{0},R_V)$, then we have $\|\hat{\gamma}_C\|_{L^2(V=O(\mathbf{0},R_V);\mathbb{C}^3)}/(kR_V) \leq 4.237+$ according to Eq. 1.33. On the other hand, we may numerically solve the equation $\partial u(R_*)/\partial R_* = 0$ where

$$u(R_*) = \frac{1}{R_V}\left\{\frac{R_*-2R_V}{\sqrt{2}}+\frac{1}{2\pi}\sqrt{\frac{8\pi R_V}{3}\iiint_{V=O(\mathbf{0},R_V)} \mathrm{d}^3\boldsymbol{r}'}\left[\frac{\sqrt{3}}{R_*-2R_V}+\frac{(2\sqrt{3}+1)R_V}{2(R_*-2R_V)^2}\right]\right\}+\sqrt{2},\quad R_*>2R_V$$

thereby verifying that the minimum of $u(R_*), R_* > 2R_V$ is 4.146+. Therefore, although we have avoided a global optimization of R_* (which inevitably involves a non-degenerate cubic equation $\partial u(R_*)/\partial R_* = 0$), the difference in the bound estimates is less than 2.5% for the spherical geometry. For generic geometries where $\iiint_V \mathrm{d}^3\boldsymbol{r} \leq 4\pi R_V^3/3$, the solution to the equation $\partial u(R_*)/\partial R_* = 0$ will not make the optimal ratio $(R_*-2R_V)/R_V$ exceed that of the spherical geometry, hence it is likely that the inverse square term $\propto (R_*-2R_V)^{-2}$ will always dominate the inverse linear term $\propto (R_*-2R_V)^{-1}$ in the expression for $u(R_*)$. Thus, the strategy employed in the proof may lead to a relatively tight upper bound for an arbitrary shape $V \subset O(\mathbf{0},R_V)$.

In principle, we may deduce an upper bound for $\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}$ by applying a similar trick to

$$\langle \boldsymbol{E},\hat{\gamma}_S\boldsymbol{E}\rangle_V = \frac{k}{(2\pi)^3}\int_0^{+\infty}\left[\iiint_{\mathbb{R}^3}\cos(|\boldsymbol{q}|\Xi)\frac{|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})|^2}{|\boldsymbol{q}|^2}\,\mathrm{d}^3\boldsymbol{q}\right]\cos(k\Xi)\,\mathrm{d}\,\Xi.$$

However, using such an approach, we at best obtain an upper bound that is greater than $\sqrt{2}kR_V$, which is less sharp than the estimate $5kR_V/8$ to be given by Lemma 1.7 in S1.3.4. ▮

S1.3.4. Upper Bounds for the Total Scattering Cross-Section

From Eq. 1.18, we know that the total scattering cross-section (up to a scaling factor) is determined by

$$\langle \hat{\gamma}_S\boldsymbol{E},\boldsymbol{E}\rangle_V = \iiint_V\left[\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}^*(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right]\cdot\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r} = \frac{k^3}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\tilde{\boldsymbol{E}}(k\boldsymbol{n})\right|^2\mathrm{d}\,\Omega, \tag{1.34}$$

where the linear mapping $\hat{\gamma}_S : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$

$$(\hat{\gamma}_S\boldsymbol{E})(\boldsymbol{r}) = \nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'$$

is a positive-semidefinite and compact operator. The operator norm $\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}$ is thus equal to the largest eigenvalue of the linear mapping $\hat{\gamma}_S : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$.

Lemma 1.6 *If* $V_1 \subset V_2$, *then* $\|\hat{\gamma}_S\|_{L^2(V_1;\mathbb{C}^3)} \leq \|\hat{\gamma}_S\|_{L^2(V_2;\mathbb{C}^3)}$.

Proof Suppose that $\boldsymbol{E}_{V_1} \in L^2(V_1;\mathbb{C}^3)$ solves the eigenequation $\hat{\gamma}_S\boldsymbol{E}_{V_1} = \|\hat{\gamma}_S\|_{L^2(V_1;\mathbb{C}^3)}\boldsymbol{E}_{V_1}$, and satisfies $\|\boldsymbol{E}_{V_1}\|_{L^2(V_1;\mathbb{C}^3)} = 1$. Then, we may take

$$\boldsymbol{E}(\boldsymbol{r}) = \begin{cases} \boldsymbol{E}_{V_1}(\boldsymbol{r}), & \boldsymbol{r} \in V_1 \\ \boldsymbol{0}, & \boldsymbol{r} \in V_2 \setminus V_1 \end{cases}$$

so that $\|\boldsymbol{E}\|_{L^2(V_2;\mathbb{C}^3)} = 1$ and

$$\|\hat{\gamma}_S\|_{L^2(V_1;\mathbb{C}^3)} = \langle \hat{\gamma}_S\boldsymbol{E}_{V_1}, \boldsymbol{E}_{V_1}\rangle_{V_1} = \langle \hat{\gamma}_S\boldsymbol{E}, \boldsymbol{E}\rangle_{V_2} \le \|\hat{\gamma}_S\|_{L^2(V_2;\mathbb{C}^3)}\|\boldsymbol{E}\|^2_{L^2(V_2;\mathbb{C}^3)} = \|\hat{\gamma}_S\|_{L^2(V_2;\mathbb{C}^3)}$$

as claimed. ▮

Proposition 1.11 *For any bounded and open volume V, we have*

$$\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)} \le \max_{\ell\in\{0,1,2,\dots\}} \int_0^{kR_V} j_\ell^2(\xi)\xi^2 \,\mathrm{d}\,\xi$$

where $R_V := \min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}|$ is the minimal radius of all the circumscribed spheres, and $j_\ell(x) := (-x)^\ell(\frac{1}{x}\frac{\mathrm{d}}{\mathrm{d}\,x})^\ell(\frac{\sin x}{x})$ denotes the ℓ-th order spherical Bessel function.

Proof As before, we may assume that the center of the circumscribed sphere with radius R_V coincides with the origin $\boldsymbol{r} = \boldsymbol{0}$ of the coordinate system, so $V \subset O(\boldsymbol{0}, R_V)$ implies $\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)} \le \|\hat{\gamma}_S\|_{L^2(O(\boldsymbol{0},R_V);\mathbb{C}^3)}$. We may further assume, without loss of generality, that $V = O(\boldsymbol{0}, R_V)$ in the computations below.

First, we have the decomposition $\hat{\gamma}_S = \hat{\gamma}_{S,\sharp} + \hat{\gamma}_{S,\sharp\sharp} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ such that

$$(\hat{\gamma}_{S,\sharp}\boldsymbol{E})(\boldsymbol{r}) \;=\; k^2\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad (\hat{\gamma}_{s,\sharp\sharp}\boldsymbol{E})(\boldsymbol{r}) = \nabla\left[\nabla\cdot\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right].$$

Then, we point out that $\hat{\gamma}_{S,\sharp} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a positive-semidefinite compact operator satisfying

$$\langle \hat{\gamma}_{S,\sharp}\boldsymbol{E}, \boldsymbol{E}\rangle_V = \frac{k^3}{16\pi^2}\oint\!\!\!\oint_{|\boldsymbol{n}|=1}\left|\tilde{\boldsymbol{E}}(k\boldsymbol{n})\right|^2 \mathrm{d}\,\Omega \ge 0;$$

whereas $\hat{\gamma}_{S,\sharp\sharp} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a negative-semidefinite compact operator satisfying

$$\langle \hat{\gamma}_{S,\sharp\sharp}\boldsymbol{E}, \boldsymbol{E}\rangle_V = -\frac{k^3}{16\pi^2}\oint\!\!\!\oint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\cdot\tilde{\boldsymbol{E}}(k\boldsymbol{n})\right|^2 \mathrm{d}\,\Omega \le 0.$$

Meanwhile, it is also clear that if $\boldsymbol{E}_\lambda \in L^2(V;\mathbb{C}^3)$ solves the eigenequation

$$(\hat{\gamma}_{S,\sharp}\boldsymbol{E}_\lambda)(\boldsymbol{r}) = k^2\iiint_V \frac{\boldsymbol{E}_\lambda(\boldsymbol{r}')\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' = \lambda\boldsymbol{E}_\lambda(\boldsymbol{r})$$

where $\lambda \ne 0$, then the solution must be twice-differentiable $\boldsymbol{E}_\lambda \in C^2(V;\mathbb{C}^3)$, and must satisfy the Helmholtz equation $(\nabla^2 + k^2)\boldsymbol{E}_\lambda(\boldsymbol{r}) = \boldsymbol{0}, \forall\boldsymbol{r} \in V$. Therefore, we have the identity:

$$\|\hat{\gamma}_{S,\sharp}\|_{L^2(V;\mathbb{C}^3)} = \max_{\substack{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\\ \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=1}} \langle \hat{\gamma}_{S,\sharp}\boldsymbol{E}, \boldsymbol{E}\rangle_V = \max_{\substack{\boldsymbol{E}\in C^2(V;\mathbb{C}^3), \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=1\\ (\nabla^2+k^2)\boldsymbol{E}(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in V}} \langle \hat{\gamma}_{S,\sharp}\boldsymbol{E}, \boldsymbol{E}\rangle_V.$$

We recall that any twice-differentiable and square-integrable vector field $\boldsymbol{E} \in C^2(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3)$ satisfying the Helmholtz equation $(\nabla^2 + k^2)\boldsymbol{E}(\boldsymbol{r}) = \boldsymbol{0}, \forall\boldsymbol{r} \in V = O(\boldsymbol{0}, R_V)$ admits a Fourier-Bessel expansion $\boldsymbol{E} = \sum_{\boldsymbol{f}_b\in\mathfrak{B}} c_b\boldsymbol{f}_b$, where the orthonormal basis $\mathfrak{B}$ can be chosen as

$$\mathfrak{B} = \left\{ \left. \frac{\boldsymbol{u}j_\ell(k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)}{\sqrt{\int_0^{R_V} j_\ell^2(kr')r'^2\,\mathrm{d}\,r'}} \right| \boldsymbol{u} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}; \ell \in \mathbb{Z}\cap[0,+\infty) = \{0,1,2,\dots\}; m \in \mathbb{Z}\cap[-\ell,\ell] \right\}.$$

Using the spherical harmonic expansion (cf. Eq. 9.98 in Ref. [9])

$$\frac{\sin(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4\pi k|\boldsymbol{r}-\boldsymbol{r}'|} = \sum_{\ell=0}^{\infty} j_\ell(k|\boldsymbol{r}'|)j_\ell(k|\boldsymbol{r}|)\sum_{m=-\ell}^{\ell} Y^*_{\ell m}(\theta',\phi')Y_{\ell m}(\theta,\phi),$$

we may deduce the eigenequation for $\boldsymbol{f}_b = \boldsymbol{u} j_\ell(k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)/[\int_0^{R_V} j_\ell^2(kr')r'^2\,\mathrm{d}\,r']^{1/2}$ as

$$\hat{\gamma}_{S,\sharp}\boldsymbol{f}_b = k^3\boldsymbol{f}_b\int_0^{R_V} j_\ell^2(kr')r'^2\,\mathrm{d}\,r' = \boldsymbol{f}_b\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi.$$

Thus, we may explicitly compute the operator norm of $\hat{\gamma}_{S,\sharp}$ by writing out its largest eigenvalue

$$\|\hat{\gamma}_{S,\sharp}\|_{L^2(V;\mathbb{C}^3)} = \max_{\substack{\boldsymbol{E}=\sum_{\boldsymbol{f}_b\in\mathfrak{B}} c_b\boldsymbol{f}_b\\ \sum_{\boldsymbol{f}_b\in\mathfrak{B}}|c_b|^2=1}} \langle\hat{\gamma}_{S,\sharp}\boldsymbol{E},\boldsymbol{E}\rangle_V = \max_{\ell\in\mathbb{Z}\cap[0,+\infty)}\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi.$$

With the inequality $\langle\hat{\gamma}_{S,\sharp\sharp}\boldsymbol{E},\boldsymbol{E}\rangle_V = \langle\hat{\gamma}_S\boldsymbol{E},\boldsymbol{E}\rangle_V - \langle\hat{\gamma}_{S,\sharp}\boldsymbol{E},\boldsymbol{E}\rangle_V \le 0$, we arrive at $\langle\hat{\gamma}_S\boldsymbol{E},\boldsymbol{E}\rangle_V \le \langle\hat{\gamma}_{S,\sharp}\boldsymbol{E},\boldsymbol{E}\rangle_V \le \|\hat{\gamma}_{S,\sharp}\|_{L^2(O(\mathbf{0},R_V);\mathbb{C}^3)}\langle\boldsymbol{E},\boldsymbol{E}\rangle_V = \langle\boldsymbol{E},\boldsymbol{E}\rangle_V \max_{\ell\in\mathbb{Z}\cap[0,+\infty)}\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi$, whence we may conclude that

$$\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)} \le \max_{\ell\in\mathbb{Z}\cap[0,+\infty)}\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi. \qquad \blacksquare$$

Lemma 1.7 *We have the inequality*

$$\max_{\ell\in\mathbb{Z}\cap[0,+\infty)}\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi \le \min\left\{\frac{1}{3}(kR_V)^3, \frac{1}{2}(kR_V)^2, \frac{5}{8}kR_V\right\}.$$

Proof First, we point out that the identity $\sum_{L=0}^{\infty}(2L+1)j_L^2(\xi) = 1$ (see Eq. 10.1.50 in Ref. [54], also Ref. [55], p. 152) entails $j_\ell^2(\xi) \le 1/(2\ell+1)$, thus resulting in

$$\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi \le \frac{1}{2\ell+1}\int_0^{kR_V}\xi^2\,\mathrm{d}\,\xi \le \frac{1}{3}(kR_V)^3, \quad \forall\ell\in\mathbb{Z}\cap[0,+\infty).$$

This inequality can be actually sharpened to $4\pi\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)} \le k^3\iiint_V \mathrm{d}^3\boldsymbol{r}$ for arbitrarily shaped V, after we plug the Cauchy-Schwarz inequality $|\tilde{\boldsymbol{E}}(\boldsymbol{q})| \le \|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\sqrt{\iiint_V \mathrm{d}^3\boldsymbol{r}}$ into the right-hand side of Eq. 1.34.

Next, we note that the following identity (see Eq. 10.1.52 in Ref. [54], also Ref. [55], p. 152)

$$\sum_{L=0}^{\infty} j_L^2(\xi) = \frac{1}{2\xi}\int_0^{2\xi}\frac{\sin\eta}{\eta}\,\mathrm{d}\,\eta = \frac{\mathrm{Si}\,(2\xi)}{2\xi},$$

along with the inequality $(\sin\eta)/\eta = \prod_{m=1}^{\infty}\cos(2^{-m}\eta) \le \cos(\eta/2), 0 \le \eta \le \pi$ lead us to $j_\ell^2(\xi) \le \mathrm{Si}\,(\pi)/(2\xi) \le 1/\xi$ for $\xi > 0$, so

$$\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi \le \int_0^{kR_V}\xi\,\mathrm{d}\,\xi = \frac{1}{2}(kR_V)^2, \quad \forall\ell\in\mathbb{Z}\cap[0,+\infty)$$

is also obvious.

Now, we will set out to prove $\int_0^{kR_V} j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi \le 5kR_V/8$ using the following Lommel series (see Ref. [55], pp. 151-152) for $x = kR_V > 0$

$$\begin{aligned}\int_0^x j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi &= \frac{\pi x^2}{4}\left[J^2_{\ell+\frac{1}{2}}(x) - J_{\ell-\frac{1}{2}}(x)J_{\ell+\frac{3}{2}}(x)\right] = \pi\sum_{L=0}^{\infty}\left(\ell+\frac{3}{2}+2L\right)J^2_{\ell+\frac{3}{2}+2L}(x)\\ &= x\sum_{L=0}^{\infty}[2(\ell+1+2L)+1]j^2_{\ell+1+2L}(x),\end{aligned}$$

where the cylindrical Bessel functions $J_\nu(x)$ are related to the spherical Bessel functions by $j_\ell(x) = \sqrt{\pi/2x}J_{\ell+\frac{1}{2}}(x)$.

With the help of the identity $\sum_{L=0}^{\infty}(-1)^L(2L+1)j_L^2(\xi) = \sin(2\xi)/(2\xi)$ (see Eq. 10.1.51 in Ref. [54], also Ref. [55], p. 152), we see that when $\ell \ge 0$ is an even number, we have

$$\frac{1}{x}\int_0^x j_\ell^2(\xi)\xi^2\,\mathrm{d}\,\xi \le \sum_{L=0}^{\infty}(4L+3)j^2_{2L+1}(x) = \frac{1}{2} - \frac{\sin 2x}{4x};$$

When $\ell \geq 0$ is an odd number, we have

$$\frac{1}{x}\int_0^x j_\ell^2(\xi)\xi^2 \,\mathrm{d}\,\xi \leq \sum_{L=1}^{\infty}(4L+1)j_{2L}^2(x) = \frac{1}{2}+\frac{\sin 2x}{4x}-j_0^2(x).$$

On one hand, as $\pi \leq 2x \leq 2\pi$, we have

$$f(x) := \frac{1}{2}-\frac{\sin 2x}{4x} = \frac{1}{2}+\frac{(2x-\pi)}{4x}\frac{\sin(2x-\pi)}{2x-\pi} = \frac{1}{2}+\frac{(2x-\pi)}{4x}\prod_{m=1}^{\infty}\left[1-\frac{(2x-\pi)^2}{m^2\pi^2}\right] \leq \frac{1}{2}+\frac{(2x-\pi)}{4x}\left[1-\frac{(2x-\pi)^2}{\pi^2}\right]$$

where the last term is a quadratic polynomial in x that attains its maximum value $5/8$ for $2x = 3\pi/2$. Meanwhile, as $2x \geq 3\pi$, we have

$$f(x) = \frac{1}{2}-\frac{1}{2}\frac{\sin 2x}{2x} \leq \frac{1}{2}+\frac{1}{6\pi} < \frac{3}{5},$$

so $f(x) \leq 5/8, \forall x > 0$. (One can use numerical computation to verify that $f(x) \leq 0.6086+$, but this numerical bound does not mark an impressive improvement as compared to $5/8 = 0.625$.)

On the other hand, from the equality

$$g(x) := \frac{1}{2}+\frac{\sin 2x}{4x}-j_0^2(x) = \frac{1}{2}+\frac{\sin 2x}{4x}-\frac{\sin^2 x}{x^2} = f(x)-\frac{\sin x\cos x}{x^2}(\tan x - x),$$

we have $g(x) \leq f(x)$ for $0 \leq x \leq \pi$. When $x \geq \pi$, we have

$$g(x) \leq \frac{1}{2}+\frac{1}{4\pi} < \frac{3}{5}.$$

Thus, we have concluded the proof by verifying that $\max(f(x), g(x)) \leq 5/8$ for all positive values $x = kR_V > 0$. ∎

Remark Combining the above results with the conclusions from S1.3.3, we immediately arrive at the following upper bound estimate for $\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}$:

$$\|\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} \;\leq\; \min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, 3\sqrt{2}kR_V\right\} + \min\left\{\frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{5}{8}kR_V\right\}.$$

It is worth pointing out that for open balls $O(\mathbf{0},R)$ with $kR \gg 1$, the linear bound $O(kR)$ indeed describes the growth rate of $\|\hat{\gamma}\|_{L^2(V=O(\mathbf{0},R);\mathbb{C}^3)}$, and cannot be reduced to a lower power of kR. To show this, we may just pick a "unit vector" $\boldsymbol{E}(\boldsymbol{r}) = \boldsymbol{e}_z e^{ik|\boldsymbol{r}|}/(i|\boldsymbol{r}|\sqrt{4\pi R})$, $|\boldsymbol{r}| < R$ satisfying $\|\boldsymbol{E}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} = 1$, perform the following computations in spherical coordinates

$$\begin{aligned}
((\hat{\mathcal{D}}+\hat{\gamma})\boldsymbol{E})(\boldsymbol{r}) &= \nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' \\
&= \frac{1}{\sqrt{4\pi R}}\nabla\times\nabla\times\left\{\frac{\boldsymbol{e}_z e^{-ik|\boldsymbol{r}|}}{k|\boldsymbol{r}|}\int_0^{|\boldsymbol{r}|}\sin(kr')e^{ikr'}\,\mathrm{d}\,r' + \frac{\boldsymbol{e}_z\sin(k|\boldsymbol{r}|)}{k|\boldsymbol{r}|}\int_{|\boldsymbol{r}|}^{R}\mathrm{d}\,r'\right\} \\
&= \frac{1}{2k\sqrt{4\pi R}}\nabla\times\nabla\times\left\{\boldsymbol{e}_z\left[ie^{ik|\boldsymbol{r}|}-\frac{i-2kR}{k|\boldsymbol{r}|}\sin(k|\boldsymbol{r}|)\right]\right\},
\end{aligned}$$

and derive

$$\begin{aligned}
&\|(\hat{\mathcal{D}}+\hat{\gamma})\boldsymbol{E}\|^2_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} \\
&= \frac{2k^2R^2}{9}+\frac{-1-3k^2R^2+6k^4R^4+(1+k^2R^2-2k^4R^4)\cos(2kR)+kR(2+3k^2R^2)\sin(2kR)}{12k^4R^4} \geq \frac{2k^2R^2}{9} \quad (1.35)
\end{aligned}$$

after some routine integrations. As we know that $\|\hat{\mathcal{D}}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} \leq 1$, we may deduce from Eq. 1.35 that

$$\|\hat{\gamma}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} \geq \frac{\sqrt{2}}{3}kR - 1, \quad kR > 3/\sqrt{2},$$

which confirms the linear growth rate $\|\hat{\gamma}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}/(kR) = O(1), kR \to +\infty$.

In the meantime, this also shows that if $O(\mathbf{0},R) \subset V \Subset \mathbb{R}^3$, then a lower bound for $\|\hat{\mathcal{D}}+\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}$ will be given by $\|\hat{\mathcal{D}}+\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)} \geq \sqrt{2}kR/3$. In other words, the radii of the inscribed sphere and the circumscribed sphere of a volume V set lower and upper bounds for the operator norm $\|\hat{\mathcal{D}}+\hat{\gamma}\|_{L^2(V;\mathbb{C}^3)}$. ∎

Corollary 1.2 *If $\chi \in \mathbb{C}$ satisfies the inequality*

$$|\chi|^2 \min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\} < \operatorname{Im}\chi,$$

then the Born operator $\hat{\mathcal{B}} = \hat{I} - \chi\hat{\mathcal{G}} = \hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C) + i\chi\hat{\gamma}_S : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is invertible, with the norm estimate of the inverse operator given by

$$\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \le \frac{|\chi|}{\operatorname{Im}\chi}\left(1 - \frac{|\chi|^2}{\operatorname{Im}\chi}\min\left\{\frac{5}{8}kR_V, \frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, \frac{k^3}{4\pi}\iiint_V \mathrm{d}^3\boldsymbol{r}\right\}\right)^{-1}.$$

Proof Now that $\operatorname{Im}\chi > 0$ and $\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a Hermitian operator, we naturally have the resolvent estimate (see Ref. [40], p. 148):

$$\|[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}\|_{L^2(V;\mathbb{C}^3)} \le \frac{1}{|\chi\operatorname{Im}(1/\chi)|} = \frac{|\chi|}{\operatorname{Im}\chi}.$$

Meanwhile, we have

$$(\hat{I} - \chi\hat{\mathcal{G}})^{-1} = \{\hat{I} + [\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1} i\chi\hat{\gamma}_S\}^{-1}[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}$$

where

$$\|\{\hat{I} + [\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1} i\chi\hat{\gamma}_S\}^{-1}\|_{L^2(V;\mathbb{C}^3)} \le \left(1 - |\chi|\|[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}\right)^{-1}.$$

The claimed norm estimate for $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)}$ then emerges after we insert the upper bounds for $\|[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}\|_{L^2(V;\mathbb{C}^3)}$ and $\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)}$ into the right-hand side of the inequality $\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \le (1 - |\chi|\|[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}\|_{L^2(V;\mathbb{C}^3)}\|\hat{\gamma}_S\|_{L^2(V;\mathbb{C}^3)})^{-1}\|[\hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C)]^{-1}\|_{L^2(V;\mathbb{C}^3)}$. ∎

By interchanging the rôles of $\hat{\gamma}_S$ and $\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C$, we may establish a similar bound estimate as in the following corollary.

Corollary 1.3 *If $\chi \in \mathbb{C}$ satisfies the inequality*

$$|\chi|^2\left(1 + \min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, 3\sqrt{2}kR_V\right\}\right) < |\operatorname{Re}\chi|,$$

then the Born operator $\hat{\mathcal{B}} = \hat{I} - \chi\hat{\mathcal{G}} = \hat{I} - \chi(\hat{\mathcal{D}} - \hat{I} + \hat{\gamma}_C) + i\chi\hat{\gamma}_S : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is invertible, with the norm estimate of the inverse operator given by

$$\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \le \frac{|\chi|}{|\operatorname{Re}\chi|}\left[1 - \frac{|\chi|^2}{|\operatorname{Re}\chi|}\left(1 + \min\left\{\frac{3}{5\pi}\left(k^3\iiint_V \mathrm{d}^3\boldsymbol{r}\right)^{2/3}, 3\sqrt{2}kR_V\right\}\right)\right]^{-1}. \quad ∎$$

S1.3.5. Analytic Continuation of the Helmholtz Equation

To complete the uniqueness theorem (Theorem 1.2) in S1.1, we need an argument based on the unique analytic continuation of Helmholtz equation, as given in the following lemma.

Lemma 1.8 *If $\mathbb{R}^3 \setminus (V \cup \partial V)$ is connected, $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$, then the scattering field*[23]

$$\begin{aligned}\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r}) &:= \chi\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\mathrm{d}^3\boldsymbol{r}' \\ &= \chi k^2\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\mathrm{d}^3\boldsymbol{r}' + \chi\iiint_V \left[\nabla'\nabla'\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\boldsymbol{E}(\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}', \boldsymbol{r} \notin V \cup \partial V\end{aligned}$$

---

[23] Here, we note that $\boldsymbol{E}_{\mathrm{sc}} \in C^\infty(\mathbb{R}^3 \setminus (V \cup \partial V);\mathbb{C}^3)$, and differentiation under the integration sign is always permissible.

vanishes identically when the far-field scattering amplitudes vanish in all directions

$$\boldsymbol{f}(\boldsymbol{n}) := -\frac{\chi k^2}{4\pi}\boldsymbol{n}\times\left[\boldsymbol{n}\times\iiint_V \boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right]\equiv\boldsymbol{0},\quad\forall|\boldsymbol{n}|=1.$$

Proof Let $R_V := \min_{\boldsymbol{r}\in\mathbb{R}^3}\max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'-\boldsymbol{r}| = \max_{\boldsymbol{r}'\in V\cup\partial V}|\boldsymbol{r}'|$ be the radius of the circumscribed sphere centered at the origin $\boldsymbol{r}=\boldsymbol{0}$ of the coordinate system. For $|\boldsymbol{r}|>R_V$, we have the following Fourier-Bessel expansions

$$\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|} = -ik\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Y^*_{\ell m}(\theta',\phi')Y_{\ell m}(\theta,\phi)j_\ell(k|\boldsymbol{r}'|)h^{(2)}_\ell(k|\boldsymbol{r}|)$$

and

$$\nabla'\nabla'\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|} = -ik\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Y_{\ell m}(\theta,\phi)h^{(2)}_\ell(k|\boldsymbol{r}|)\nabla'\nabla'[Y^*_{\ell m}(\theta',\phi')j_\ell(k|\boldsymbol{r}'|)],$$

both converging uniformly and absolutely for $(\boldsymbol{r},\boldsymbol{r}')\in B\times(V\cup\partial V)\subset\mathbb{R}^6$ where $B\Subset\mathbb{R}^3\setminus(V\cup\partial V)$ is any compact subset. This is because the rate of convergence of the Fourier-Bessel expansions may be compared to $\sum_{\ell=1}^{\infty}\ell^N|z|^\ell<+\infty, |z|<1, N>0$, as

$$\lim_{\ell\to\infty}\frac{(2\ell+1)j_\ell(k|\boldsymbol{r}'|)h^{(2)}_\ell(k|\boldsymbol{r}|)k|\boldsymbol{r}|}{i}\left(\frac{|\boldsymbol{r}|}{|\boldsymbol{r}'|}\right)^\ell=1,\quad\left|\sum_{m=-\ell}^{\ell}Y^*_{\ell m}(\theta',\phi')Y_{\ell m}(\theta,\phi)\right|\le\frac{2\ell+1}{4\pi},\quad\max_{(\boldsymbol{r},\boldsymbol{r}')\in B\times(V\cup\partial V)}\frac{|\boldsymbol{r}'|}{|\boldsymbol{r}|}<1,$$

and the Clebsch-Gordon coefficients that express $\nabla'\nabla'[Y^*_{\ell m}(\theta',\phi')j_\ell(k|\boldsymbol{r}'|)]$ as linear combinations of $j_{\ell'}(k|\boldsymbol{r}'|)Y^*_{\ell'm'}(\theta',\phi'), |\ell-\ell'|\le 2, |m-m'|\le 2$ are of order $O(\ell^2)$.

This allows us to interchange the infinite sum with the integration in the representation of $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})$, and obtain a Fourier-Bessel expansion of the scattering field

$$\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})=\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}\boldsymbol{E}_{\ell m}Y_{\ell m}(\theta,\phi)h^{(2)}_\ell(k|\boldsymbol{r}|),\quad|\boldsymbol{r}|>R_V,\tag{1.36}$$

where the coefficients $\boldsymbol{E}_{\ell m}$ are completely determined by the far-field asymptotic behavior of $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})$ as

$$\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})\sim\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}\boldsymbol{E}_{\ell m}Y_{\ell m}(\theta,\phi)\frac{i^{\ell+1}e^{-ik|\boldsymbol{r}|}}{k|\boldsymbol{r}|}=\boldsymbol{f}(\boldsymbol{n}(\theta,\phi))\frac{e^{-ik|\boldsymbol{r}|}}{|\boldsymbol{r}|},\quad|\boldsymbol{r}|\to\infty$$

or

$$\boldsymbol{E}_{\ell m}=\lim_{|\boldsymbol{r}|\to+\infty}\frac{k|\boldsymbol{r}|e^{ik|\boldsymbol{r}|}}{i^{\ell+1}}\oint\!\!\!\oint\boldsymbol{E}_{\mathrm{sc}}(|\boldsymbol{r}|,\theta,\phi)Y^*_{\ell m}(\theta,\phi)\,\mathrm{d}\,\Omega=\frac{k}{i^{\ell+1}}\oint\!\!\!\oint\boldsymbol{f}(\boldsymbol{n}(\theta,\phi))Y^*_{\ell m}(\theta,\phi)\,\mathrm{d}\,\Omega.$$

Now that the far-field scattering amplitudes $\boldsymbol{f}(\boldsymbol{n})\equiv\boldsymbol{0}$ vanish in all directions $|\boldsymbol{n}|=1$, we obtain a vanishing scattering field $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})\equiv\boldsymbol{0}, |\boldsymbol{r}|>R_V$ in a non-empty open subset $\mathbb{R}^3\setminus\mathrm{Cl}\,(O(\boldsymbol{0},R_V))$ of the connected open set $\mathbb{R}^3\setminus(V\cup\partial V)$. We claim that this implies a vanishing scattering field $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$ in its entire domain. To see this, we may establish a spherical coordinate system $\boldsymbol{r}(|\boldsymbol{r}-\boldsymbol{r}_\star|,\theta_\star,\phi_\star)$ based on any point $\boldsymbol{r}_\star\in\mathbb{R}^3\setminus(V\cup\partial V)$. So long as the open ball $O(\boldsymbol{r}_\star,\varepsilon_\star)\subset\mathbb{R}^3\setminus(V\cup\partial V)$ lies outside the dielectric volume, we have $\min_{\boldsymbol{r}'\in\partial V}|\boldsymbol{r}'-\boldsymbol{r}_\star|>\varepsilon_\star$, so we can use

$$\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}=-ik\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Y^*_{\ell m}(\theta'_\star,\phi'_\star)Y_{\ell m}(\theta_\star,\phi_\star)j_\ell(k|\boldsymbol{r}-\boldsymbol{r}_\star|)h^{(2)}_\ell(k|\boldsymbol{r}'-\boldsymbol{r}_\star|)$$

and

$$\nabla'\nabla'\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}=-ik\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}Y_{\ell m}(\theta_\star,\phi_\star)j_\ell(k|\boldsymbol{r}-\boldsymbol{r}_\star|)\nabla'\nabla'[Y^*_{\ell m}(\theta'_\star,\phi'_\star)h^{(2)}_\ell(k|\boldsymbol{r}'-\boldsymbol{r}_\star|)],$$

to deduce a uniformly and absolutely convergent Fourier-Bessel expansion based on point $\boldsymbol{r}_\star$

$$\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})=\sum_{\ell=0}^{\infty}\sum_{m=-\ell}^{\ell}\boldsymbol{E}_{\ell m,\star}Y_{\ell m}(\theta_\star,\phi_\star)j_\ell(k|\boldsymbol{r}-\boldsymbol{r}_\star|),\quad\forall\boldsymbol{r}\in O(\boldsymbol{r}_\star,\varepsilon_\star),$$

where the coefficient $\boldsymbol{E}_{\ell m,\star}$ can also be uniquely determined by integration of the field $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})=\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r}_\star+\varepsilon'_\star\boldsymbol{n}),|\boldsymbol{n}|=1$ over a spherical surface $\partial O(\boldsymbol{r}_\star,\varepsilon'_\star)\subset O(\boldsymbol{r}_\star,\varepsilon_\star)$ such that $j_\ell(k\varepsilon'_\star)\neq0$:

$$\boldsymbol{E}_{\ell m,\star}=\frac{1}{j_\ell(k\varepsilon'_\star)}\oiint_{|\boldsymbol{n}|=1}\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r}_\star+\varepsilon'_\star\boldsymbol{n}(\theta,\phi))Y^*_{\ell m}(\theta,\phi)\,\mathrm{d}\,\Omega.$$

As the external volume $\mathbb{R}^3\setminus(V\cup\partial V)$ is a connected open set, any two points can be joined by an unbroken chain of piecewise line segments of finite total length. This chain of line segments, in turn, can be covered by an unbroken chain of spheres (where two consecutive members have non-empty intersection) that lies inside $\mathbb{R}^3\setminus(V\cup\partial V)$. In each of these spheres, we can construct a convergent Fourier-Bessel expansion of $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})$ based on the center of the sphere, thus forming an analytic continuation of the field $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})$ along the unbroken chain of spheres. Neighbor after neighbor, the Fourier-Bessel coefficients $\boldsymbol{E}_{\ell m,\star}$ based on any point $\boldsymbol{r}_\star\in\mathbb{R}^3\setminus(V\cup\partial V)$ can be related to a sphere that intersects the volume $\mathbb{R}^3\setminus\mathrm{Cl}\,(O(\boldsymbol{0},R_V))$. As $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})\equiv\boldsymbol{0},|\boldsymbol{r}|>R_V$, we can see that all the Fourier-Bessel coefficients $\boldsymbol{E}_{\ell m,\star}$ thus obtained have to vanish. Therefore, we have an identically vanishing scattering field $\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$ in the connected region outside the dielectric volume. ■

Remark The unique analytic continuation of the electric field satsifying the Helmholtz equation $(\nabla^2+k^2)\boldsymbol{E}_{\mathrm{sc}}(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V)$ is a specific case of the "unique continuation principle" (Theorem 8.6 in Ref. [14] with a more sophisticated proof) that applies to a broader class of partial differential equations $\nabla^2f(\boldsymbol{r})+k^2N(\boldsymbol{r})f(\boldsymbol{r})=0$, where $N(\boldsymbol{r})$ is continuous.

Lord Rayleigh once conjectured that the Fourier-Bessel expansion outside $O(\boldsymbol{0},R_V)$ (Eq. 1.36) would be applicable to the entire volume $\mathbb{R}^3\setminus(V\cup\partial V)$. If such a "Rayleigh hypothesis" were true, we would have a more direct way to conclude that the far-field scattering amplitudes determine the scattering field. However, the last decade saw strong numerical evidence against the Rayleigh hypothesis for light scattering on non-spherical particles — there is now a consensus (see Ref. [36], p. 40) that the Rayleigh hypothesis "could be wrong". As seen above, the proof of Lemma 1.8 does not have to rely on the Rayleigh hypothesis, and only draws on the connectedness of the open set $\mathbb{R}^3\setminus(V\cup\partial V)$.

The conclusion of this lemma is identical to that of Corollary 4.10 in Ref. [18], in which the authors provided a proof based on Taylor expansions in $1/|\boldsymbol{r}|$ as $\boldsymbol{E}_{sc}(\boldsymbol{r})=(e^{ik|\boldsymbol{r}|}/|\boldsymbol{r}|)\sum_{\ell=0}^{\infty}\boldsymbol{E}_\ell(\theta,\phi)/|\boldsymbol{r}|^\ell$ and the recursive relations for $\boldsymbol{E}_\ell(\theta,\phi),\ell=0,1,2,\dots$, without explicitly using the spherical Hankel function. ■

S1.3.6. Generalizations to Inhomogeneous and Anisotropic Media

Here, we briefly state some natural generalizations of Theorem 1.1 and Theorem 1.2 to inhomogeneous and anisotropic media, with the self-evident proofs omitted.

First, we mention some obvious extensions of the generalized optical theorem proved for homogeneous media (Theorem 1.1).

Corollary 1.4 (Generalized Optical Theorem — Inhomogeneous Medium) *Consider a spatial distribution of dielectric susceptibility* $\chi(\boldsymbol{r}),\boldsymbol{r}\in\mathbb{R}^3$ *that is bounded* $\sup_{\boldsymbol{r}\in\mathbb{R}^3}|\chi(\boldsymbol{r})|<+\infty$ *and has compact support* $\operatorname{supp}\chi:=\mathrm{Cl}\,\{\boldsymbol{r}\in\mathbb{R}^3|\chi(\boldsymbol{r})\neq0\}\Subset\mathbb{R}^3$*. Let the corresponding Born operator* $\hat{\mathcal{B}}:L^2(\operatorname{supp}\chi;\mathbb{C}^3)\longrightarrow L^2(\operatorname{supp}\chi;\mathbb{C}^3)$ *be defined by*

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r})=\boldsymbol{E}(\boldsymbol{r})+\frac{\boldsymbol{P}(\boldsymbol{r})}{\epsilon_0}+\frac{1}{\epsilon_0}\lim_{\varepsilon\to0^+}\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{P}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}\tag{1.37}$$

where $\tilde{\boldsymbol{P}}(\boldsymbol{q})=\epsilon_0\iiint_{\operatorname{supp}\chi}\chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r}$ *is the Fourier transform of the electric polarization* $\boldsymbol{P}(\boldsymbol{r})=\epsilon_0\chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})$*. Then, we have the following form of generalized optical theorem:*

$$\begin{aligned}\operatorname{Im}\langle\boldsymbol{E}-\hat{\mathcal{B}}\boldsymbol{E},\chi k^2\boldsymbol{E}\rangle&:=\operatorname{Im}\left\{k^2\iiint_{\operatorname{supp}\chi}\Big[\boldsymbol{E}(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r})\Big]^*\cdot\chi(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right\}\\&=\frac{k^5}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\iiint_{\operatorname{supp}\chi}\chi(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right|^2\mathrm{d}\,\Omega,\forall\boldsymbol{E}\in L^2(\operatorname{supp}\chi;\mathbb{C}^3).\end{aligned}$$ ■

Remark Here, we may consider a piecewise constant $\chi(\boldsymbol{r})$ that takes only finitely many different values. Suppose that $\chi(\boldsymbol{r})$ assumes a non-zero constant $\chi_{(j)}$ value in a bounded open set $V_{(j)}$, then we may still regard the Born operator $\hat{\mathcal{B}} : L^2(\operatorname{supp}\chi;\mathbb{C}^3) \longrightarrow L^2(\operatorname{supp}\chi;\mathbb{C}^3)$ defined in Eq. 1.37 as the continuous extension of $\hat{\mathcal{B}} : \bigcap_{(j)}[C^1(V_{(j)};\mathbb{C}^3)\cap L^2(V_{(j)};\mathbb{C}^3)] \longrightarrow [\bigcap_{(j)} C^2(V_{(j)};\mathbb{C}^3)]\cap L^2(\operatorname{supp}\chi;\mathbb{C}^3)$ that is defined by classical derivatives:

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) = [1+\chi(\boldsymbol{r})]\boldsymbol{E}(\boldsymbol{r}) - \nabla\times\nabla\times\iiint_{\mathbb{R}^3}\frac{\chi(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \forall\boldsymbol{E}\in\bigcap_{(j)}\left[C^1(V_{(j)};\mathbb{C}^3)\cap L^2(V_{(j)};\mathbb{C}^3)\right].$$

For a generic $\chi(\boldsymbol{r}), \boldsymbol{r}\in\mathbb{R}^3$, due to the possible lack of certain analytic attributes in $\chi(\boldsymbol{r})$, the Born equation as in Eq. 1 may not be a well-defined differo-integral equation in terms of classical derivatives. In this case, we may regard Eq. 1.37 as an axiomatic generalization of the Born operator to a bounded and compactly supported spatial distribution of dielectric susceptibility $\chi(\boldsymbol{r}), \boldsymbol{r}\in\mathbb{R}^3$. $\blacksquare$

In the above, we assumed that the electric polarization $\boldsymbol{P}(\boldsymbol{r})$ is pointwise parallel to the electric field $\boldsymbol{E}(\boldsymbol{r})$, and the proportionality coefficient $\epsilon_0\chi(\boldsymbol{r})$ is a scalar field. This is a correct assumption for isotropic media. For anisotropic media, we generally have $\boldsymbol{P}(\boldsymbol{r}) = \epsilon_0\overleftrightarrow{\chi}(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})$, where the dielectric susceptibility tensor $\overleftrightarrow{\chi}(\boldsymbol{r}) = \{\chi_{ij}(\boldsymbol{r})\}_{i,j=1}^3$ is a 3×3 matrix. Even in this case, we can use the momentum space method to arrive at a generalized optical theorem with ease.

Corollary 1.5 (Generalized Optical Theorem — Anisotropic Medium) *Consider a spatial distribution of dielectric susceptibility tensor* $\overleftrightarrow{\chi}(\boldsymbol{r}), \boldsymbol{r}\in\mathbb{R}^3$ *that is bounded* $\sup_{\boldsymbol{r}\in\mathbb{R}^3}\max_{1\leq i\leq j\leq 3}|\chi_{ij}(\boldsymbol{r})| < +\infty$ *and has compact support* $\operatorname{supp}\overleftrightarrow{\chi} := \operatorname{Cl}\{\boldsymbol{r}\in\mathbb{R}^3|\overleftrightarrow{\chi}(\boldsymbol{r})\neq\overleftrightarrow{0}\}\Subset\mathbb{R}^3$. *Let the corresponding Born operator* $\hat{\mathcal{B}} : L^2(\operatorname{supp}\overleftrightarrow{\chi};\mathbb{C}^3)\longrightarrow L^2(\operatorname{supp}\overleftrightarrow{\chi};\mathbb{C}^3)$ *be defined by*

$$(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r}) = \boldsymbol{E}(\boldsymbol{r}) + \frac{\boldsymbol{P}(\boldsymbol{r})}{\epsilon_0} + \frac{1}{\epsilon_0}\lim_{\varepsilon\to 0^+}\frac{1}{(2\pi)^3}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{P}}(\boldsymbol{q})]}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{q}$$

where $\tilde{\boldsymbol{P}}(\boldsymbol{q}) = \epsilon_0\iiint_{\operatorname{supp}\overleftrightarrow{\chi}}\overleftrightarrow{\chi}(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^3\boldsymbol{r}$. *Then, we have the following form of generalized optical theorem:*

$$\begin{aligned}\operatorname{Im}\langle\boldsymbol{E}-\hat{\mathcal{B}}\boldsymbol{E},\overleftrightarrow{\chi}k^2\boldsymbol{E}\rangle &:= \operatorname{Im}\left\{k^2\iiint_{\operatorname{supp}\overleftrightarrow{\chi}}\Big[\boldsymbol{E}(\boldsymbol{r})-(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r})\Big]^*\cdot\overleftrightarrow{\chi}(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}\right\}\\ &= \frac{k^5}{16\pi^2}\oiint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\iiint_{\operatorname{supp}\overleftrightarrow{\chi}}\overleftrightarrow{\chi}(\boldsymbol{r}')\boldsymbol{E}(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right|^2\mathrm{d}\,\Omega, \forall\boldsymbol{E}\in L^2(\operatorname{supp}\overleftrightarrow{\chi};\mathbb{C}^3).\end{aligned}$$ $\blacksquare$

Next, we point out that the uniqueness theorem proved for dissipative homogeneous media (Theorem 1.2, situation (A)) admits an immediate generalization to inhomogeneous media where $\sup_{\boldsymbol{r}\in\operatorname{supp}\chi}\operatorname{Im}\chi(\boldsymbol{r}) < 0$, and to anisotropic media where $\sup_{\boldsymbol{r}\in\operatorname{supp}\overleftrightarrow{\chi}}\lambda_{\max}[\operatorname{Im}\overleftrightarrow{\chi}(\boldsymbol{r})] < 0$. Here, $\lambda_{\max}[\operatorname{Im}\overleftrightarrow{\chi}(\boldsymbol{r})]$ stands for the maximum eigenvalue of the real-valued and symmetric matrix $\operatorname{Im}\overleftrightarrow{\chi}(\boldsymbol{r})$.

Nonetheless, it should be noted that situation (B) in Theorem 1.2 for non-dissipative homogeneous dielectric media does not allow direct generalizations to inhomogeneous dielectric. In other words, the topological condition about the connectedness of $\mathbb{R}^3\setminus(V\cup\partial V)$ will no longer be conducive to the uniqueness of the solution. On pp. 259-264 of Ref. [14], D. Colton and R. Kress discussed the non-uniqueness of the solution to electromagnetic scattering problem in a spherically stratified sphere, in which case $\chi(\boldsymbol{r}) = \chi(|\boldsymbol{r}|)$ only depends on the radial distance $|\boldsymbol{r}|$ and $\operatorname{Im}\chi(\boldsymbol{r}) = 0$ inside the inhomogeneous dielectric sphere.

In S1.2.2, we demonstrated the existence theorem for the light scattering problem in homogeneous media using the Riesz-Fredholm theory for compact perturbations of invertible mappings. It should also be noted that the existence theorems for certain types of inhomogeneous dielectric media have also been established in the literature using the Riesz-Fredholm theory. When the susceptibility $\chi(\boldsymbol{r})$ is Lipschitz-Hölder differentiable in the entire space, satisfying $|\nabla\chi(\boldsymbol{r})-\nabla'\chi(\boldsymbol{r}')|\leq L_1|\boldsymbol{r}-\boldsymbol{r}'|^\alpha, \forall\boldsymbol{r},\boldsymbol{r}'\in\mathbb{R}^3$ $(0<\alpha\leq 1)$, the Fredholm operator in question can be resolved into the form of "identity + compact", as shown in Chapter 9 of Ref. [14]. When the susceptibility $\chi(\boldsymbol{r})$ is merely Lipschitz-Hölder continuous in the entire space, satisfying $|\chi(\boldsymbol{r})-\chi(\boldsymbol{r}')|\leq L_0|\boldsymbol{r}-\boldsymbol{r}'|^\alpha, \forall\boldsymbol{r},\boldsymbol{r}'\in\mathbb{R}^3$ $(0<\alpha\leq 1)$, the Riesz-Fredholm theory can still be carried out by considering the "symbol of a Hölder continuous multiplier", as shown in Ref. [15].

S2. EVOLUTION SEMIGROUPS FOR ELECTROMAGNETIC SCATTERING

From now on, we will mainly concentrate on light scattering problems for homogeneous medium occupying a bounded and open volume, with dielectric susceptibility χ satisfying $\operatorname{Im}\chi\leq 0$.

In S2, we refine the spectral analysis of the Born equation from the perspective of evolution semigroups. In S2.1, we start by representing the solution to the Born equation using evolution semigroups (S2.1.1) and then proceed to demonstrate that the uniqueness theorem of the Born equation can be extended to the special point $\chi = -1$ under reasonable geometric and physical constraints (S2.1.2). In S2.2, we will prove a surprising extension of the existence theorem of the Born equation to the half-line $\chi \leq -1$ excluding a critical point $\chi = -2$, where the electromagnetic enhancement becomes unbounded.

S2.1. Integral Representation of the Solution to Light Scattering

S2.1.1. Evolution Semigroups, Mean Ergodicity, and Fixed Points

Proposition 2.1 (Integral Representation of the Solution) *The solution to the Born equation* $(\hat{I} - \chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ *has the following integral representation for* $\operatorname{Im}\chi < 0$*:*

$$\boldsymbol{E} = (\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}} = \frac{1}{i\chi}\int_0^{+\infty} e^{i\tau/\chi}\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{E}_{\text{inc}}\,\mathrm{d}\,\tau \tag{2.1}$$

where $\{\exp(-i\tau\hat{\mathcal{G}}) := \sum_{s=0}^{\infty}(-i\tau\hat{\mathcal{G}})^s/s!|\tau \geq 0\}$ *is an evolution semigroup*[24] *with infinitesimal generator* $-i\hat{\mathcal{G}}$.

Proof From the existence theorem, we know that for every $\lambda' > 0$, the operator $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1} : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is a well-defined bounded linear mapping, satisfying $\|\lambda'(\lambda'\hat{I} + i\hat{\mathcal{G}})^{-1}\|_{L^2(V;\mathbb{C}^3)} \leq 1$.[25] Using the Hille-Yosida theorem (see Refs. [58, 59], as well as Ref. [44], Chap. IX, Ref. [56], p. 249 and Ref. [60], pp. 73-75), we may thus conclude that $\{\exp(-i\tau\hat{\mathcal{G}})|\tau \geq 0\}$ is a contraction semigroup, which satisfies $\|\exp(-i\tau\hat{\mathcal{G}})\|_{L^2(V;\mathbb{C}^3)} \leq 1, \forall\tau \geq 0$, and this semigroup can be related to the resolvent by the following Bochner-Dunford integral: [61, 62, 63]

$$(\lambda'\hat{I} + i\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}} = \int_0^{+\infty} e^{-\lambda'\tau}\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{E}_{\text{inc}}\,\mathrm{d}\,\tau, \quad \text{where } \operatorname{Re}\lambda' > 0.$$

Now, setting $\lambda = -i/\chi$, one obtains

$$(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}} = \frac{1}{i\chi}\int_0^{+\infty} e^{i\tau/\chi}\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{E}_{\text{inc}}\,\mathrm{d}\,\tau, \quad \text{where } \operatorname{Re}(i\chi) = \operatorname{Re}\lambda' > 0. \quad \blacksquare$$

Remark Here, if we define $\boldsymbol{\psi}_\tau(\boldsymbol{r}) \equiv \boldsymbol{\psi}(\boldsymbol{r};\tau) := (\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{E}_{\text{inc}})(\boldsymbol{r})$, then we have the differential equation

$$i\frac{\partial\boldsymbol{\psi}(\boldsymbol{r};\tau)}{\partial\tau} = \hat{\mathcal{G}}\boldsymbol{\psi}(\boldsymbol{r};\tau) = \nabla\times\nabla\times\iiint_V \frac{\boldsymbol{\psi}(\boldsymbol{r}';\tau)e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' - \boldsymbol{\psi}(\boldsymbol{r};\tau) \tag{2.2}$$

with initial condition $\boldsymbol{\psi}_0(\boldsymbol{r}) \equiv \boldsymbol{\psi}(\boldsymbol{r};0) = \boldsymbol{E}_{\text{inc}}(\boldsymbol{r})$. By the generalized optical theorem, it is clear that we have the evolution equation for $\langle\boldsymbol{\psi}_\tau,\boldsymbol{\psi}_\tau\rangle_V = \|\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{E}_{\text{inc}}\|^2_{L^2(V;\mathbb{C}^3)}$ as follows:

$$\frac{\mathrm{d}}{\mathrm{d}\,\tau}\langle\boldsymbol{\psi}_\tau,\boldsymbol{\psi}_\tau\rangle_V = -2\operatorname{Im}\left\langle i\frac{\partial\boldsymbol{\psi}_\tau}{\partial\tau},\boldsymbol{\psi}_\tau\right\rangle_V = -2\operatorname{Im}\langle\hat{\mathcal{G}}\boldsymbol{\psi}_\tau,\boldsymbol{\psi}_\tau\rangle_V = -\frac{k^3}{8\pi^2}\oint\!\!\!\oint_{|\boldsymbol{n}|=1}\left|\boldsymbol{n}\times\iiint_V \boldsymbol{\psi}_\tau(\boldsymbol{r}')e^{ik\boldsymbol{n}\cdot\boldsymbol{r}'}\,\mathrm{d}^3\boldsymbol{r}'\right|^2\mathrm{d}\,\Omega \leq 0. \tag{2.3}$$

We note that the monotonic decay in the amplitude $\|\exp(\tau\hat{\mathcal{A}})\boldsymbol{F}\|_{\mathfrak{H}}$ of the trajectory $\{\exp(\tau\hat{\mathcal{A}})\boldsymbol{F},\tau \geq 0|\boldsymbol{F} \in \mathfrak{H}\}$ is a universal property for all contraction semigroups $\{\exp(\tau\hat{\mathcal{A}})|\tau \geq 0\}$ acting on a Hilbert space $\mathfrak{H}$, because in a

---

[24] In our context, the variable τ is dimensionless and has nothing to do with the elapse of physical time, so the term "evolution" has only a formal meaning. In general, the evolution semigroup $\{\exp(-i\tau\hat{\mathcal{G}})|\tau \geq 0\}$ acting on $L^2(V;\mathbb{C}^3)$ may also bear little resemblance to the Lax-Phillips evolution semigroup $\{\hat{Z}(t) = \hat{P}_+\hat{U}(t)|t \geq 0\}$ acting on a generic Hilbert space $\mathfrak{H}$ in quantum scattering theory (see for example, Ref. [56], Chaps. I, III and V, or Ref. [57], Chap. 9), where $\hat{U}(t)$ is a unitary transformation, and $\hat{P}_+$ an orthogonal projection. The major difference is that the strong stability result $\lim_{t\to+\infty}\langle\hat{Z}(t)\boldsymbol{X},\hat{Z}(t)\boldsymbol{X}\rangle_{\mathfrak{H}} = 0$ holds for all $\boldsymbol{X} \in \mathfrak{H}$ in quantum scattering theory (see Ref. [56], p. 12), while the validity of $\lim_{\tau\to+\infty}\langle\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F},\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}\rangle_V = 0$ hinges on some subtle qualifications of $\boldsymbol{F}$ and V (see Theorem 2.1).

[25] Hereafter, we juxtapose λ' with $-i\hat{\mathcal{G}}$ and reserve the symbol $\lambda = i\lambda'$ for the spectral analysis of the operator $\hat{\mathcal{G}}$.

Hilbert space, the Hille-Yosida condition $\|\lambda'(\lambda'\hat{I}+\hat{\mathcal{A}})^{-1}\|_{\mathfrak{H}}\leq 1,\lambda'>0$ is equivalent to the dissipation condition $\langle\hat{\mathcal{A}}\boldsymbol{F},\boldsymbol{F}\rangle_{\mathfrak{H}}+\langle\boldsymbol{F},\hat{\mathcal{A}}\boldsymbol{F}\rangle_{\mathfrak{H}}\leq 0,\forall\boldsymbol{F}\in\mathfrak{H}$, according to the Lumer-Phillips theorem (see Ref. [64] or Ref. [60], pp. 83-84). In the context of light scattering, the dissipation condition follows directly from the generalized optical theorem (Theorem 1.1). ∎

Proposition 2.2 (Mean Ergodicity) *So long as the volume V is open and bounded, we have the following limit relations*

$$\lim_{T\to+\infty}\frac{1}{T}\int_0^T\exp(-i\tau(\hat{\mathcal{G}}+\hat{I}))\boldsymbol{F}\,\mathrm{d}\,\tau=\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F},\quad\forall\boldsymbol{F}\in L^2(V;\mathbb{C}^3)$$

where the operator $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}$ projects a square-integrable vector field to the kernel space

$$\ker(\hat{\mathcal{G}}+\hat{I})=\left\{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\left|\|(\hat{\mathcal{G}}+\hat{I})\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0,\ \textit{where}\ ((\hat{\mathcal{G}}+\hat{I})\boldsymbol{E})(\boldsymbol{r}):=\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right.\right\}.$$

If $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected and $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$, then we further have

$$\lim_{T\to+\infty}\frac{1}{T}\int_0^T\exp(-i\tau(\hat{\mathcal{G}}-\lambda\hat{I}))\boldsymbol{F}\,\mathrm{d}\,\tau=\boldsymbol{0},\quad\forall\boldsymbol{F}\in L^2(V;\mathbb{C}^3),\quad\forall\lambda\in(-\infty,-1)\cup(-1,+\infty).$$

Proof We note that for the bounded strongly continuous semigroups $\exp(\tau\hat{\mathcal{A}})$ (where $\hat{\mathcal{A}}=-i(\hat{\mathcal{G}}-\lambda\hat{I})$ for some real number $\lambda\in\mathbb{R}$) acting on a Hilbert space, the mean ergodicity theorem holds (see Ref. [60], pp. 338-340). Accordingly, as $T\to+\infty$, the Cesàro mean $\frac{1}{T}\int_0^T\exp(\tau\hat{\mathcal{A}})\boldsymbol{F}\,\mathrm{d}\,\tau$ converges to the projection $\hat{P}_{\ker(\hat{\mathcal{A}})}\boldsymbol{F}$, where $\ker(\hat{\mathcal{A}})$ is the kernel space of the infinitesimal generator $\hat{\mathcal{A}}$. In other words, the limit of the Cesàro mean $\lim_{T\to+\infty}\frac{1}{T}\int_0^T\exp(\tau\hat{\mathcal{A}})\boldsymbol{F}\,\mathrm{d}\,\tau=\hat{P}_{\ker(\hat{\mathcal{A}})}\boldsymbol{F}\in\ker(\hat{\mathcal{A}})$ is a fixed point of the entire semigroup: $\exp(\tau\hat{\mathcal{A}})\hat{P}_{\ker(\hat{\mathcal{A}})}\boldsymbol{F}=\hat{P}_{\ker(\hat{\mathcal{A}})}\boldsymbol{F},\forall\tau\geq 0$.

According to Proposition 1.5, so long as the volume V is open and bounded, the kernel space $\ker(\hat{\mathcal{G}}+\hat{I})$ is infinite-dimensional, thus the projection operator $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}:L^2(V;\mathbb{C}^3)\longrightarrow\ker(\hat{\mathcal{G}}+\hat{I})$ is non-trivial. According to Proposition 1.6, if $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected and $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$, the kernel space $\ker(\hat{\mathcal{G}}-\lambda\hat{I})=\{\boldsymbol{0}\}$ is zero-dimensional for $\lambda\in(-\infty,-1)\cup(-1,+\infty)$, hence the associated projection operator $\hat{P}_{\ker(\hat{\mathcal{G}}-\lambda\hat{I})}:L^2(V;\mathbb{C}^3)\longrightarrow\{\boldsymbol{0}\}$ trivializes to an annihilation. ∎

From the mean ergodicity result, it is clear that the the kernel space $\ker(\hat{\mathcal{G}}+\hat{I})$ (i.e. the totality of fixed points of the entire semigroup $\{\exp(-i\tau(\hat{\mathcal{G}}+\hat{I}))|\tau\geq 0\}$) calls for special attention.

Proposition 2.3 (Characterization of Fixed Points) *So long as the volume V is open and bounded, we have the following set inclusion relation*

$$\ker(\hat{\mathcal{G}}+\hat{I})\supset\{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\boldsymbol{0},\forall\boldsymbol{q}\in\mathbb{R}^3\};$$

if $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected and $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$, then we further have the identity

$$\ker(\hat{\mathcal{G}}+\hat{I})=\{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\boldsymbol{0},\forall\boldsymbol{q}\in\mathbb{R}^3\}.$$

Proof First, we note that so long as the volume $V\subset\mathbb{R}^3$ is open and bounded, the set inclusion relation $\ker(\hat{\mathcal{G}}+\hat{I})\supset\{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\boldsymbol{0},\forall\boldsymbol{q}\in\mathbb{R}^3\}$ is an immediate consequence of the Fourier inversion formula

$$((\hat{\mathcal{G}}+\hat{I})\boldsymbol{E})(\boldsymbol{r}):=-\frac{1}{(2\pi)^3}\lim_{\varepsilon\to 0^+}\iiint_{\mathbb{R}^3}\frac{\boldsymbol{q}\times[\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})]e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}\,\mathrm{d}^3\boldsymbol{q}.$$

Then, we can see that if $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected, the boundary has zero volume $\iiint_{\partial V}\mathrm{d}^3\boldsymbol{r}=0$, and $\|(\hat{\mathcal{G}}+\hat{I})\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$, Eq. 1.21 in the proof of situation (B) in the uniqueness theorem (Theorem 1.2) may be adapted for the $\chi=-1$ case, thus giving rise to

$$|\boldsymbol{q}|^2\frac{\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-k^2}=\boldsymbol{0},\quad\forall\boldsymbol{q}\in\mathbb{R}^3\setminus V_0'',\quad\text{where}\ \iiint_{V_0''}\mathrm{d}^3\boldsymbol{q}=0.$$

As $\tilde{\boldsymbol{E}}\in C^\infty(\mathbb{R}^3;\mathbb{C}^3)$ is continuous in $\boldsymbol{q}$, we may extend the equality $\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\boldsymbol{0}$ to all $\boldsymbol{q}\in\mathbb{R}^3$. This proves the reverse set inclusion relation $\ker(\hat{\mathcal{G}}+\hat{I})\subset\{\boldsymbol{E}\in L^2(V;\mathbb{C}^3)|\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\boldsymbol{0},\forall\boldsymbol{q}\in\mathbb{R}^3\}$. ∎

While $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}$ does not necessarily vanish for an arbitrarily given square-integrable field, we will show in S2.1.2 that $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}=\boldsymbol{0}$ is true for all *transverse vector fields* $\boldsymbol{F}\in T(V;\mathbb{C}^3):=L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$ defined in a broad class of dielectric shapes. Given the indispensability of transversality in the physical formulation of the light scattering problem, we will highlight the Hilbert subspace $\Phi(V;\mathbb{C}^3):=\operatorname{Cl}(T(V;\mathbb{C}^3))$ as the totality of *physically admissible vector fields.* This Hilbert subspace $\Phi(V;\mathbb{C}^3)$ forms the totality of vector fields that can be approximated, in L^2-norm, by a sequence of transverse vector fields in the function space $T(V;\mathbb{C}^3)=L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$.

S2.1.2. Evolution of Physically Admissible Vector Fields — Extended Uniqueness Theorem

Here, we will present a detailed investigation of the evolution of a physically admissible vector field $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$ under the action of the semigroup $\{\exp(-i\tau\hat{\mathcal{G}})=\sum_{s=0}^{\infty}(-i\tau\hat{\mathcal{G}})^s/s!|\tau\geq 0\}$.

To start, we point out that it is meaningful to narrow down the integral representation of the solution (Eq. 2.1) to the Hilbert subspace $\Phi(V;\mathbb{C}^3)$, as the following two lemmata will reveal that $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1}\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3)$ whenever $\operatorname{Re}\lambda'>0$, and $\exp(-i\tau\hat{\mathcal{G}})\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3),\forall\tau\geq 0$.

Lemma 2.1 *For any bounded and open volume V, the operator $\hat{\mathcal{G}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ is an endomorphism, satisfying $\hat{\mathcal{G}}\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3)$. Accordingly, the semigroup action* $\exp(-i\tau\hat{\mathcal{G}}):\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ *is also an endomorphism, satisfying* $\exp(-i\tau\hat{\mathcal{G}})\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3),\forall\tau\geq 0$.

Proof First, we show that $\hat{\mathcal{G}}(T(V;\mathbb{C}^3))\subset T(V;\mathbb{C}^3)$.

For any $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)$, we define

$$(\hat{\mathcal{C}}\boldsymbol{E})(\boldsymbol{r}):=\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}',$$

as in S1.0.2. We may use integration by parts to verify that $\hat{\mathcal{C}}\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)$, and $(\nabla^2+k^2)(\hat{\mathcal{C}}\boldsymbol{E})(\boldsymbol{r})=-\boldsymbol{E}(\boldsymbol{r}),\forall\boldsymbol{r}\in V$. Therefore, we have $\hat{\mathcal{G}}\boldsymbol{E}\in L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)$. To demonstrate the transversality condition $\hat{\mathcal{G}}\boldsymbol{E}\in\ker(\nabla\cdot)$, it would suffice to compute $\nabla\cdot\nabla\times\nabla\times(\hat{\mathcal{C}}\boldsymbol{E})=0$.

Now that we have confirmed $\hat{\mathcal{G}}(T(V;\mathbb{C}^3))\subset T(V;\mathbb{C}^3)$, and the continuity of $\hat{\mathcal{G}}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ entails that $\hat{\mathcal{G}}(\operatorname{Cl}(\mathfrak{A}))\subset\operatorname{Cl}(\hat{\mathcal{G}}(\mathfrak{A}))$ for any subset $\mathfrak{A}\subset L^2(V;\mathbb{C}^3)$, we have $\hat{\mathcal{G}}\operatorname{Cl}(T(V;\mathbb{C}^3))\subset\operatorname{Cl}(\hat{\mathcal{G}}(T(V;\mathbb{C}^3)))\subset\operatorname{Cl}(T(V;\mathbb{C}^3))=\Phi(V;\mathbb{C}^3)$, as claimed. Furthermore, as the infinitesimal generator $-i\hat{\mathcal{G}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ is a continuous endomorphism, it generates a semigroup that satisfies $\exp(-i\tau\hat{\mathcal{G}})\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3),\forall\tau\geq 0$. ▮

Lemma 2.2 *For any bounded and open volume V, the operator $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ is an endomorphism for* $\operatorname{Re}\lambda'>0$, *satisfying $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1}\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3)$.*

Proof We will prove, *a fortiori*, that $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1}\Phi(V;\mathbb{C}^3)=\Phi(V;\mathbb{C}^3)$.

Consider the operator $\lambda'\hat{I}+i\hat{\mathcal{G}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$, which is a well-defined endomorphism between Hilbert spaces, owing to the fact that $\hat{\mathcal{G}}\Phi(V;\mathbb{C}^3)\subset\Phi(V;\mathbb{C}^3)$. Now, one may apply the spectral analysis to the Hilbert space $\Phi(V;\mathbb{C}^3)$, and decompose the complex plane into $\mathbb{C}=\rho^\Phi(\hat{\mathcal{G}})\cup\sigma_p^\Phi(\hat{\mathcal{G}})\cup\sigma_c^\Phi(\hat{\mathcal{G}})\cup\sigma_r^\Phi(\hat{\mathcal{G}})$. As the uniqueness theorem (situation (A) in Theorem 1.2) necessarily holds on $\Phi(V;\mathbb{C}^3)$, which is a subspace of $L^2(V;\mathbb{C}^3)$, we have $\lambda=i\lambda'\notin\sigma_p^\Phi(\hat{\mathcal{G}})$ for $\operatorname{Re}\lambda'>0$.

Meanwhile, it can be verified that $\hat{I}-\hat{\mathcal{D}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ remains a positive-semidefinite operator satisfying $\sigma^\Phi(\hat{I}-\hat{\mathcal{D}})\subset[0,1]$, and $\hat{\gamma}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ is a compact endomorphism. Therefore, the Riesz-Fredholm argument is still valid for the operator $\lambda'\hat{I}+i\hat{\mathcal{G}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$, $\lambda=i\lambda'\notin[0,1]$, leading to $i\lambda'\in\rho^\Phi(\hat{\mathcal{G}})$ for $\operatorname{Re}\lambda'>0$, i.e. $(\lambda'\hat{I}+i\hat{\mathcal{G}})^{-1}\Phi(V;\mathbb{C}^3)=\Phi(V;\mathbb{C}^3)$. ▮

Remark The same argument in this proof can be extended to confirm the relation $\rho(\hat{\mathcal{G}})\setminus[0,1]\subset\rho^\Phi(\hat{\mathcal{G}})\setminus[0,1]$. Meanwhile, it is also evident that $\sigma_p(\hat{\mathcal{G}})\supset\sigma_p^\Phi(\hat{\mathcal{G}})$, as an eigenvector in the function space $L^2(V;\mathbb{C}^3)$ may not belong to $\Phi(V;\mathbb{C}^3)$. Generally speaking, the Born equation $(\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E}=\boldsymbol{E}_{\text{inc}}$ with a physically admissible input $\boldsymbol{E}_{\text{inc}}\in\Phi(V;\mathbb{C}^3)$ is solvable for a wider range of χ values than the Born equation with a mere energetically admissible (square-integrable) input $\boldsymbol{E}_{\text{inc}}\in L^2(V;\mathbb{C}^3)$. ▮

Hereafter, our major interest will be devoted to *smooth dielectrics* satisfying the following topological and geometric requirements:

(a) The dielectric volume $V = \bigcup_{j=1}^{N_V} V_j \Subset \mathbb{R}^3$ is a bounded open set with finitely many connected components $V_1, \dots, V_{N_V}$ ($N_V < +\infty$);

(b) Each connected component V_j has a smooth boundary surface ∂V_j;[26]

(c) The dielectric boundary $\partial V = \bigsqcup_{j=1}^{N_V} \partial V_j$ is the union of mutually disjoint subsets $\partial V_1, \dots, \partial V_{N_V}$ satisfying $\partial V_j \cap \partial V_{j'} = \emptyset, 1 \leq j < j' \leq N_V$.

Proposition 2.4 *For a smooth dielectric with connected exterior volume* $\mathbb{R}^3 \setminus (V \cup \partial V)$, *we have* $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})} \boldsymbol{F} = \boldsymbol{0}$ *for all* $\boldsymbol{F} \in \Phi(V; \mathbb{C}^3)$.

Proof There are two immediate consequences of the geometric and topological conditions set forth in this proposition. First, as the compact smooth surface ∂V_j has zero volume (see Corollary 2.3 in S2.3.1), we have $\iiint_{\partial V} \mathrm{d}^3\boldsymbol{r} = \sum_{j=1}^{N_V} \iiint_{\partial V_j} \mathrm{d}^3\boldsymbol{r} = 0$. Hence the smooth boundary surface ∂V must have zero volume, and we may arrive at the identity $\ker(\hat{\mathcal{G}} + \hat{I}) = \{\boldsymbol{E} \in L^2(V; \mathbb{C}^3) | \boldsymbol{q} \times \tilde{\boldsymbol{E}}(\boldsymbol{q}) = \boldsymbol{0}, \forall \boldsymbol{q} \in \mathbb{R}^3\}$ according to Proposition 2.3. Second, with a connected exterior volume $\mathbb{R}^3 \setminus (V \cup \partial V)$ and mutually disjoint smooth boundary surfaces of distinct components $\partial V_j \cap \partial V_{j'} = \emptyset, 1 \leq j < j' \leq N_V < +\infty$, we may verify that each connected component V_j has a connected boundary ∂V_j, using the Alexander relation which counts the number of connected components. (See Ref. [65], pp. 384-387 for an analytic proof based on potential theory, or Lemma 2.3 in S2.3.2 for an algebraic proof based on the Mayer-Vietoris sequence in the de Rham cohomology.)

Now, for any $\boldsymbol{E} \in \ker(\hat{\mathcal{G}}+\hat{I})$, we smooth it out as $\boldsymbol{E}_{(\delta)}(\boldsymbol{r}) = \iiint_V \boldsymbol{E}(\boldsymbol{r}') \varkappa_\delta(\boldsymbol{r}-\boldsymbol{r}') \, \mathrm{d}^3\boldsymbol{r}'$ using the mollifier convolution kernel $\varkappa_\delta$ (see Ref. [66], p. 36) given by

$$\varkappa_\delta(\boldsymbol{r}) = \frac{\varkappa(\boldsymbol{r}/\delta)}{\delta^3}, \quad \text{with} \quad \varkappa(\boldsymbol{r}) = \begin{cases} \frac{\exp[-(1-|\boldsymbol{r}|^2)^{-1}]}{\iiint_{|\boldsymbol{r}|<1} \exp[-(1-|\boldsymbol{r}|^2)^{-1}] \, \mathrm{d}^3\boldsymbol{r}}, & |\boldsymbol{r}| < 1; \\ 0, & |\boldsymbol{r}| \geq 1. \end{cases} \tag{2.4}$$

Here, the positive parameter δ is chosen to be suitably small as compared to the spacings between different connected components $V_j, j = 1, \dots, N_V$. Concretely speaking, by choosing $6\delta \leq \delta_0 := \min\{\delta_V, \min_{1 \leq j < j' \leq N_V} \min_{\boldsymbol{r} \in \partial V_j, \boldsymbol{r}' \in \partial V_{j'}} |\boldsymbol{r} - \boldsymbol{r}'|\}$, we may ensure that $\operatorname{supp} \boldsymbol{E}_{(\delta)} \Subset V^+_{(2\delta)} := V \cup \{\boldsymbol{r} + \varepsilon \boldsymbol{n}(\boldsymbol{r}) | \boldsymbol{r} \in \partial V, 0 \leq \varepsilon < 2\delta\}$. Here, the "swollen volume" $V^+_{(2\delta)}$ is constructed by expansion along the outward normal vectors $\boldsymbol{n}(\boldsymbol{r})$ of ∂V, and the parameter $\delta_V > 0$ is determined by the geometry of the boundary surface ∂V (see the proof of Proposition 2.6 in S2.3.1). Using a corollary of the tubular neighborhood theorem (Corollary 2.4 in S2.3.1), one can verify that $V^+_{(2\delta)}$ has N_V connected components $V_j \cup \{\boldsymbol{r} + \varepsilon \boldsymbol{n}(\boldsymbol{r}) | \boldsymbol{r} \in \partial V_j, 0 \leq \varepsilon < 2\delta\}, j = 1, 2, \dots, N_V$, with their respective boundaries given by $\{\boldsymbol{r} + 2\delta \boldsymbol{n}(\boldsymbol{r}) | \boldsymbol{r} \in \partial V_j\}, j = 1, 2, \dots, N_V$, which are all smooth and connected surfaces.

Hereafter, we may assume that $N_V = 1$, without loss of generality. Now that we have $\boldsymbol{E}_{(\delta)} \in C_0^\infty(V^+_{(2\delta)}; \mathbb{C}^3)$, and $\nabla \times \boldsymbol{E}_{(\delta)} = \boldsymbol{0}$ holds on a connected open set $V^+_{(2\delta)}$ with smooth and connected boundary $\partial V^+_{(2\delta)}$, we can always pick a scalar potential with compact support $f_{(\delta)} \in C_0^\infty(V^+_{(2\delta)}; \mathbb{C})$ such that $\boldsymbol{E}_{(\delta)}(\boldsymbol{r}) = \nabla f_{(\delta)}(\boldsymbol{r}), \forall \boldsymbol{r} \in V^+_{(2\delta)}$. The existence of such a scalar potential is guaranteed by the Poincaré duality relation between the de Rham cohomology and the Alexander-Spanier cohomology, as explained by Proposition 2.7 in S2.3.2.

For any $\boldsymbol{F} \in T(V; \mathbb{C}^3) = L^2(V; \mathbb{C}^3) \cap C^\infty(V; \mathbb{C}^3) \cap \ker(\nabla \cdot)$, and $\boldsymbol{E} \in \ker(\hat{\mathcal{G}} + \hat{I})$, we may evaluate $\langle \boldsymbol{F}, \boldsymbol{E} \rangle_V = \lim_{\delta \to 0^+} \langle \boldsymbol{F}, \boldsymbol{E}_{(\delta)} \rangle_V$ with the help of the following generalized Green identity (see Ref. [67], p. 206):

$$\langle \boldsymbol{F}, \boldsymbol{E}_{(\delta)} \rangle_V = \iiint_V \boldsymbol{F}^*(\boldsymbol{r}) \cdot \nabla f_{(\delta)}(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r} + \iiint_V f_{(\delta)}(\boldsymbol{r}) \nabla \cdot \boldsymbol{F}^*(\boldsymbol{r}) \, \mathrm{d}^3\boldsymbol{r} = {}_{H^{-1/2}(\partial V; \mathbb{C})} \langle \boldsymbol{n} \cdot \boldsymbol{F} | f_{(\delta)} \rangle_{H^{1/2}(\partial V; \mathbb{C})}.$$

Here, the last item represents the canonical pairing between $\boldsymbol{n} \cdot \boldsymbol{F} \in H^{-1/2}(\partial V; \mathbb{C})$ and $f_{(\delta)} \in H^{1/2}(\partial V; \mathbb{C})$, and it necessarily reduces to a usual Lebesgue integral ${}_{H^{-1/2}(\partial V; \mathbb{C})} \langle \boldsymbol{n} \cdot \boldsymbol{F} | f_{(\delta)} \rangle_{H^{1/2}(\partial V; \mathbb{C})} = \oiint_{\partial V} f_{(\delta)}(\boldsymbol{r}) \boldsymbol{n} \cdot \boldsymbol{F}^*(\boldsymbol{r}) \, \mathrm{d}S$ if $\boldsymbol{F}$ is smooth up to the boundary. For any bounded open set $M \Subset \mathbb{R}^3$ with smooth boundary surface $\Sigma = \partial M \subset \mathbb{R}^3$, the Hilbert space $H^s(\Sigma; \mathbb{C}), s \geq 0$ is defined as (see Ref. [68], p. 37 and Ref. [69], p. 147)

$$H^s(\Sigma; \mathbb{C}) := \left\{ f : \Sigma \longrightarrow \mathbb{C} \,\middle|\, \|f\|_{H^s(\Sigma; \mathbb{C})} = \sqrt{\|f\|^2_{L^2(\Sigma; \mathbb{C})} + k^{-2s} \|(-\Delta_\Sigma)^{s/2} f\|^2_{L^2(\Sigma; \mathbb{C})}} < +\infty \right\}$$

---

[26] See S2.3.1 for the definition of a smooth surface.

where Δ_Σ is the Laplace-Beltrami operator on the surface Σ (see S2.3.1 for details),[27] and $H^{-s}(\Sigma;\mathbb{C}), s\geq 0$ is the dual space of $H^{s}(\Sigma;\mathbb{C})$, i.e. the totality of continuous linear functionals acting on $H^{s}(\Sigma;\mathbb{C})$ via the canonical pairing. For fixed $\boldsymbol{n}\cdot\boldsymbol{F}\in H^{-1/2}(\partial V;\mathbb{C})$, the canonical pairing ${}_{H^{-1/2}(\partial V;\mathbb{C})}\langle \boldsymbol{n}\cdot\boldsymbol{F}|f_{(\delta)}\rangle_{H^{1/2}(\partial V;\mathbb{C})}$ is continuous with respect to variations of $f_{(\delta)}$ in the Hilbert space $H^{1/2}(\partial V;\mathbb{C})$ equipped with the norm $\|\cdot\|_{H^{1/2}(\partial V;\mathbb{C})}$. In particular, to show that $\langle\boldsymbol{F},\boldsymbol{E}\rangle_V=\lim_{\delta\to0^+}\langle\boldsymbol{F},\boldsymbol{E}_{(\delta)}\rangle_V=0$, it would suffice to prove that $\lim_{\delta\to0^+}\|f_{(\delta)}\|_{H^{1/2}(\partial V;\mathbb{C})}=0$.

For any $\delta\leq\delta_0/6$, we have $\|f_{(\delta)}\|_{H^{1/2}(\partial V;\mathbb{C})}=\|f_{(\delta)}\|_{H^{1/2}(\partial(V^+_{(\delta_0/3)}\setminus(V\cup\partial V));\mathbb{C})}$ because the boundary $\partial(V^+_{(\delta_0/3)}\setminus(V\cup\partial V))=\partial V^+_{(\delta_0/3)}\cup\partial V$ has two disjoint components, and $f_{(\delta)}\in C_0^\infty(V^+_{(2\delta)};\mathbb{C})\subset C_0^\infty(V^+_{(\delta_0/3)};\mathbb{C})$ has to vanish identically on the component $\partial V^+_{(\delta_0/3)}$. The values of $\|f_{(\delta)}\|_{L^2(\Sigma;\mathbb{C})}$ and $\|(-\Delta_\Sigma)^{1/4}f_{(\delta)}\|_{L^2(\Sigma;\mathbb{C})}$ are not affected by the choice of either $\Sigma=\partial V$ or $\Sigma=\partial(V^+_{(\delta_0/3)}\setminus(V\cup\partial V))$. (See S2.3.1 for the localization of $(-\Delta_\Sigma)^s, 0<s<1$ on a connected component of Σ.) Meanwhile, quoting the trace theorem (see Ref. [27], p. 287 or Ref. [68], p. 39) for a bounded open set $M_0=V^+_{(\delta_0/3)}\setminus(V\cup\partial V)$ with smooth boundary ∂M_0, we have

$$\|f_{(\delta)}\|_{H^{1/2}(\partial(V^+_{(\delta_0/3)}\setminus(V\cup\partial V));\mathbb{C})}=\|f_{(\delta)}\|_{H^{1/2}(\partial M_0;\mathbb{C})}\leq C_{\delta_0;V}\sqrt{k^2\|f_{(\delta)}\|^2_{L^2(M_0;\mathbb{C})}+\|\nabla f_{(\delta)}\|^2_{L^2(M_0;\mathbb{C}^3)}},$$

where the finite constant $C_{\delta_0;V}<+\infty$ only depends on the fixed shape $M_0=V^+_{(\delta_0/3)}\setminus(V\cup\partial V)$. On one hand, the approximation capability of the convolution kernel $\varkappa_\delta$ entails the following estimate

$$\|\nabla f_{(\delta)}\|^2_{L^2(M_0;\mathbb{C}^3)}\leq\|\nabla f_{(\delta)}-\boldsymbol{E}\|^2_{L^2(V^+_{(\delta_0/3)};\mathbb{C}^3)}=\|\boldsymbol{E}_{(\delta)}-\boldsymbol{E}\|^2_{L^2(V^+_{(\delta_0/3)};\mathbb{C}^3)}\to0,\quad\text{as }\delta\to0^+,$$

where we have zero-padded $\boldsymbol{E}(\boldsymbol{r})$ for $\boldsymbol{r}\notin V$. On the other hand, we have the relation

$$\|f_{(\delta)}\|_{L^6(V^+_{(2\delta)}\setminus(V\cup\partial V);\mathbb{C})}=\|f_{(\delta)}\|_{L^6(M_0;\mathbb{C})}\leq\|f_{(\delta)}\|_{L^6(V^+_{(\delta_0/3)};\mathbb{C})}\leq\frac{1}{\sqrt{3}}\left(\frac{2}{\pi}\right)^{2/3}\|\nabla f_{(\delta)}\|_{L^2(V^+_{(\delta_0/3)};\mathbb{C}^3)},\tag{2.5}$$

which can be derived from the Gagliardo-Nirenberg-Sobolev inequality $\|f\|_{L^{dp/(d-p)}(\mathbb{R}^d;\mathbb{C})}\leq C_{d,p}\|\nabla f\|_{L^p(\mathbb{R}^d;\mathbb{C}^d)}$ (see Ref. [26], p. 158, Ref. [28], p. 263, or Ref. [66], p. 103) for smooth and compactly supported scalar fields $f\in C_0^\infty(\mathbb{R}^d;\mathbb{C})$. Here, for $d=3,p=2$, we have $dp/(d-p)=6$ and the constant $C_{3,2}=(2/\pi)^{2/3}/\sqrt{3}$ can be computed from the Bliss-Rodemich-Talenti formula [70, 71, 72]. After we combine Eq. 2.5 with the nonexpansivity property of the convolution $\|\nabla f_{(\delta)}\|_{L^2(V^+_{(\delta_0/3)};\mathbb{C}^3)}=\|\boldsymbol{E}_{(\delta)}\|_{L^2(V^+_{(\delta_0/3)};\mathbb{C}^3)}\leq\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}$ (see Theorem 2.16 in Ref. [29]) and the Hölder inequality that relates L^p-norms with different powers $\|f_{(\delta)}\|_{L^2(V^+_{(2\delta)}\setminus(V\cup\partial V);\mathbb{C})}\leq\|f_{(\delta)}\|_{L^6(V^+_{(2\delta)}\setminus(V\cup\partial V);\mathbb{C})}(\iiint_{V^+_{(2\delta)}\setminus(V\cup\partial V)}\mathrm{d}^3\tilde{\boldsymbol{r}})^{1/3}$ (Theorem 2.3 in Ref. [29]), we obtain the estimate

$$\begin{aligned}\|f_{(\delta)}\|_{L^2(M_0;\mathbb{C})}&=\|f_{(\delta)}\|_{L^2(V^+_{(2\delta)}\setminus(V\cup\partial V);\mathbb{C})}\\&\leq\frac{1}{\sqrt{3}}\left(\frac{2}{\pi}\right)^{2/3}\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\left(\iiint_{V^+_{(2\delta)}\setminus(V\cup\partial V)}\mathrm{d}^3\tilde{\boldsymbol{r}}\right)^{1/3}\\&=\frac{1}{\sqrt{3}}\left(\frac{2}{\pi}\right)^{2/3}\|\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\left[\int_0^{2\delta}\left(\oint\!\!\!\oint_{\partial V}[1-2\varepsilon H(\boldsymbol{r})+\varepsilon^2K(\boldsymbol{r})]\,\mathrm{d}S\right)\mathrm{d}\varepsilon\right]^{1/3}\to0,\quad\text{as }\delta\to0^+\end{aligned}\tag{2.6}$$

Here, with the substitution $\tilde{\boldsymbol{r}}=\boldsymbol{r}+\varepsilon\boldsymbol{n}(\boldsymbol{r})$, we have represented the volume element as $\mathrm{d}^3\tilde{\boldsymbol{r}}=[1-2\varepsilon H(\boldsymbol{r})+\varepsilon^2K(\boldsymbol{r})]\,\mathrm{d}S\,\mathrm{d}\varepsilon$, where $H(\boldsymbol{r})$ and $K(\boldsymbol{r})$ stand for the mean curvature and Gaussian curvature of the boundary surface at point $\boldsymbol{r}\in\partial V$, respectively. (See S2.3.1 for detailed explanations.) In passage to the $\delta\to0^+$ limit in Eq. 2.6, we have also exploited the fact that the surface area of ∂V is finite $\oint\!\!\oint_{\partial V}\mathrm{d}S<+\infty$ (see Corollary 2.3 in S2.3.1 for a proof).

Clearly, the above quantitative arguments lead us to the conclusion that

$$\langle\boldsymbol{F},\boldsymbol{E}\rangle_V=\lim_{\delta\to0^+}\langle\boldsymbol{F},\boldsymbol{E}_{(\delta)}\rangle_V=0,\quad\forall\boldsymbol{E}\in\ker(\hat{\mathcal{G}}+\hat{I}),\forall\boldsymbol{F}\in T(V;\mathbb{C}^3).\tag{2.7}$$

---

[27] The Hilbert spaces $H^s, s\geq0$ may be regarded as generalizations of the Sobolev spaces $W^{m,2}, m=0,1,2,\dots$ to accommodate "fractional-order derivatives".

In particular, for $\boldsymbol{E}=\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}\in\ker(\hat{\mathcal{G}}+\hat{I})$, we have $\langle\boldsymbol{F},\boldsymbol{E}\rangle_V=\langle\boldsymbol{F},\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}\rangle_V=\langle\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F},\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}\rangle_V=0$, which indicates that $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}=\mathbf{0}$ for all $\boldsymbol{F}\in T(V;\mathbb{C}^3)$. Meanwhile, by continuity, we may extend Eq. 2.7 to all $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)=\mathrm{Cl}\,(T(V;\mathbb{C}^3))$, which also implies the result $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}=\mathbf{0}$ for all $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$. $\blacksquare$

Remark From this proof, we see that if a bounded, open and connected set V has a smooth and connected boundary ∂V, the orthogonality relation $\langle\boldsymbol{E},\boldsymbol{F}\rangle_V=0$ holds for any $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$ satisfying $\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\mathbf{0},\forall\boldsymbol{q}\in\mathbb{R}^3$ and any $\boldsymbol{F}\in T(V;\mathbb{C}^3)=L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$.

It should be noted that this procedure may fail if we dispense with the topological constraint on a connected boundary. Consider, for example, a spherical shell $V=O(\mathbf{0},R_2)\setminus\mathrm{Cl}\,(O(\mathbf{0},R_1))\Subset\mathbb{R}^3$ with an inner radius $R_1>0$ and an outer radius $R_2>R_1$, with smooth but disconnected boundary surface ∂V. The vector field $\boldsymbol{F}(\boldsymbol{r})=\boldsymbol{r}/|\boldsymbol{r}|^3,\boldsymbol{r}\in V$ is both divergence-free $\nabla\cdot\boldsymbol{F}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in V$ and irrotational $\nabla\times\boldsymbol{F}(\boldsymbol{r})=\mathbf{0},\forall\boldsymbol{r}\in V$. It has vanishing tangential components on the boundary $\boldsymbol{n}\times\boldsymbol{F}(\boldsymbol{r})=\mathbf{0},\forall\boldsymbol{r}\in\partial V$, so

$$\mathbf{0}=\oiint_{\partial V}\boldsymbol{n}\times[\boldsymbol{F}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}]\,\mathrm{d}S=\iiint_V\nabla\times[\boldsymbol{F}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}]\,\mathrm{d}^3\boldsymbol{r}=\iiint_V e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\nabla\times\boldsymbol{F}(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}+i\boldsymbol{q}\times\tilde{\boldsymbol{F}}(\boldsymbol{q})=i\boldsymbol{q}\times\tilde{\boldsymbol{F}}(\boldsymbol{q})$$

also holds for any $\boldsymbol{q}\in\mathbb{R}^3$. Now, setting $\boldsymbol{E}=\boldsymbol{F}\in L^2(V;\mathbb{C}^3)$ which satisfies $\boldsymbol{q}\times\tilde{\boldsymbol{E}}(\boldsymbol{q})=\mathbf{0},\forall\boldsymbol{q}\in\mathbb{R}^3$ and $\boldsymbol{F}\in T(V;\mathbb{C}^3)$, we have $\langle\boldsymbol{E},\boldsymbol{F}\rangle_V=\langle\boldsymbol{F},\boldsymbol{F}\rangle_V>0$. According to Proposition 2.7 of S2.3.2, a *connected* and smooth boundary surface ∂V suffices to forestall the anomalies in such a counterexample, i.e. a divergence-free and irrotational vector field $\boldsymbol{F}$ that has vanishing tangential components on the boundary must vanish identically in that scenario.

During the proof of Proposition 2.4, the smoothness assumption for the boundary surface ∂V is convenient, but not totally indispensable. In S2.3.3, we show that for a certain class of dielectric shapes without a smooth boundary, the conclusion $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F}=\mathbf{0},\forall\boldsymbol{F}\in T(V;\mathbb{C}^3)$ remains valid. In the light of this, we may say that the driving force for the orthogonality condition (Eq. 2.7) is more topological than geometrical. $\blacksquare$

Proposition 2.4 shows that the action of the evolution semigroup on physically admissible vector fields leads to an asymptotically vanishing (in the sense of τ-average) trajectory $\{\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}|\tau\geq0\}$.

Corollary 2.1 (Mean Ergodicity for Physically Admissible Fields) *For a smooth dielectric with connected exterior volume $\mathbb{R}^3\setminus(V\cup\partial V)$, we have $\sigma_p^T(\hat{\mathcal{G}})\cap\mathbb{R}=\emptyset$. Accordingly, we have the mean ergodicity result for physically admissible fields*

$$\lim_{T\to+\infty}\frac{1}{T}\int_{\tau_0}^{\tau_0+T}\exp(-i\tau(\hat{\mathcal{G}}-\lambda\hat{I}))\boldsymbol{F}\,\mathrm{d}\tau=\mathbf{0},\quad\tau_0\geq0,\quad\forall\boldsymbol{F}\in\Phi(V;\mathbb{C}^3),\quad\forall\lambda\in\mathbb{R}.$$

Proof As shown in Proposition 2.4, any square-integrable solution $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$ to the equation $\|(\hat{\mathcal{G}}+\hat{I})\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$ must be orthogonal to $\Phi(V;\mathbb{C}^3)$: $\langle\boldsymbol{E},\boldsymbol{F}\rangle_V=0,\forall\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$. Therefore, there is no non-zero member $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$ that may solve the eigenequation $(\hat{\mathcal{G}}+\hat{I})\boldsymbol{F}=\mathbf{0}$, meaning that $-1\notin\sigma_p^\Phi(\hat{\mathcal{G}})$. For $\lambda\in\mathbb{R}\setminus\{-1\}$, we have $\lambda\notin\sigma_p^\Phi(\hat{\mathcal{G}})$ according to the uniqueness theorem (situation (B) in Theorem 1.2) and Proposition 1.6. This proves that $\sigma_p^\Phi(\hat{\mathcal{G}})\cap\mathbb{R}=\emptyset$.

According to Lemma 2.1, we have $\exp(-i\tau_0(\hat{\mathcal{G}}-\lambda\hat{I}))\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$ for every $\tau_0\geq0$ and $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$, so the mean ergodicity statement in Proposition 2.2 can be adapted to show that the vanishing limit

$$\lim_{T\to+\infty}\frac{1}{T}\int_0^T\exp(-i\tau(\hat{\mathcal{G}}-\lambda\hat{I}))\exp(-i\tau_0(\hat{\mathcal{G}}-\lambda\hat{I}))\boldsymbol{F}\,\mathrm{d}\tau=\mathbf{0}$$

must hold. $\blacksquare$

Remark It is worth noting that we have $-1\in\sigma_p(\hat{\mathcal{G}}),\dim\ker(\hat{I}+\hat{\mathcal{G}})=+\infty$ for $\hat{\mathcal{G}}:L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ according to Proposition 1.5, while we have $-1\notin\sigma_p^\Phi(\hat{\mathcal{G}}),\dim[\ker(\hat{I}+\hat{\mathcal{G}})\cap\Phi(V;\mathbb{C}^3)]=0$ for $\hat{\mathcal{G}}:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ according to Propositions 2.8 and 2.4. This shows that the spectral rôle of the point $1/\chi=-1$ has shifted dramatically from an eigenvalue of infinite multiplicity with respect to the Hilbert space $L^2(V;\mathbb{C}^3)$ of *energetically* admissible vector fields to a non-eigenvalue with respect to the Hilbert subspace $\Phi(V;\mathbb{C}^3)$ of *physically* admissible vector fields. Although the Born equation may appear "seriously singular" for vanishing relative permittivity $\epsilon_r=1+\chi=0$ in the Hilbert space $L^2(V;\mathbb{C}^3)$ (which has been classified as "essential resonance" in Ref. [73]), we see from Propositions 2.8 and 2.4 that such a "serious singularity" is effectively mitigated by the physical admissibility condition in the Hilbert subspace $\Phi(V;\mathbb{C}^3)$. In a moment, we will see that Proposition 2.4 can actually be sharpened into $-1\in\rho^\Phi(\hat{\mathcal{G}})$, so the case $\epsilon_r=1+\chi=0$ is not physically singular, after all. $\blacksquare$

S2.2. Smooth Solution, Optical Resonance and Strong Stability — Extended Existence Theorem

We note that restricting the Born equation to physically admissible fields $\boldsymbol{F} \in \Phi(V;\mathbb{C}^3)$ brings some additional benefits that are not warranted by the energy admissibility (square-integrability) alone. Three notable bonuses are the smoothness (infinite differentiability) of the solution, universality of spectral structure $\sigma^{\Phi}(\hat{\mathcal{G}}) \setminus \sigma^{\Phi}_{p}(\hat{\mathcal{G}}) \subset \{0,-1/2\}$ and the strong stability $\lim_{\tau\to+\infty} \|\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)} = 0, \forall \boldsymbol{F} \in \Phi(V;\mathbb{C}^3)$ of the associated evolution semigroup.

Proposition 2.5 (Smooth Solution to Light Scattering) *If the incident field is a smooth and divergence-free vector field defined in a bounded and open volume* $V \Subset \mathbb{R}^3$*, i.e.* $\boldsymbol{E}_{\rm inc} \in T(V;\mathbb{C}^3) = L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) \cap \ker(\nabla\cdot)$*, then the solution to the Born equation* $(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc}$ *is also smooth and divergence-free, i.e. we have* $(\hat{I}-\chi\hat{\mathcal{G}})^{-1}T(V;\mathbb{C}^3) \subset T(V;\mathbb{C}^3)$ *whenever* $1/\chi \in \rho^{\Phi}(\hat{\mathcal{G}})$*.*

Proof It would suffice to show that when $\boldsymbol{E}_{\rm inc} \in T(V;\mathbb{C}^3)$ and $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in \Phi(V;\mathbb{C}^3)$ both hold, we have $(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in T(V;\mathbb{C}^3)$ as a result. To confirm this, we use the decomposition $\hat{\mathcal{G}} = (\hat{\mathcal{D}}-\hat{I}) + \hat{\gamma}$ and show that $(\hat{\mathcal{D}}-\hat{I})\boldsymbol{E}$ is always a smooth vector field satisfying the Laplace equation, for all $\boldsymbol{E} \in \Phi(V;\mathbb{C}^3)$.

First, we note that for a twice continuously differentiable vector field $\hat{\mathcal{D}}\boldsymbol{E} \in C^2(V;\mathbb{C}^3)$, the vector identity $\nabla\times\nabla\times(\hat{\mathcal{D}}\boldsymbol{E}) = -\nabla^2(\hat{\mathcal{D}}\boldsymbol{E}) + \nabla[\nabla\cdot(\hat{\mathcal{D}}\boldsymbol{E})]$ holds, where all the derivatives are taken in the classical sense. Therefore, for any $\boldsymbol{E} \in T(V;\mathbb{C}^3) = L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) \cap \ker(\nabla\cdot)$, we have

$$((\hat{\mathcal{D}}-\hat{I})\boldsymbol{E})(\boldsymbol{r}) = \nabla\times\nabla\times\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' - \boldsymbol{E}(\boldsymbol{r}') = \nabla\left[\nabla\iiint_V \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right], \quad \boldsymbol{r} \in V.$$

We claim that $(\hat{\mathcal{D}}-\hat{I})\boldsymbol{E} \in \mathfrak{b}^2(V;\mathbb{C}^3) := L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) \cap \ker(\nabla^2) = \{\boldsymbol{F} \in L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) | \nabla^2\boldsymbol{F}(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in V\}$, where $\mathfrak{b}^p(V;\mathbb{C}^3)$ denotes the Bergman space of power $p \geq 1$ (see Ref. [74], p. 171 or Ref. [75]). This can be shown by picking any enclosed open ball $O(\boldsymbol{r}_*,\varepsilon) \subset V$, and applying the Gauss theorem in vector calculus, as follows:

$$((\hat{\mathcal{D}}-\hat{I})\boldsymbol{E})(\boldsymbol{r}) = \nabla\left[\nabla\cdot\iiint_{V\setminus O(\boldsymbol{r}_*,\varepsilon)} \frac{\boldsymbol{E}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right] + \oiint_{\partial O(\boldsymbol{r}_*,\varepsilon)} [\boldsymbol{n}'\cdot\boldsymbol{E}(\boldsymbol{r}')]\nabla'\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}S'.$$

For $\boldsymbol{r} \in O(\boldsymbol{r}_*,\varepsilon/2)$, one can always differentiate under the integral sign to arbitrarily high order, so as to conclude that $(\hat{\mathcal{D}}-\hat{I})\boldsymbol{E} \in C^\infty(O(\boldsymbol{r}_*,\varepsilon/2);\mathbb{C}^3)$, and the Laplace equation $\nabla^2((\hat{\mathcal{D}}-\hat{I})\boldsymbol{E})(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in O(\boldsymbol{r}_*,\varepsilon/2)$ holds. As $O(\boldsymbol{r}_*,\varepsilon)$ is chosen arbitrarily, this shows that $\nabla^2((\hat{\mathcal{D}}-\hat{I})\boldsymbol{E})(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in V$.

Then, we point out that for all $\boldsymbol{E} \in \Phi(V;\mathbb{C}^3) = \mathrm{Cl}\,(T(V;\mathbb{C}^3))$, we have $(\hat{\mathcal{D}}-\hat{I})\boldsymbol{E} \in (\hat{\mathcal{D}}-\hat{I})\mathrm{Cl}\,(T(V;\mathbb{C}^3)) \subset \mathrm{Cl}\,((\hat{\mathcal{D}}-\hat{I})T(V;\mathbb{C}^3)) \subset \mathrm{Cl}\,(\mathfrak{b}^2(V;\mathbb{C}^3)) = \mathfrak{b}^2(V;\mathbb{C}^3) = L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) \cap \ker(\nabla^2)$. This is because $(\hat{\mathcal{D}}-\hat{I}) : L^2(V;\mathbb{C}^3) \longrightarrow L^2(V;\mathbb{C}^3)$ is continuous, and the Bergman space $\mathfrak{b}^2(V;\mathbb{C}^3) = \mathrm{Cl}\,(\mathfrak{b}^2(V;\mathbb{C}^3))$ is a closed subspace of the Hilbert space $L^2(V;\mathbb{C}^3)$ (see Proposition 8.3 in Ref. [74]). Thus, for $1/\chi \in \rho(\hat{\mathcal{G}})$, we have the Born equation

$$\boldsymbol{E}(\boldsymbol{r}) = \boldsymbol{E}_{\rm inc}(\boldsymbol{r}) + \chi(\hat{\gamma}\boldsymbol{E})(\boldsymbol{r}) + \chi((\hat{\mathcal{D}}-\hat{I})\boldsymbol{E})(\boldsymbol{r}), \quad \forall \boldsymbol{r} \in V,$$

where $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in \Phi(V;\mathbb{C}^3) = \mathrm{Cl}\,(T(V;\mathbb{C}^3))$ and $(\hat{\mathcal{D}}-\hat{I})\boldsymbol{E} \in \mathfrak{b}^2(V;\mathbb{C}^3)$. Using the Cauchy-Schwarz inequality, one can verify that $\hat{\gamma}\boldsymbol{E} \in C^0(\mathbb{R}^3;\mathbb{C}^3)$ for all square-integrable vector fields $\boldsymbol{E} \in L^2(V;\mathbb{C}^3)$. Therefore, all the three terms on the right-hand side of the Born equation belongs to $C^0(V;\mathbb{C}^3)$, hence $\boldsymbol{E} \in C^0(V;\mathbb{C}^3)$. Using differentiation under the integral sign, one can check that $\boldsymbol{E} \in C^0(V;\mathbb{C}^3) \cap L^2(V;\mathbb{C}^3)$ entails $\hat{\gamma}\boldsymbol{E} \in C^1(V;\mathbb{C}^3)$, so $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in C^1(V;\mathbb{C}^3)$. By induction, we may deduce $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in C^{m+1}(V;\mathbb{C}^3)$ from the assumption that $\boldsymbol{E} \in C^m(V;\mathbb{C}^3)$ for all $m = 0,1,2,\dots$, using differentiation under the integral sign and integration by parts. Hence, we may conclude that the solution to the Born equation is a smooth vector field $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3)$.

Taking divergence on both sides of the Born equation, one thus confirms that $\boldsymbol{E} \in \ker(\nabla\cdot)$ as well. This shows that the solution $\boldsymbol{E} = (\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\rm inc} \in L^2(V;\mathbb{C}^3) \cap C^\infty(V;\mathbb{C}^3) \cap \ker(\nabla\cdot)$ must also be a transverse field. ▮

Now we will state and prove the "optical resonance theorem" — a technical result that both outlines the structure of the spectrum $\sigma^{\Phi}(\hat{\mathcal{G}})$ and implies the strong stability of the operator semigroup $\{\exp(-i\tau\hat{\mathcal{G}}) | \tau \geq 0\}$ acting on the function space of physically admissible vector fields $\Phi(V;\mathbb{C}^3)$.

Theorem 2.1 (Optical Resonance in Light Scattering) *For a smooth dielectric volume* $V \Subset \mathbb{R}^3$*, the operator* $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$ *is compact, and* $\sigma^\Phi(\hat{\mathcal{G}}) \subset \sigma^\Phi_p(\hat{\mathcal{G}}) \cup \{0,-1/2\}$*. The solution to the Born equation* $(\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\mathrm{inc}} \in \Phi(V;\mathbb{C}^3)$ *can be represented as*

$$\begin{aligned} \boldsymbol{E}(\boldsymbol{r}) &= \boldsymbol{F}(\zeta;\boldsymbol{r}) + 2\frac{n^2-1}{n^2+1}\nabla\times\nabla\times\iiint_V \frac{\boldsymbol{F}(\zeta;\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad \forall \boldsymbol{r}\in V, \\ &\textit{where } \zeta = \frac{(n^2-1)^2}{n^2+1}, \quad n^2 = 1+\chi, \quad 1/\chi \in \mathbb{C}\setminus\sigma^\Phi(\hat{\mathcal{G}}) \end{aligned} \tag{2.8}$$

whenever $\boldsymbol{F}(\zeta;\boldsymbol{r}) = ((\hat{I}-\zeta\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}))^{-1}\boldsymbol{E}_{\mathrm{inc}})(\boldsymbol{r})$ *is well-defined. Each Cartesian component of the vector* $\boldsymbol{F}(\zeta;\boldsymbol{r})$ *is a meromorphic function of* $\zeta \in \mathbb{C}$*, with a discrete set of isolated singularities* $\{\zeta_s|s=1,2,\dots\}$ *corresponding to the "optical resonance modes".*

Proof The proof will be divided into two parts. In Part I, we show how the compactness of the operator $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$ leads to the rest of the conclusions; in Part II, we elaborate on a long and technical proof that the operator $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$ is indeed compact.

$\boxed{\text{PART I}}$ Recalling the Riesz-Schauder theory for the spectrum of compact operators (see Ref. [43] or Ref. [44], pp. 283-284), we have a countable set $\sigma^\Phi(\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}})) = \{0\} \cup \sigma^\Phi_p(\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}})) = \{0\} \cup \{\mu_s|s=1,2,\dots\}$ with $\{0\}$ being the only possible accumulation point. Owing to the spectral mapping theorem (see Ref. [44], p. 227), we have

$$\sigma^\Phi(\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}})) = \{\lambda(1+2\lambda)|\lambda \in \sigma^\Phi(\hat{\mathcal{G}})\}. \tag{2.9}$$

We note that every $\lambda(1+2\lambda) \in \sigma^\Phi(\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}))\setminus\{0\}$ leads to a pole of the resolvent $[z\hat{I}-\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}})]^{-1} : \Phi(O(\mathbf{0},R);\mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0},R);\mathbb{C}^3)$ at $z = \lambda(1+2\lambda)$ (see Ref. [44], p. 286). Consequently, the algebraic factorization

$$(\lambda\hat{I}-\hat{\mathcal{G}})^{-1} = [\lambda(1+2\lambda)\hat{I}-\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}})]^{-1}[(1+2\lambda)\hat{I}+2\hat{\mathcal{G}}] \tag{2.10}$$

reveals that every $\lambda \in \sigma^\Phi(\hat{\mathcal{G}}) \setminus \{0,-1/2\}$ must also lead to a pole of the resolvent $(z\hat{I}-\hat{\mathcal{G}})^{-1} : \Phi(O(\mathbf{0},R);\mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0},R);\mathbb{C}^3)$ at $z=\lambda$, thereby giving rise to an eigenvalue of $\hat{\mathcal{G}}$ with finite multiplicity. In other words, we have the following relation

$$\sigma^\Phi(\hat{\mathcal{G}}) \setminus \sigma^\Phi_p(\hat{\mathcal{G}}) \subset \{0,-1/2\}. \tag{2.11}$$

Combining the clues in Eqs. 2.9-2.11, we may conclude that $\sigma^\Phi(\hat{\mathcal{G}}) \subset \sigma^\Phi_p(\hat{\mathcal{G}}) \cup \{0,-1/2\}$ is a countable set, with $\{0,-1/2\}$ being the only possible accumulation points in $\sigma^\Phi(\hat{\mathcal{G}})$, as well as the only possible points in the continuous spectrum $\sigma^\Phi_c(\hat{\mathcal{G}}) \subset \{0,-1/2\}$.

Dividing the algebraic factorization (Eq. 2.10) by χ, we obtain Eq. 2.8 as claimed. The absence of finite accumulation points in the set $\{\zeta \in \mathbb{C}|1/\zeta \in \sigma^\Phi_p(\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}))\}$ indicates that the totality of singularities for the vector field $\boldsymbol{F}(\zeta;\boldsymbol{r}) = ((\hat{I}-\zeta\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}))^{-1}\boldsymbol{E}_{\mathrm{inc}})(\boldsymbol{r})$ is a discrete and isolated set in the complex ζ-plane. Each singularity of $\boldsymbol{F}(\zeta;\boldsymbol{r})$ in the complex ζ-plane is a pole of finite order (as opposed to essential singularity), because $1/\zeta$ corresponds to an eigenvalue of the compact operator $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$. Therefore, $\boldsymbol{F}(\zeta;\boldsymbol{r})$ is a meromorphic function of $\zeta \in \mathbb{C}$, with isolated singularities.

$\boxed{\text{PART II}}$ We note that

$$\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) - (\hat{\mathcal{G}}-\hat{\gamma})(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma})) = \hat{\gamma}+2\hat{\mathcal{G}}\hat{\gamma}+2\hat{\gamma}\hat{\mathcal{G}}-2\hat{\gamma}^2 \tag{2.12}$$

is a compact operator, as each addend on the right-hand side of Eq. 2.12 is a composition of a compact operator with a bounded operator, thus also a compact operator in its own right. Therefore, we may establish the compactness of the operator $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$ by proving that the operator $(\hat{\mathcal{G}}-\hat{\gamma})(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma})) : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)$ is compact. We will fulfill this task by the justification of the following two claims:

(i) Define a family of cut-off operators parameterized by $\varepsilon \in (0,\varepsilon_V]$ as follows:

$$(\hat{\mathcal{A}}_{(\varepsilon)}\boldsymbol{F})(\boldsymbol{r}) = \begin{cases} ((\hat{\mathcal{G}}-\hat{\gamma}))\boldsymbol{F})(\boldsymbol{r}), & \boldsymbol{r} \in V^-_{(\varepsilon)} := V \setminus \{\tilde{\boldsymbol{r}} = \boldsymbol{r}^\sharp - \varepsilon'\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp \in \partial V, 0 < \varepsilon' \le \varepsilon\}; \\ \mathbf{0}, & \boldsymbol{r} \in V \setminus V^-_{(\varepsilon)}, \end{cases}$$

where the positive parameter $\varepsilon_V > 0$ is determined by the shape of V according to the tubular neighborhood theorem (Proposition 2.6 in S2.3.1), such that $\partial V^-_{(\varepsilon)} = \{\tilde{\boldsymbol{r}} = \boldsymbol{r}^\sharp - \varepsilon\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp \in \partial V\}$ is a smooth surface for all

$\varepsilon\in(0,2\varepsilon_V]$, with bounded area $\oint\!\!\!\oint_{\partial V^-_{(\varepsilon)}}\mathrm{d}\,\tilde S\le 2\sum_{j=1}^{N_V}\oint\!\!\!\oint_{\partial V_j}\mathrm{d}\,S<+\infty$.[28] Then the operator $\hat{\mathcal{A}}_{(\varepsilon)}:\Phi(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ is compact for $\varepsilon\in(0,\varepsilon_V]$.

(ii) Let $\hat P: L^2(V;\mathbb{C}^3)\longrightarrow \mathrm{Cl}\,(\operatorname{ran}^{\Phi}(\hat{\mathcal{G}}-\hat\gamma))$ be the orthogonal projection operator, and $\hat{\mathcal{R}}_{(\varepsilon)}: L^2(V;\mathbb{C}^3)\longrightarrow L^2(V_{(\varepsilon)};\mathbb{C}^3)\subset L^2(V;\mathbb{C}^3)$ be the restriction operator defined by

$$(\hat{\mathcal{R}}_{(\varepsilon)}\boldsymbol{F})(\boldsymbol{r})=\begin{cases}\boldsymbol{F}(\boldsymbol{r}), & \boldsymbol{r}\in V^-_{(\varepsilon)};\\ \boldsymbol{0}, & \boldsymbol{r}\in V\setminus V^-_{(\varepsilon)},\end{cases}$$

then both operators $\hat P,\hat{\mathcal{R}}_{(\varepsilon)}: L^2(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ are Hermitian, and the family of compact operators $\{\hat{\mathcal{K}}_{(\varepsilon)}=\hat P\hat{\mathcal{R}}_{(\varepsilon)}(\hat I+2(\hat{\mathcal{G}}-\hat\gamma))\hat{\mathcal{A}}_{(\varepsilon)}\hat P:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)|\varepsilon\in(0,\varepsilon_V]\}$ converges uniformly to the operator $(\hat{\mathcal{G}}-\hat\gamma)(\hat I+2(\hat{\mathcal{G}}-\hat\gamma)):\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$, in the following sense

$$\lim_{\varepsilon\to0^+}\left[\sup_{\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)\setminus\{\boldsymbol{0}\}}\frac{\|(\hat{\mathcal{G}}-\hat\gamma)(\hat I+2(\hat{\mathcal{G}}-\hat\gamma))\boldsymbol{F}-\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}}{\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}}\right]=0. \tag{2.13}$$

Hence, the operator $(\hat{\mathcal{G}}-\hat\gamma)(\hat I+2(\hat{\mathcal{G}}-\hat\gamma)):\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)$ is compact as well.

Proof of claim (i): For any $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$, we may express $(\hat{\mathcal{G}}-\hat\gamma)\boldsymbol{F}$ as the gradient of a harmonic function $(\hat{\mathcal{G}}-\hat\gamma)\boldsymbol{F}=\nabla h, h\in\mathfrak{b}^6(V;\mathbb{C})\cap W^{1,2}(V;\mathbb{C})$. It satisfies the bound estimate $\|\nabla h\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)}=\|(\hat{\mathcal{G}}-\hat\gamma)\boldsymbol{F}\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)}\le\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}$, according to Lemma 1.1. Using the Gagliardo-Nirenberg-Sobolev inequality $\|h\|_{L^6(\mathbb{R}^3;\mathbb{C})}\le(2/\pi)^{2/3}3^{-1/2}\|\nabla h\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)}$ and the Hölder inequality $\|h\|_{L^2(V;\mathbb{C})}\le\|h\|_{L^6(V;\mathbb{C})}\left(\iiint_V\mathrm{d}^3\boldsymbol{r}\right)^{1/3}$, we then arrive at

$$\|h\|_{L^2(V;\mathbb{C})}\le\|h\|_{L^6(\mathbb{R}^3;\mathbb{C})}\left(\iiint_V\mathrm{d}^3\boldsymbol{r}\right)^{1/3}\le\frac{1}{\sqrt3}\left(\frac{2}{\pi}\right)^{2/3}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\left(\iiint_V\mathrm{d}^3\boldsymbol{r}\right)^{1/3}. \tag{2.14}$$

For the "pinched volume" $V^-_{(\varepsilon)}\Subset V$, we have the interior regularity estimate (see Ref. [30], p. 237 or Ref. [27], p. 379)

$$\|h\|_{W^{2,2}(V^-_{(\varepsilon)};\mathbb{C})}\le C_{\varepsilon,k;V}\left[\|h\|_{L^2(V;\mathbb{C})}+\frac{\|\nabla^2h\|_{L^2(V;\mathbb{C})}}{k^2}\right],\quad C_{\varepsilon,k;V}<+\infty. \tag{2.15}$$

Combining the Laplace equation $\nabla^2h(\boldsymbol{r})=0,\forall\boldsymbol{r}\in V$ with Eqs. 2.14 and 2.15, we obtain the estimate

$$\|\hat{\mathcal{A}}_{(\varepsilon)}\boldsymbol{F}\|_{W^{1,2}(V^-_{(\varepsilon)};\mathbb{C}^3)}=\|\nabla h\|_{W^{1,2}(V^-_{(\varepsilon)};\mathbb{C}^3)}\le k\|h\|_{W^{2,2}(V^-_{(\varepsilon)};\mathbb{C})}\le\frac{kC_{\varepsilon,k;V}}{\sqrt3}\left(\frac{2}{\pi}\right)^{2/3}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\left(\iiint_V\mathrm{d}^3\boldsymbol{r}\right)^{1/3}.$$

In other words, the mapping $\hat{\mathcal{A}}_{(\varepsilon)}:\Phi(V;\mathbb{C}^3)\longrightarrow W^{1,2}(V^-_{(\varepsilon)};\mathbb{C}^3)$ is continuous. Meanwhile, the natural inclusion map $\iota: W^{1,2}(V^-_{(\varepsilon)};\mathbb{C}^3)\longrightarrow L^2(V^-_{(\varepsilon)};\mathbb{C}^3)$ is a compact operator provided that the boundary surface $\partial V^-_{(\varepsilon)}$ is smooth (see Ref. [26], p. 171 or Ref. [27], p. 286 for the Rellich-Kondrachov theorem of compact imbedding). Accordingly, the mapping $\hat{\mathcal{A}}_{(\varepsilon)}:\Phi(V;\mathbb{C}^3)\longrightarrow L^2(V^-_{(\varepsilon)};\mathbb{C}^3)$ is the composition of a continuous operator with a compact operator, hence $\hat{\mathcal{A}}_{(\varepsilon)}:\Phi(V;\mathbb{C}^3)\longrightarrow L^2(V;\mathbb{C}^3)$ is a compact operator.

Proof of claim (ii): Our main goal is to use an integral representation of the sesquilinear form $\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V,\boldsymbol{F},\boldsymbol{F}^\sharp\in\Phi(V;\mathbb{C}^3)$ to prove that the set of operators $\{\hat{\mathcal{K}}_{(\varepsilon_V/s)}=\hat P\hat{\mathcal{R}}_{(\varepsilon_V/s)}(\hat I+2(\hat{\mathcal{G}}-\hat\gamma))\hat{\mathcal{A}}_{(\varepsilon_V/s)}\hat P:\Phi(V;\mathbb{C}^3)\longrightarrow\Phi(V;\mathbb{C}^3)|s=1,2,\dots\}$ forms a Cauchy sequence with respect to operator norm:

$$\lim_{s,s'\to\infty}\left[\sup_{\boldsymbol{F}^\natural\in\Phi(V;\mathbb{C}^3)\setminus\{\boldsymbol{0}\}}\frac{\|\hat{\mathcal{K}}_{(\varepsilon_V/s')}\boldsymbol{F}^\natural-\hat{\mathcal{K}}_{(\varepsilon_V/s)}\boldsymbol{F}^\natural\|_{L^2(V;\mathbb{C}^3)}}{\|\boldsymbol{F}^\natural\|_{L^2(V;\mathbb{C}^3)}}\right]=0. \tag{2.16}$$

---

[28] During the later stages of this proof, we keep using $\mathrm{d}\,\tilde S$ for the surface element of a parallel surface other than the boundary surface ∂V. Meanwhile, we will drop the tilde for $\tilde{\boldsymbol{r}}$ and use $\boldsymbol{r}$ to denote a point in the dielectric volume V, so long as this substitution does not lead to ambiguities.

Therefore, the sequence of operators $\{\hat{\mathcal{K}}_{(\varepsilon_V/s)} : \Phi(V;\mathbb{C}^3) \longrightarrow \Phi(V;\mathbb{C}^3)|s = 1,2,\dots\}$ is convergent with respect to operator norm. We then claim that Eq. 2.13 is an immediate consequence, because we have the following strong convergence result for every individual vector field $\boldsymbol{F}^\natural \in \Phi(V;\mathbb{C}^3)$:

$$\lim_{\varepsilon\to0^+}\|(\hat{\mathcal{G}}-\hat{\gamma})(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma}))\boldsymbol{F}^\natural-\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F}^\natural\|_{L^2(V;\mathbb{C}^3)}=\lim_{\varepsilon\to0^+}\|\hat{P}(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma}))(\hat{\mathcal{G}}-\hat{\gamma})\hat{P}\boldsymbol{F}^\natural-\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F}^\natural\|_{L^2(V;\mathbb{C}^3)}=0,$$

according to the absolute continuity of convergent integrals. The main goal in the proof of claim (ii) will be accomplished in two steps, as elaborated below.

$\boxed{1}$ In the first step, we will establish the identity

$$\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V=\oiint_{\partial V^-_{(\varepsilon)}}\{[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot[\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]-[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})][\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]\}\,\mathrm{d}\tilde{S}\tag{2.17}$$

where

$$\boldsymbol{U}(\boldsymbol{r}):=((\hat{\mathcal{G}}-\hat{\gamma})(\hat{P}\boldsymbol{F})^*)(\boldsymbol{r})=\nabla\left[\nabla\cdot\iiint_V\frac{(\hat{P}\boldsymbol{F})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right],\quad \boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r}):=(\hat{\mathcal{N}}\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp)(\boldsymbol{r})=-\iiint_{V^-_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^\sharp)(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'.$$

Before proving Eq. 2.17, we need to work out some analytic properties of the vector fields $\boldsymbol{U}$ and $\boldsymbol{Q}_{(\varepsilon)}$, and derive surface integral representations for the scalar field $\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}$ as well as the vector field $\nabla\times\boldsymbol{Q}_{(\varepsilon)}$. Following the arguments carried out in Proposition 2.5, we may check that $\boldsymbol{U}\in C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)\cap\ker(\nabla\times)\cap L^2(V;\mathbb{C}^3)\subset\mathfrak{b}^2(V;\mathbb{C}^3)$. Meanwhile, by definition, for every $\boldsymbol{E}\in\operatorname{ran}^\Phi(\hat{\mathcal{G}}-\hat{\gamma})\subset\mathfrak{b}^2(V;\mathbb{C}^3)$, we may write $\boldsymbol{E}=-\nabla u$ where the scalar potential $u\in\mathfrak{b}^6(V;\mathbb{C})\cap W^{1,2}(V;\mathbb{C})$ is a smooth function, so we have $\nabla\times\boldsymbol{E}(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in V$, whenever $\boldsymbol{E}\in\operatorname{ran}^\Phi(\hat{\mathcal{G}}-\hat{\gamma})$. We can further show that $\nabla\times(\hat{P}\boldsymbol{F}^\natural)(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in V$ for all $\boldsymbol{F}^\natural\in\Phi(V;\mathbb{C}^3)$, by approximating the harmonic vector field $\hat{P}\boldsymbol{F}^\natural\in\operatorname{Cl}(\operatorname{ran}^\Phi(\hat{\mathcal{G}}-\hat{\gamma}))\subset\mathfrak{b}^2(V;\mathbb{C}^3)$ with a sequence of harmonic irrotational vector fields $\{\boldsymbol{E}_s\in\operatorname{ran}^\Phi(\hat{\mathcal{G}}-\hat{\gamma})\subset\mathfrak{b}^2(V;\mathbb{C}^3)\cap\ker(\nabla\times)|s=1,2,\dots\}$ and exploit the Bergman space inequality (see Ref. [74], p. 172)

$$|(D^{\boldsymbol{\mu}}\hat{P}\boldsymbol{F}^\natural)(\boldsymbol{r})-(D^{\boldsymbol{\mu}}\boldsymbol{E}_s)(\boldsymbol{r})|\le\frac{C'\|\hat{P}\boldsymbol{F}^\natural-\boldsymbol{E}_s\|_{L^2(V;\mathbb{C}^3)}}{[\operatorname{dist}(\boldsymbol{r},\partial V)]^{5/2}},\quad C'<+\infty,|\boldsymbol{\mu}|=1,\operatorname{dist}(\boldsymbol{r},\partial V):=\min_{\boldsymbol{r}'\in\partial V}|\boldsymbol{r}-\boldsymbol{r}'|.$$

Here, we note that the L^2-convergence $\lim_{s\to\infty}\|\hat{P}\boldsymbol{F}^\natural-\boldsymbol{E}_s\|_{L^2(V;\mathbb{C}^3)}=0$ implies the pointwise convergence $|\nabla\times(\hat{P}\boldsymbol{F}^\natural)(\boldsymbol{r})|=\lim_{s\to\infty}|\nabla\times(\hat{P}\boldsymbol{F}^\natural)(\boldsymbol{r})-\nabla\times\boldsymbol{E}_s(\boldsymbol{r})|=0,\forall\boldsymbol{r}\in V$. In the light of this, we may verify that $\hat{P}\boldsymbol{F}^\sharp\in C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)\cap\ker(\nabla\times)$ is a smooth, divergence-free and irrotational vector field, leading to the expressions

$$\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})=\oiint_{\partial V^-_{(\varepsilon)}}\frac{\boldsymbol{n}'\cdot(\hat{P}\boldsymbol{F})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\tilde{S}'\quad\text{and}\quad\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})=\oiint_{\partial V^-_{(\varepsilon)}}\frac{\boldsymbol{n}'\times(\hat{P}\boldsymbol{F})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\tilde{S}',\quad\forall\boldsymbol{F}\in\Phi(V;\mathbb{C}^3),\tag{2.18}$$

as well as the assertion that both $\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\in\mathfrak{b}^6(V^-_{(\varepsilon)};\mathbb{C})\cap W^{1,6}(V^-_{(\varepsilon)};\mathbb{C})\cap C^0(V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)};\mathbb{C})$ and $\nabla\times\boldsymbol{Q}_{(\varepsilon)}\in\mathfrak{b}^6(V^-_{(\varepsilon)};\mathbb{C}^3)\cap W^{1,6}(V^-_{(\varepsilon)};\mathbb{C}^3)\cap C^0(V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)};\mathbb{C}^3)$ are continuous up to the boundary $\partial V^-_{(\varepsilon)}$. To explain the last assertion in more detail, we need to clarify three points: (1) We have the relation

$$\|\hat{P}\boldsymbol{F}\|_{L^6(V^-_{(\varepsilon)};\mathbb{C}^3)}\le\sqrt{\frac{3}{4\pi}}\left(\iiint_{V^-_{(\varepsilon)}}\frac{\mathrm{d}^3\boldsymbol{r}}{[\operatorname{dist}(\boldsymbol{r},\partial V)]^9}\right)^{1/6}\|\hat{P}\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)}\le A_{\varepsilon;V}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)},\quad\text{where }A_{\varepsilon;V}<+\infty,$$

according to the Bergman space inequality (see Ref. [74], p. 172)

$$|h(\boldsymbol{r})|^2\le\frac{3\|h\|^2_{L^2(V;\mathbb{C})}}{4\pi[\operatorname{dist}(\boldsymbol{r},\partial V)]^3},\quad\forall h\in\mathfrak{b}^2(V;\mathbb{C});\tag{2.19}$$

(2) We have the estimate $\|(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\boldsymbol{Q}_{(\varepsilon)}\|_{L^6(V^-_{(\varepsilon)};\mathbb{C}^3)}=\|(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\hat{\mathcal{N}}\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp\|_{L^6(V^-_{(\varepsilon)};\mathbb{C}^3)}\le C'_{3,6}\|\hat{P}\boldsymbol{F}^\sharp\|_{L^6(V^-_{(\varepsilon)};\mathbb{C}^3)},\forall\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}$ according to the Calderón-Zygmund inequality (see Theorem 9.9 in Ref. [26] or Theorem 10.2.1 in Ref. [30]); (3) When $M\Subset\mathbb{R}^3$ is a bounded open set with smooth boundary, we have

the Sobolev imbedding relation $W^{1,6}(M;\mathbb{C}^{d'})\subset C^0(M\cup\partial M;\mathbb{C}^{d'})$ for $d'=1,3$ (see Ref. [66], p. 85), along with the Sobolev inequalities

$$\|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\|_{C^0(V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)};\mathbb{C})} := \max_{\boldsymbol{r}\in V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)}}|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})|\le C''\|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\|_{W^{1,6}(V^-_{(\varepsilon)};\mathbb{C})}\le C''_{\varepsilon;V}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)}, \tag{2.20a}$$

$$\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{C^0(V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)};\mathbb{C}^3)} := \max_{\boldsymbol{r}\in V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)}}|\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})|\le C''\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{W^{1,6}(V^-_{(\varepsilon)};\mathbb{C}^3)}\le C''_{\varepsilon;V}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)} \tag{2.20b}$$

with a finite constant $C''_{\varepsilon;V}<+\infty$ applicable to all $\boldsymbol{F}\in\Phi(V;\mathbb{C}^3)$.

To start the proof of Eq. 2.17, we use the abbreviations $f^*:=-\nabla\cdot(\hat{\mathcal{N}}(\hat{P}\boldsymbol{F})^*)$ and $f^\sharp_{(\varepsilon)}:=-\nabla\cdot(\hat{\mathcal{N}}\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp)=-\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}$ in the following vector algebra computations

$$\begin{aligned}&[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot[\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]-[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})][\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]=f^\sharp_{(\varepsilon)}(\boldsymbol{r})[\boldsymbol{n}\cdot\nabla f^*(\boldsymbol{r})]+\boldsymbol{n}\cdot\{\nabla f^*(\boldsymbol{r})\times[\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]\}\\
=\;&f^\sharp_{(\varepsilon)}(\boldsymbol{r})[\boldsymbol{n}\cdot\nabla f^*(\boldsymbol{r})]-\boldsymbol{n}\cdot[f^*(\boldsymbol{r})\nabla\times\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]+\boldsymbol{n}\cdot\nabla\times[f^*(\boldsymbol{r})\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]\\
=\;&\boldsymbol{n}\cdot[f^*(\boldsymbol{r})\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp(\boldsymbol{r})+f^*(\boldsymbol{r})\nabla f^\sharp_{(\varepsilon)}(\boldsymbol{r})+f^\sharp_{(\varepsilon)}(\boldsymbol{r})\nabla f^*(\boldsymbol{r})]+\boldsymbol{n}\cdot\nabla\times[f^*(\boldsymbol{r})\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})],\quad\forall\boldsymbol{r}\in V^-_{(\varepsilon)}.\end{aligned}\tag{2.21}$$

As we have $\oiint_{\partial V^-_{(\varepsilon')}}\boldsymbol{n}\cdot\nabla\times[f^*(\boldsymbol{r})\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}=0,\forall\varepsilon'\in(\varepsilon,2\varepsilon_V]$ for a smooth vector field $f^*\nabla\times\boldsymbol{Q}_{(\varepsilon)}$ on a smooth boundary surface $\partial V^-_{(\varepsilon')}$, we may establish the truthfulness of Eq. 2.17 by integrating Eq. 2.21 in the following manner

$$\begin{aligned}&\oiint_{\partial V^-_{(\varepsilon)}}\{[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot[\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]-[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})][\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]\}\,\mathrm{d}\,\tilde{S}\\
=\;&\lim_{2\varepsilon_V\ge\varepsilon'\to\varepsilon+0}\oiint_{\partial V^-_{(\varepsilon')}}\boldsymbol{n}\cdot[f^*(\boldsymbol{r})\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp(\boldsymbol{r})+f^*(\boldsymbol{r})\nabla f^\sharp_{(\varepsilon)}(\boldsymbol{r})+f^\sharp_{(\varepsilon)}(\boldsymbol{r})\nabla f^*(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}\\
=\;&\lim_{2\varepsilon_V\ge\varepsilon'\to\varepsilon+0}\iiint_{V^-_{(\varepsilon')}}\{\nabla\cdot[f^*(\boldsymbol{r})\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp(\boldsymbol{r})]+\nabla\cdot[f^*(\boldsymbol{r})\nabla f^\sharp_{(\varepsilon)}(\boldsymbol{r})+f^\sharp_{(\varepsilon)}(\boldsymbol{r})\nabla f^*(\boldsymbol{r})]\}\,\mathrm{d}^3\boldsymbol{r}\\
=\;&\iiint_{V^-_{(\varepsilon)}}[\nabla f^*(\boldsymbol{r})\cdot\hat{P}\boldsymbol{F}^\sharp(\boldsymbol{r})+2\nabla f^*(\boldsymbol{r})\cdot\nabla f^\sharp_{(\varepsilon)}(\boldsymbol{r})]\,\mathrm{d}^3\boldsymbol{r}\\
=\;&\langle\hat{\mathcal{A}}_{(\varepsilon)}\hat{P}\boldsymbol{F},(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma}))\hat{\mathcal{R}}_{(\varepsilon)}\hat{P}\boldsymbol{F}^\sharp\rangle_V=\langle\hat{P}\hat{\mathcal{R}}_{(\varepsilon)}(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma}))\hat{\mathcal{A}}_{(\varepsilon)}\hat{P}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V=\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V.\end{aligned}\tag{2.22}$$

Here, in the first step of Eq. 2.22, we have employed the fact that both $\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}$ and $\nabla\times\boldsymbol{Q}_{(\varepsilon)}$ are continuous in the compact set $V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon)}$.

$\boxed{2}$ In the second step, we will estimate the convergence rate of Eq. 2.17 as

$$\left|\frac{\partial\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V}{\partial\varepsilon}\right|\le C_\star\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)},\quad\forall\boldsymbol{F},\boldsymbol{F}^\sharp\in\Phi(V;\mathbb{C}^3),\varepsilon\in(0,\varepsilon_V]\tag{2.23}$$

for some finite constant $C_\star<+\infty$ solely determined by the geometry of the volume V.

By differentiation under the integral sign on the right-hand side of Eq. 2.17, we can show that

$$\begin{aligned}\frac{\partial\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^\sharp\rangle_V}{\partial\varepsilon}=\;&\oiint_{\partial V^-_{(\varepsilon+0)}}\left\{[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot\left[\nabla\times\oiint_{\partial V^-_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^\sharp)(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]-[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})]\left[\nabla\cdot\oiint_{\partial V^-_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^\sharp)(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\right\}\mathrm{d}\,\tilde{S}\\
&+\oiint_{\partial V^-_{(\varepsilon+0)}}\{\boldsymbol{U}(\boldsymbol{r})\cdot\nabla[\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})]+\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\times\nabla\times\boldsymbol{Q}_{(\varepsilon)}(\boldsymbol{r})\}\,\mathrm{d}\,\tilde{S},\quad\forall\varepsilon\in(0,\varepsilon_V],\end{aligned}\tag{2.24}$$

where the notation "$\oiint_{\partial V^-_{(\varepsilon+0)}}$" is short-hand for "$\lim_{2\varepsilon_V\ge\varepsilon'\to\varepsilon+0}\oiint_{\partial V^-_{(\varepsilon')}}$". The first term on the right-hand side of Eq. 2.24 arises from the dependence of $\boldsymbol{r}'$ on ε, while the second term is attributed to the dependence of $\boldsymbol{r}$ on ε. Using the vector identities $\nabla(\boldsymbol{a}\cdot\boldsymbol{b})=(\boldsymbol{a}\cdot\nabla)\boldsymbol{b}+(\boldsymbol{b}\cdot\nabla)\boldsymbol{a}+\boldsymbol{a}\times(\nabla\times\boldsymbol{b})+\boldsymbol{b}\times(\nabla\times\boldsymbol{a})$, $\nabla\times(\boldsymbol{a}\times\boldsymbol{b})=\boldsymbol{a}(\nabla\cdot\boldsymbol{b})-\boldsymbol{b}(\nabla\cdot\boldsymbol{a})+(\boldsymbol{b}\cdot\nabla)\boldsymbol{a}-(\boldsymbol{a}\cdot\nabla)\boldsymbol{b}$, the geometric relations $\nabla\times\boldsymbol{n}=\boldsymbol{0},\nabla\cdot\boldsymbol{n}=-2H(\boldsymbol{r})$ (see Eq. 2.37 in S2.3.4), together with the integral expressions for $\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}$ and $\nabla\times\boldsymbol{Q}_{(\varepsilon)}$ (Eq. 2.18), we may convert Eq. 2.24 into the following form

$$\begin{aligned}
&\frac{\partial\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^{\sharp}\rangle_V}{\partial\varepsilon}\\
={}&\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\mathrm{d}\,\tilde{S}+\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\times\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}'-\boldsymbol{n})\times(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\mathrm{d}\,\tilde{S}\\
&+\oiint_{\partial V^{-}_{(\varepsilon+0)}}2H(\boldsymbol{r})\boldsymbol{U}(\boldsymbol{r})\cdot\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\mathrm{d}\,\tilde{S}+2\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\left\{\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\cdot\nabla\right]\boldsymbol{n}\right\}\mathrm{d}\,\tilde{S}. \quad (2.25)
\end{aligned}$$

Now, we may use the decomposition of normal and tangential components of a vector field to arrive at the following expression

$$\begin{aligned}
\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}' &= \boldsymbol{n}\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{\boldsymbol{n}\cdot(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'-\boldsymbol{n}\times\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{\boldsymbol{n}\times(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\\
&= \boldsymbol{n}(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})-\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})+\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{[\boldsymbol{n}(\boldsymbol{n}-\boldsymbol{n}')-\boldsymbol{n}\times(\boldsymbol{n}-\boldsymbol{n}')]\cdot(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}',
\end{aligned}$$

and combine it with the geometric identities $\boldsymbol{n}\cdot[(\boldsymbol{a}\cdot\nabla)\boldsymbol{n}]=0$ and $\hat{W}(\boldsymbol{a}):=-(\boldsymbol{a}\cdot\nabla)\boldsymbol{n}=-\{[(\boldsymbol{n}\times\boldsymbol{a})\times\boldsymbol{n}]\cdot\nabla\}\boldsymbol{n}$ (see Lemma 2.4 in S2.3.4) to rewrite Eq. 2.25 as below:

$$\begin{aligned}
&\frac{\partial\langle\hat{\mathcal{K}}_{(\varepsilon)}\boldsymbol{F},\boldsymbol{F}^{\sharp}\rangle_V}{\partial\varepsilon}\\
={}&\oiint_{\partial V^{-}_{(\varepsilon+0)}}2H(\boldsymbol{r})\boldsymbol{U}(\boldsymbol{r})\cdot[\boldsymbol{n}(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})-\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}+2\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))\,\mathrm{d}\,\tilde{S}\\
&+\oiint_{\partial V^{-}_{(\varepsilon+0)}}2H(\boldsymbol{r})\left\{[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})]\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}-\boldsymbol{n}')\cdot(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]+[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}-\boldsymbol{n}')\times(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\right\}\mathrm{d}\,\tilde{S}\\
&+2\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\hat{W}\left(\boldsymbol{n}\times\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}-\boldsymbol{n}')\times(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right)\mathrm{d}\,\tilde{S}+\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\mathrm{d}\,\tilde{S}\\
&+\oiint_{\partial V^{-}_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\times\left[\oiint_{\partial V^{-}_{(\varepsilon)}}\frac{(\boldsymbol{n}'-\boldsymbol{n})\times(\hat{P}\boldsymbol{F}^{\sharp})(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,\tilde{S}'\right]\mathrm{d}\,\tilde{S}. \quad (2.26)
\end{aligned}$$

We can prove Eq. 2.23 by deriving bound estimates, term by term, on the right-hand side of Eq. 2.26.

Using the Gauss theorem in vector calculus, we may argue that

$$\begin{aligned}
&\oiint_{\partial V^{-}_{(\varepsilon+0)}}2H(\boldsymbol{r})\boldsymbol{U}(\boldsymbol{r})\cdot[\boldsymbol{n}(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})-\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}\\
={}&\oiint_{\partial V^{-}_{(2\varepsilon_V)}}2H(\boldsymbol{r})\boldsymbol{U}(\boldsymbol{r})\cdot[\boldsymbol{n}(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})-\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}\\
&+\iiint_{V^{-}_{(\varepsilon)}\setminus V^{-}_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\{\nabla[2H(\boldsymbol{r})(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]+\nabla\times[2H(\boldsymbol{r})(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\}\,\mathrm{d}^3\boldsymbol{r},\quad\forall\varepsilon\in(0,\varepsilon_V] \quad (2.27)
\end{aligned}$$

is bounded by a certain constant multiple of $\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^{\sharp}\|_{L^2(V;\mathbb{C}^3)}$. Here, the Bergman space estimate (Eq. 2.19) is applicable to the surface integral on the right-hand side of Eq. 2.27 as

$$\begin{aligned}
&\left|\oiint_{\partial V^{-}_{(2\varepsilon_V)}}2H(\boldsymbol{r})\boldsymbol{U}(\boldsymbol{r})\cdot[\boldsymbol{n}(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})-\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\,\mathrm{d}\,\tilde{S}\right|\\
\leq{}& 2\left(\max_{\boldsymbol{r}\in\partial V^{-}_{(2\varepsilon_V)}}|H(\boldsymbol{r})|\oiint_{\partial V^{-}_{(2\varepsilon_V)}}\mathrm{d}\,\tilde{S}\right)\max_{\boldsymbol{r}\in\partial V^{-}_{(2\varepsilon_V)}}|\boldsymbol{U}(\boldsymbol{r})|\max_{\boldsymbol{r}\in\partial V^{-}_{(2\varepsilon_V)}}[|(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})|+|(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})|]\\
\leq{}& \frac{3}{2\pi}\left(\max_{\boldsymbol{r}\in\partial V^{-}_{(2\varepsilon_V)}}|H(\boldsymbol{r})|\oiint_{\partial V^{-}_{(2\varepsilon_V)}}\mathrm{d}\,\tilde{S}\right)\frac{\|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)}\left(\|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^{-}_{(\varepsilon)};\mathbb{C}^3)}+\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^{-}_{(\varepsilon)};\mathbb{C}^3)}\right)}{(2\varepsilon_V)^{3/2}(2\varepsilon_V-\varepsilon)^{3/2}}.
\end{aligned}$$

We may write this bound as $B_{\varepsilon_V;V}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)}$, where a finite constant $B_{\varepsilon_V;V}$ applies to all $\varepsilon\in(0,\varepsilon_V]$. The Cauchy-Schwarz inequality is applicable to the volume integral on the right-hand side of Eq. 2.27 as

$$\begin{aligned}
&\left|\iiint_{V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\{\nabla[2H(\boldsymbol{r})(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]+\nabla\times[2H(\boldsymbol{r})(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]\}\,\mathrm{d}^3\boldsymbol{r}\right|\\
\le\;&\|\boldsymbol{U}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C}^3)}\sqrt{\iiint_{V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}|\nabla[2H(\boldsymbol{r})(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]+\nabla\times[2H(\boldsymbol{r})(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})]|^2\,\mathrm{d}^3\boldsymbol{r}}\\
\le\;&2\|\boldsymbol{U}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C}^3)}\left\{\max_{\boldsymbol{r}\in V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}|H(\boldsymbol{r})|\left[\|\nabla(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C}^3)}+\|\nabla\times\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C}^3)}\right]\right.\\
&\left.+\max_{\boldsymbol{r}\in V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}|\nabla H(\boldsymbol{r})|\left[\|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C})}+\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)};\mathbb{C}^3)}\right]\right\}\\
\le\;&2\|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)}\left\{\max_{\boldsymbol{r}\in V\setminus V^-_{(2\varepsilon_V)}}|H(\boldsymbol{r})|\left[\|\nabla(\nabla\cdot\boldsymbol{Q}_{(\varepsilon)})\|_{L^2(V^-_{(\varepsilon)};\mathbb{C}^3)}+\|\nabla\times\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)};\mathbb{C}^3)}\right]\right.\\
&\left.+\max_{\boldsymbol{r}\in V\setminus V^-_{(2\varepsilon_V)}}|\nabla H(\boldsymbol{r})|\left[\|\nabla\cdot\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)};\mathbb{C})}+\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)};\mathbb{C}^3)}\right]\right\}\le B'_{\varepsilon_V;V}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^\sharp\|^2_{L^2(V;\mathbb{C}^3)}.
\end{aligned}$$

Employing the Gauss theorem for the second term on the right-hand side of Eq. 2.26, we find

$$\begin{aligned}
&2\oiint_{\partial V^-_{(\varepsilon+0)}}\boldsymbol{U}(\boldsymbol{r})\cdot\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))\,\mathrm{d}\tilde{S}\\
=\;&2\oiint_{\partial V^-_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))\,\mathrm{d}\tilde{S}-2\iiint_{V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))]\,\mathrm{d}^3\boldsymbol{r}.
\end{aligned}$$

To deduce bound estimates of the above expression, we may employ the principal curvature coordinate system (ξ,η) on the parallel surfaces $\partial V^-_{(\varepsilon')},\varepsilon'\in(\varepsilon,2\varepsilon_V]$ (see S2.3.1 for details) so that $\hat{W}(\partial\boldsymbol{r}/\partial\xi)=\kappa_1(\boldsymbol{r})(\partial\boldsymbol{r}/\partial\xi),\hat{W}(\partial\boldsymbol{r}/\partial\eta)=\kappa_2(\boldsymbol{r})(\partial\boldsymbol{r}/\partial\eta)$, where $\kappa_1(\boldsymbol{r})$ and $\kappa_2(\boldsymbol{r})$ are the principal curvatures satisfying $\kappa_1(\boldsymbol{r})+\kappa_2(\boldsymbol{r})=2H(\boldsymbol{r})$, $\kappa_1(\boldsymbol{r})\kappa_2(\boldsymbol{r})=K(\boldsymbol{r})$. With the Bergman space inequality (Eq. 2.19) and the geometric relation $|\hat{W}(\boldsymbol{a})|^2\le(\kappa_1^2+\kappa_2^2)|\boldsymbol{a}|^2=(4H^2-2K)|\boldsymbol{a}|^2$, we may establish the estimate

$$\begin{aligned}
&2\left|\oiint_{\partial V^-_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))\,\mathrm{d}\tilde{S}\right|\\
\le\;&2\max_{\boldsymbol{r}\in\partial V^-_{(2\varepsilon_V)}}|\boldsymbol{U}(\boldsymbol{r})|\max_{\boldsymbol{r}\in\partial V^-_{(2\varepsilon_V)}}|(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r})|\max_{\boldsymbol{r}\in\partial V^-_{(2\varepsilon_V)}}\sqrt{4H^2(\boldsymbol{r})-2K(\boldsymbol{r})}\oiint_{\partial V^-_{(2\varepsilon_V)}}\mathrm{d}\tilde{S}\\
\le\;&\frac{3}{2\pi}\frac{\|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)}\|\nabla\times\boldsymbol{Q}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)};\mathbb{C}^3)}}{(2\varepsilon_V)^{3/2}(2\varepsilon_V-\varepsilon)^{3/2}}\max_{\boldsymbol{r}\in\partial V^-_{(2\varepsilon_V)}}\sqrt{4H^2(\boldsymbol{r})-2K(\boldsymbol{r})}\oiint_{\partial V^-_{(2\varepsilon_V)}}\mathrm{d}\tilde{S}\le B''_{\varepsilon_V;V}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)}.
\end{aligned}$$

Meanwhile, with the Cauchy-Schwarz inequality, the geometric relation (see Eq. 2.43 in S2.3.4)

$$\begin{aligned}
&|\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\nabla\times\boldsymbol{Q}_{(\varepsilon)})]|\\
\le\;&\left(\sqrt{4H^2-2K}+2|H|\right)|(\boldsymbol{n}\cdot\nabla)\nabla\times\boldsymbol{Q}_{(\varepsilon)}|+2|\nabla H||\nabla\times\boldsymbol{Q}_{(\varepsilon)}|+2|H|\left(\sqrt{4H^2-2K}|\nabla\times\boldsymbol{Q}_{(\varepsilon)}|+2|\nabla\times\nabla\times\boldsymbol{Q}_{(\varepsilon)}|\right)\\
&+2\sqrt{4H^2-2K}\left[|\nabla(\boldsymbol{e}_x\cdot\nabla\times\boldsymbol{Q}_{(\varepsilon)})|+|\nabla(\boldsymbol{e}_y\cdot\nabla\times\boldsymbol{Q}_{(\varepsilon)})|+|\nabla(\boldsymbol{e}_z\cdot\nabla\times\boldsymbol{Q}_{(\varepsilon)})|\right],
\end{aligned}$$

the Calderón-Zygmund identity $\sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}\boldsymbol{Q}_{(\varepsilon)}\|^2_{L^2(\mathbb{R}^3;\mathbb{C}^3)}=\|\hat{P}\boldsymbol{F}^\sharp\|^2_{L^2(V^-_{(\varepsilon)};\mathbb{C}^3)}\le\|\boldsymbol{F}^\sharp\|^2_{L^2(V;\mathbb{C}^3)}$, we may arrive at the estimate

$$2\left|\iiint_{V^-_{(\varepsilon)}\setminus V^-_{(2\varepsilon_V)}}\boldsymbol{U}(\boldsymbol{r})\cdot\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times(\nabla\times\boldsymbol{Q}_{(\varepsilon)})(\boldsymbol{r}))]\,\mathrm{d}^3\boldsymbol{r}\right|\le B'''_{\varepsilon_V;V}\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{F}^\sharp\|_{L^2(V;\mathbb{C}^3)}.$$

Now that we have obtained a bound estimate

$$2\left|\oiint_{\partial V^-_{(2\varepsilon_V)}} H(\boldsymbol r)\boldsymbol U(\boldsymbol r)\cdot[\boldsymbol n(\nabla\cdot\boldsymbol Q_{(\varepsilon)})(\boldsymbol r)-\boldsymbol n\times(\nabla\times\boldsymbol Q_{(\varepsilon)})(\boldsymbol r)]\,\mathrm d\,\tilde S\right|+2\left|\oiint_{\partial V^-_{(\varepsilon+0)}}\boldsymbol U(\boldsymbol r)\cdot\hat W(\boldsymbol n\times(\nabla\times\boldsymbol Q_{(\varepsilon)})(\boldsymbol r))\,\mathrm d\,\tilde S\right|$$
$$\le\ (B_{\varepsilon_V;V}+B'_{\varepsilon_V;V}+B''_{\varepsilon_V;V}+B'''_{\varepsilon_V;V})\|\boldsymbol F\|_{L^2(V;\mathbb C^3)}\|\boldsymbol F^\sharp\|_{L^2(V;\mathbb C^3)},\quad\text{where } B_{\varepsilon_V;V}+B'_{\varepsilon_V;V}+B''_{\varepsilon_V;V}+B'''_{\varepsilon_V;V}<+\infty,$$

we only need to work on the remaining contributions to Eq. 2.26, which all involve a term related to

$$\oiint_{\partial V^-_{(\varepsilon)}}\frac{(\boldsymbol n'-\boldsymbol n)\cdot(\hat P\boldsymbol F^\sharp)(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d\,\tilde S'\quad\text{or}\quad\oiint_{\partial V^-_{(\varepsilon)}}\frac{(\boldsymbol n'-\boldsymbol n)\times(\hat P\boldsymbol F^\sharp)(\boldsymbol r')}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d\,\tilde S'.$$

We can eventually establish the bound estimate in Eq. 2.23 by investigating the above two surface integrals via the Calderón-Zygmund theory in harmonic analysis. The routine but lengthy derivation is deferred to S2.3.4.

Finally, we may integrate the relation in Eq. 2.23 to derive

$$\sup_{\boldsymbol F^\natural\in\Phi(V;\mathbb C^3)\setminus\{\boldsymbol 0\}}\frac{\|\hat{\mathcal K}_{(\varepsilon_V/s')}\boldsymbol F^\natural-\hat{\mathcal K}_{(\varepsilon_V/s)}\boldsymbol F^\natural\|_{L^2(V;\mathbb C^3)}}{\|\boldsymbol F^\natural\|_{L^2(V;\mathbb C^3)}}=\sup_{\boldsymbol F,\boldsymbol F^\sharp\in\Phi(V;\mathbb C^3)\setminus\{\boldsymbol 0\}}\frac{\langle\hat{\mathcal K}_{(\varepsilon_V/s')}\boldsymbol F-\hat{\mathcal K}_{(\varepsilon_V/s)}\boldsymbol F,\boldsymbol F^\sharp\rangle_V}{\|\boldsymbol F\|_{L^2(V;\mathbb C^3)}\|\boldsymbol F^\sharp\|_{L^2(V;\mathbb C^3)}}\le C_\star\varepsilon_V\left|\frac{1}{s'}-\frac{1}{s}\right|,$$

which entails the Cauchy criterion stated in Eq. 2.16, and hence the uniform convergence as claimed in Eq 2.13. $\blacksquare$

Remark The universal expression of the internal field (Eq. 2.8) may appear as striking. It shows that the complex susceptibility $\chi\in\mathbb C$ is not a well-behaved coordinate for studying the light scattering problem, whereas its rational transform $\zeta\in\chi^2/(\chi+2)$ is the appropriate choice. In other words, in the complex χ-plane, the solution $(\hat I-\chi\hat{\mathcal G})^{-1}\boldsymbol E_{\text{inc}}\in\Phi(V;\mathbb C^3)$ may not necessarily be meromorphic — there could be an inevitable barrier at $\chi=-2$ that obstructs the analytic continuation of the solution across the $\operatorname{Re}\chi$-axis. While static polarization of dieletric media invokes a Clausius-Mossotti factor $(n^2-1)/(n^2+2)=(\epsilon_r-1)/(\epsilon_r+2)$ (see Ref. [9], p. 162) that characterize the dielectric response, the dynamic polarization process in the light scattering problem seems to call for a totally different set of factors (say, $(n^2-1)/(n^2+1)$ and $(n^2-1)^2/(n^2+1)$ as appeared in Eq. 2.8). It is thus conceivable that any perturbation series for light scattering based on the Clausius-Mossotti factor $(n^2-1)/(n^2+2)$ will not converge well for a wide range of refractive indices n, because such an expansion fails to capture the correct location of the singularity in the scattering problem.

The light scattering problem for smooth dielectrics with connected exterior volume $\mathbb R^3\setminus(V\cup\partial V)$ is well-posed for all $\operatorname{Im}\chi\le0$ except (perhaps) the specific point $\chi=-2$. When the Born equation becomes ill-conditioned for $\chi=-2$, it does not fail the uniqueness theorem, but the existence theorem, leading to an unbounded inverse

$$\|(\hat I+2\hat{\mathcal G})^{-1}\|_{(\hat I+2\hat{\mathcal G})\Phi(V;\mathbb C^3)}:=\sup_{\boldsymbol E\in(\hat I+2\hat{\mathcal G})\Phi(V;\mathbb C^3)\setminus\{\boldsymbol 0\}}\frac{\|(\hat I+2\hat{\mathcal G})^{-1}\boldsymbol E\|_{L^2(V;\mathbb C^3)}}{\|\boldsymbol E\|_{L^2(V;\mathbb C^3)}}=+\infty.$$

In particular, in the case of light scattered by a spherical dielectric (Mie scattering), one can use the explicit expression of the Mie series to verify all the conclusions of the "optical resonance theorem" (Theorem 2.1), to reveal $\chi=-2$ as an accumulation point of the singularities in the Mie series, and to demonstrate the singular behavior of $(\hat I+2\hat{\mathcal G})^{-1}$ with the aid of special functions. (See S2.3.5 for detailed derivations.)

The peculiar behavior of electromagnetic scattering forms a stark contrast to the acoustic scattering problem concerning scalar waves. The acoustic scattering equation in three-dimensional space

$$((\hat I-\chi k^2\hat{\mathcal C})u)(\boldsymbol r):=u(\boldsymbol r)-\chi k^2\iiint_V\frac{u(\boldsymbol r')e^{-ik|\boldsymbol r-\boldsymbol r'|}}{4\pi|\boldsymbol r-\boldsymbol r'|}\,\mathrm d^3\boldsymbol r'=u_{\text{inc}}(\boldsymbol r),\quad u_{\text{inc}}\in L^2(V;\mathbb C)$$

involves a perturbation of the identity operator $\hat I$ by a compact operator $-\chi k^2\hat{\mathcal C}:L^2(V;\mathbb C)\longrightarrow L^2(V;\mathbb C)$, and thus has drastically different spectral properties as compared to the Born equation for light scattering. In particular, $(\hat I-\chi k^2\hat{\mathcal C})^{-1}u_{\text{inc}}$ is indeed meromorphic in the complex χ-plane, with a discrete set of isolated singularities. The critical difference between the scalar and vector wave scattering problems may well have algorithmic implications. While the scalar wave approximation has been sometimes employed to "simplify" the light scattering problem, it is clear from Theorem 2.1 that such a "simplification" may carry potential quantitative hazard, as well as qualitative misinformation, especially when used without caution in the neighborhood of the critical point $\chi=-2$. (See S2.3.6 for a detailed comparison of acoustic and electromagnetic scattering problems in various spatial dimensions.)

We have relied on the working hypothesis that the dielectric boundary ∂V is a smooth surface of class C^∞ during the proof of Theorem 2.1, but the validity of the proof remains unshaken if the smoothness condition is reduced to class

C^4. Proving the countability of the spectrum $\sigma^{\Phi}(\hat{\mathcal{G}})$ (and the related strong stability result) may require considerably more effort when the dielectric boundary ∂V ceases to be smooth of class C^4 or higher.[29] Unlike Proposition 2.4 that is largely of topological nature, the optical resonance theorem (Theorem 2.1) has a heavy geometric flavor. The smooth geometry of the dielectric boundary plays (nearly) an indispensable rôle in many critical steps (the Rellich-Kondrachov compact imbedding theorem, the bound estimate in Eq. 2.23, etc.) during the proof of Theorem 2.1.

While the "generalized optical theorem" (Theorem 1.1) may be regarded as the *energy conservation* for energetically admissible vector fields in $L^2(V;\mathbb{C}^3)$, the "optical resonance theorem" (Theorem 2.1) may qualify as a subtler result of *spectral quantization* for physically admissible vector fields in $\Phi(V;\mathbb{C}^3)$. Central to the physical admissibility criterion is the transversality condition, which has been heavily exploited (in the interior regularity estimate, Bergman space inequalities, etc.) during the establishment of the optical resonance theorem. This is remotely reminiscent of the quantization procedure of electromagnetic field in quantum field theory (see Ref. [76], p. 79 or Ref. [77], p. 23), where the transversality constraint is *condicio sine qua non*.

It is worth noting that the transversality of the internal electric field derives from one of the Maxwell equations $\nabla\cdot\boldsymbol{D}(\boldsymbol{r})=\nabla\cdot(\epsilon(\boldsymbol{r})\boldsymbol{E}(\boldsymbol{r}))=0$ and the homogeneity condition $\epsilon(\boldsymbol{r})\equiv 1+\chi,\boldsymbol{r}\in V$. For inhomogeneous media with continuously varying values of $\epsilon(\boldsymbol{r})$, the transversality constraint may not be available to the electric field $\boldsymbol{E}$, and one usually has to directly cope with the light scattering problem in a larger Hilbert space $L^2(V;\mathbb{C}^3)$. This may lead to the absence of spectral quantization for light scattered by inhomogeneous media, as pointed out in Ref. [78]. ∎

Corollary 2.2 (Strong Stability in Light Scattering) *If the smooth dielectric has a connected exterior volume* $\mathbb{R}^3\setminus(V\cup\partial V)$*, then we have* $\sigma^{\Phi}(\hat{\mathcal{G}})\cap\mathbb{R}\subset\{0,-1/2\}$ *and the related evolution semigroup* $\{\exp(-i\tau\hat{\mathcal{G}})|\tau\geq 0\}$ *is strongly stable:*

$$\lim_{\tau\to+\infty}\|\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}=0,\quad\forall\boldsymbol{F}\in\Phi(V;\mathbb{C}^3).$$

Proof According to Proposition 2.4, we have $\sigma^{\Phi}_p(\hat{\mathcal{G}})\cap\mathbb{R}=\emptyset$ if the exterior volume $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected. Using Theorem 2.1, we may combine the relation $\sigma^{\Phi}(\hat{\mathcal{G}})\setminus\sigma^{\Phi}_p(\hat{\mathcal{G}})\subset\{0,-1/2\}$ (Eq. 2.11) with $\sigma^{\Phi}_p(\hat{\mathcal{G}})\cap\mathbb{R}=\emptyset$, and obtain $\sigma^{\Phi}(\hat{\mathcal{G}})\cap\mathbb{R}\subset\{0,-1/2\}$. Now, the absence of point spectrum on the real axis $\sigma^{\Phi}_p(\hat{\mathcal{G}})\cap\mathbb{R}=\emptyset$ and the countability of the spectrum on the real axis $\sigma^{\Phi}(\hat{\mathcal{G}})\cap\mathbb{R}\subset\{0,-1/2\}$ will allow us to deduce the strong stability of the operator semigroup $\{\exp(-i\tau\hat{\mathcal{G}})|\tau\geq 0\}$, according to the Arendt-Batty-Lyubich-Vũ theorem (see Refs. [79, 80], as well as Ref. [60], pp. 326-327). In other words, the trajectory $\{\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}|\tau\geq 0\}$ not only "converges on average" to zero according the mean ergodicity result in Corollary 2.1, it also converges in L^2-norm to zero: $\lim_{\tau\to+\infty}\|\exp(-i\tau\hat{\mathcal{G}})\boldsymbol{F}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}=0,\forall\boldsymbol{F}\in\Phi(O(\mathbf{0},R);\mathbb{C}^3)$. ∎

S2.3. Additional Comments and Details

Some geometrical/topological concepts and terminologies will be introduced in S2.3.1 and S2.3.2, and serve as background materials for the proof of Proposition 2.4 regarding the evolution of transverse vector fields in smooth dielectrics. In S2.3.3, a certain class of non-smooth dielectrics are discussed, as a logical parallel to Proposition 2.4. The "optical resonance theorem" (Theorem 2.1) is supplemented by some geometric identities derived in S2.3.4, as well as extended by some discussions in S2.3.5 and S2.3.6.

S2.3.1. Tubular Neighborhoods of a Smooth Surface

In the following, we will consider some geometric properties of a volume V that is bounded and open, with its boundary ∂V being a smooth surface of class C^∞.

Definition (Smooth surface) We call a subset $\Sigma\subset\mathbb{R}^3$ a smooth surface of class C^∞ if it can be locally parametrized by well-behaved curvilinear coordinates that are continuously differentiable to arbitrarily high order.[30] More precisely, we require that for every point $\boldsymbol{r}_\star \in \Sigma$, there exists a three-dimensional open set $O_\star \subset \mathbb{R}^3$ which contains the point $\boldsymbol{r}_\star$, and a non-empty two-dimensional open rectangle $U_\star = \{(\xi,\eta) \in \mathbb{R}^2 | \xi_- < \xi < \xi_+, \eta_- < \eta < \eta_+\}$ such that every point $\boldsymbol{r} \in O_\star \cap \Sigma$ can be assigned curvilinear coordinates $\boldsymbol{r}(\xi,\eta)$ for some point $(\xi,\eta) \in U_\star$, satisfying the following properties (see Ref. [81], p. 52 for a more technically-phrased, yet practically equivalent definition):

---

[29] When the boundary smoothness is lower than order C^4, the curvature of the surface does not match up to second order at certain points. From a heuristic ray-optics perspective, these "curvature-mismatching" points may have additional "lensing capabilities", thus adding complications to the study of light-matter interaction.

[30] If we relax the condition to the existence of continuous partial derivatives up to m-th order ($m<\infty$), we have "class C^m" instead of "class C^∞".

(a) **(Differentiability)** The partial derivatives of $\boldsymbol{r}(\xi,\eta)$ are continuous up to arbitrarily high order for all $(\xi,\eta) \in U_\star$;

(b) **(Non-degeneracy)** The cross product $(\partial \boldsymbol{r}(\xi,\eta)/\partial\xi) \times (\partial \boldsymbol{r}(\xi,\eta)/\partial\eta)$ does not vanish for any $(\xi,\eta) \in U_\star$, and its normalization defines the normal vector $\boldsymbol{n}(\boldsymbol{r}) = \boldsymbol{n}(\xi,\eta)$;

(c) **(Injectivity)** If $\boldsymbol{r}(\xi,\eta) = \boldsymbol{r}(\xi',\eta')$ for $(\xi,\eta), (\xi',\eta') \in U_\star$, then $(\xi,\eta) = (\xi',\eta')$;

(d) **(Continuity of the Inverse)** If a sequence $\{\boldsymbol{r}(\xi_s,\eta_s) | s = 1, 2, \dots\} \subset O_\star \cap \Sigma$ converges to a point $\boldsymbol{r}(\xi,\eta) \in O_\star \cap \Sigma$, i.e. $\lim_{s\to\infty} |\boldsymbol{r}(\xi_s,\eta_s) - \boldsymbol{r}(\xi,\eta)| = 0$, then $\lim_{s\to\infty} \sqrt{(\xi_s - \xi)^2 + (\eta_s - \eta)^2} = 0$. ▮

Remark For a bounded open set $V \Subset \mathbb{R}^3$ with smooth boundary ∂V, the orientation of the *outward* normal $\boldsymbol{n}(\boldsymbol{r})$ on the boundary surface ∂V (i.e. pointing from the "inside part" $V \setminus \partial V$ to the "outside part" $\mathbb{R}^3 \setminus (V \cup \partial V)$) is not ambiguous — "single-sided surface" anomalies like the Möbius strip or the Klein bottle will not occur — See Ref. [82], also Ref. [27], p. 411, Ref. [81], p. 114, and Ref. [83], p. 440 for the orientability (two-sidedness) of compact surfaces. One may also view the orientability from the perspective of the Jordan-Brouwer separation theorem, which implies that any compact, smooth and connected surface $S \Subset \mathbb{R}^3$ separates its complement $\mathbb{R}^3 \setminus S$ into two connected components.

On a compact smooth surface $\Sigma = \partial V \subset \mathbb{R}^3$ endowed with the metric $\mathrm{d}\boldsymbol{r} \cdot \mathrm{d}\boldsymbol{r} = E(\xi,\eta)\,\mathrm{d}\xi^2 + 2F(\xi,\eta)\,\mathrm{d}\xi\,\mathrm{d}\eta + G(\xi,\eta)\,\mathrm{d}\eta^2$, the Laplace-Beltrami operator Δ_Σ can be expressed in local curvilinear coordinates as

$$\begin{aligned}&\Delta_\Sigma f(\boldsymbol{r}(\xi,\eta))\\ =\ &\frac{1}{\sqrt{EG-F^2}}\left[\frac{\partial}{\partial\xi}\left(\frac{G}{\sqrt{EG-F^2}}\frac{\partial f}{\partial\xi}\right) - \frac{\partial}{\partial\xi}\left(\frac{F}{\sqrt{EG-F^2}}\frac{\partial f}{\partial\eta}\right) - \frac{\partial}{\partial\eta}\left(\frac{F}{\sqrt{EG-F^2}}\frac{\partial f}{\partial\xi}\right) + \frac{\partial}{\partial\eta}\left(\frac{E}{\sqrt{EG-F^2}}\frac{\partial f}{\partial\eta}\right)\right].\end{aligned}$$

It can be verified that $\langle f, \Delta_\Sigma f\rangle_{L^2(\Sigma;\mathbb{C})} := \oiint_\Sigma f^* \Delta_\Sigma f\,\mathrm{d}S \leq 0$ for all complex-valued smooth scalar fields f. Thus, $(-\Delta_\Sigma)$ is a positive-semidefinite operator, and its power $(-\Delta_\Sigma)^s$ is well-defined for all positive-valued exponents s. In particular, for $0 < s < 1$, the fractional powers can be expressed by the Balakrishnan formula (see Ref. [44], Chap. IX, or Ref. [84]):

$$(-\Delta_\Sigma)^s f = \frac{1}{\Gamma(-s)}\int_0^{+\infty} \frac{e^{\tau\Delta_\Sigma} f - f}{\tau^{s+1}}\,\mathrm{d}\tau$$

where $u(\boldsymbol{r},\tau) = (e^{\tau\Delta_\Sigma} f)(\boldsymbol{r})$ is the unique solution to the diffusion equation $\partial u(\boldsymbol{r},\tau)/\partial\tau = \Delta_\Sigma u(\boldsymbol{r},\tau)$ with initial condition $u(\boldsymbol{r},0) = f(\boldsymbol{r})$. Moreover, if the smooth surface Σ can be decomposed into $N_\Sigma < +\infty$ connected components $\Sigma_1, \Sigma_2, \dots, \Sigma_{N_\Sigma}$, and $f(\boldsymbol{r}) = f_j(\boldsymbol{r}), \forall \boldsymbol{r} \in \Sigma_j$, then we have $(e^{\tau\Delta_\Sigma} f)(\boldsymbol{r}) = (e^{\tau\Delta_{\Sigma_j}} f_j)(\boldsymbol{r}), \forall \boldsymbol{r} \in \Sigma_j$ and $((-\Delta_\Sigma)^s f)(\boldsymbol{r}) = ((-\Delta_{\Sigma_j})^s f_j)(\boldsymbol{r}), \forall \boldsymbol{r} \in \Sigma_j$. ▮

Definition (Equidistant Sets and Parallel Surfaces) The distance $\operatorname{dist}(\boldsymbol{r}, \partial V)$ from a point $\boldsymbol{r} \in \mathbb{R}^3$ to the smooth boundary surface $\partial V \Subset \mathbb{R}^3$ is given by $T(\boldsymbol{r}) \equiv \operatorname{dist}(\boldsymbol{r}, \partial V) := \min_{\boldsymbol{r}' \in \partial V} |\boldsymbol{r} - \boldsymbol{r}'|$. It is a continuous function satisfying $|\operatorname{dist}(\boldsymbol{r}, \partial V) - \operatorname{dist}(\boldsymbol{r}^\sharp, \partial V)| \leq |\boldsymbol{r} - \boldsymbol{r}^\sharp|$. It vanishes $\operatorname{dist}(\boldsymbol{r}, \partial V) = 0$ if and only if $\boldsymbol{r} \in \partial V$. For any $\varepsilon \geq 0$, the two equidistant sets are given by

$$\Sigma^+_{(\varepsilon)} := \{\boldsymbol{r} \in \mathbb{R}^3 \setminus V | \operatorname{dist}(\boldsymbol{r}, \partial V) = \varepsilon\}, \quad \Sigma^-_{(\varepsilon)} := \{\boldsymbol{r} \in V | \operatorname{dist}(\boldsymbol{r}, \partial V) = \varepsilon\};$$

and the two parallel surfaces are given by (cf. Ref. [81], p. 212)

$$S^\pm_{(\varepsilon)} := \{\tilde{\boldsymbol{r}} \in \mathbb{R}^3 | \exists \boldsymbol{r} \in \partial V \text{ such that } \tilde{\boldsymbol{r}} = \boldsymbol{r} \pm \varepsilon \boldsymbol{n}(\boldsymbol{r})\}.$$

Here, $\Sigma^\pm_{(0)} = S^\pm_{(0)} = \partial V$. ▮

Following the spirit of the tubular neighborhood theorem in differential topology [81, 85], we claim that the equidistant sets $\Sigma^\pm_{(\varepsilon)}$ and the parallel surfaces $S^\pm_{(\varepsilon)}$ satisfy the following properties.

Proposition 2.6 *Suppose that ∂V is a smooth surface of class C^∞ satisfying properties (a)-(d), then*

- *(i) For any $\varepsilon > 0$, we have $\Sigma^{+}_{(\varepsilon)} \subset S^{+}_{(\varepsilon)}$ and $\Sigma^{-}_{(\varepsilon)} \subset S^{-}_{(\varepsilon)}$; for sufficiently small $\varepsilon > 0$, $S^{\pm}_{(\varepsilon)}$ represent two surfaces satisfying properties (a) and (b);*
- *(ii) For sufficiently small $\varepsilon > 0$, we have $\Sigma^{+}_{(\varepsilon)} = S^{+}_{(\varepsilon)}$ and $\Sigma^{-}_{(\varepsilon)} = S^{-}_{(\varepsilon)}$;*
- *(iii) For sufficiently small $\varepsilon > 0$, the sets $\Sigma^{\pm}_{(\varepsilon)} = S^{\pm}_{(\varepsilon)}$ represent two smooth surfaces of class C^∞ satisfying properties (a)-(d), and are topologically equivalent to ∂V.*

Proof It would be sufficient to demonstrate the case for $\Sigma^{-}_{(\varepsilon)}$ and $S^{-}_{(\varepsilon)}$.

Proof of (i): To start, we may fix a point $\boldsymbol{r}_{(\varepsilon)} \in \Sigma^{-}_{(\varepsilon)}$ and consider the continuously differentiable scalar field $f(\boldsymbol{r}) = (\boldsymbol{r}_{(\varepsilon)} - \boldsymbol{r}) \cdot (\boldsymbol{r}_{(\varepsilon)} - \boldsymbol{r}), \forall \boldsymbol{r} \in \partial V$ that achieves minimum $f(\boldsymbol{r}_*) = [\operatorname{dist}(\boldsymbol{r}_{(\varepsilon)}, \partial V)]^2$ for some $\boldsymbol{r}_* \in \partial V$. We may parameterize the surface ∂V in a neighborhood of $\boldsymbol{r}_*$ using non-degenerate curvilinear coordinates $\boldsymbol{r}(\xi, \eta)$ such that $(\partial \boldsymbol{r}/\partial \xi) \times (\partial \boldsymbol{r}/\partial \eta) \neq \boldsymbol{0}$ in the corresponding neighborhood. In such a neighborhood, the normal vector $\pm \boldsymbol{n}$ is parallel to $(\partial \boldsymbol{r}/\partial \xi) \times (\partial \boldsymbol{r}/\partial \eta)$. As the continuously differentiable scalar field $f(\boldsymbol{r})$ attains minimum for $\boldsymbol{r} = \boldsymbol{r}_*$, both partial derivatives $\partial f(\boldsymbol{r}(\xi, \eta))/\partial \xi$ and $\partial f(\boldsymbol{r}(\xi, \eta))/\partial \eta$ must vanish for $\boldsymbol{r}(\xi, \eta) = \boldsymbol{r}_*$. In other words, owing to the results

$$[\boldsymbol{r}_{(\varepsilon)} - \boldsymbol{r}(\xi, \eta)] \cdot \left.\frac{\partial \boldsymbol{r}(\xi, \eta)}{\partial \xi}\right|_{\boldsymbol{r}(\xi,\eta)=\boldsymbol{r}_*} = 0, \quad [\boldsymbol{r}_{(\varepsilon)} - \boldsymbol{r}(\xi, \eta)] \cdot \left.\frac{\partial \boldsymbol{r}(\xi, \eta)}{\partial \eta}\right|_{\boldsymbol{r}(\xi,\eta)=\boldsymbol{r}_*} = 0,$$

we know that $\boldsymbol{r}_* - \boldsymbol{r}_{(\varepsilon)}$ must be parallel to the outward unit normal vector $\boldsymbol{n}(\boldsymbol{r}_*)$, which is in turn parallel to $\pm(\partial \boldsymbol{r}/\partial \xi) \times (\partial \boldsymbol{r}/\partial \eta)|_{\boldsymbol{r}=\boldsymbol{r}_*}$. Therefore, we have

$$\Sigma^{-}_{(\varepsilon)} \subset S^{-}_{(\varepsilon)} := \left\{ \tilde{\boldsymbol{r}} \in \mathbb{R}^3 \,|\, \exists \boldsymbol{r} \in \partial V \text{ such that } \tilde{\boldsymbol{r}} = \boldsymbol{r} - \varepsilon \boldsymbol{n}(\boldsymbol{r}) \right\}.$$

At this point, it should be noted that although $\Sigma^{-}_{(\varepsilon)} \subset V$, there is no guarantee that $S^{-}_{(\varepsilon)} \subset V$ for arbitrarily large $\varepsilon > 0$.

We can deduce the geometric properties of $S^{-}_{(\varepsilon)}$ from that of the smooth surface ∂V. The Weingarten transformation $\hat{W}(\boldsymbol{r})$ is defined by $\mathrm{d}\boldsymbol{n} = -\hat{W}(\boldsymbol{r})\,\mathrm{d}\boldsymbol{r}$ on the surface ∂V, which can be represented in local curvilinear coordinates as

$$\begin{pmatrix} \partial \boldsymbol{n}(\xi, \eta)/\partial \xi \\ \partial \boldsymbol{n}(\xi, \eta)/\partial \eta \end{pmatrix} = -\hat{W}(\boldsymbol{r}(\xi, \eta)) \begin{pmatrix} \partial \boldsymbol{r}(\xi, \eta)/\partial \xi \\ \partial \boldsymbol{r}(\xi, \eta)/\partial \eta \end{pmatrix}. \tag{2.28}$$

Owing to the smoothness of the surface ∂V (properties (a) and (d)), the mean curvature $H(\boldsymbol{r}) := \frac{1}{2}\operatorname{tr}\hat{W}(\boldsymbol{r})$ and the Gaussian curvature $K(\boldsymbol{r}) := \det \hat{W}(\boldsymbol{r})$ are both continuous functions of $\boldsymbol{r} \in \partial V$. Thus, both $|H(\boldsymbol{r})|$ and $|K(\boldsymbol{r})|$ are bounded as $\boldsymbol{r}$ runs over the compact set ∂V (which is a bounded and closed subset of $\mathbb{R}^3$). It follows that there is a finite constant $\varepsilon' > 0$, such that $2\varepsilon|H(\boldsymbol{r})| + \varepsilon^2|K(\boldsymbol{r})| < 1/2$ holds for all $\varepsilon < \varepsilon'$ and $\boldsymbol{r} \in \partial V$. Now, for $\varepsilon < \varepsilon'$, we may introduce local curvilinear coordinates on the set $S^{-}_{(\varepsilon)}$ as $\tilde{\boldsymbol{r}}(\xi, \eta) = \boldsymbol{r}(\xi, \eta) - \varepsilon \boldsymbol{n}(\xi, \eta)$. It can be verified that (Ref. [81], p. 212)

$$\frac{\partial \tilde{\boldsymbol{r}}(\xi, \eta)}{\partial \xi} \times \frac{\partial \tilde{\boldsymbol{r}}(\xi, \eta)}{\partial \eta} = [1 + 2\varepsilon H(\boldsymbol{r}) + \varepsilon^2 K(\boldsymbol{r})] \frac{\partial \boldsymbol{r}(\xi, \eta)}{\partial \xi} \times \frac{\partial \boldsymbol{r}(\xi, \eta)}{\partial \eta} \tag{2.29}$$

does not vanish so long as $(\partial \boldsymbol{r}/\partial \xi) \times (\partial \boldsymbol{r}/\partial \eta) \neq \boldsymbol{0}$. Consequently, $S^{-}_{(\varepsilon)}$ is a surface of class C^∞ whose local curvilinear coordinates satisfy properties (a) and (b).

Proof of (ii): We need to verify that for sufficiently small ε and any given point $\tilde{\boldsymbol{r}} \in S^{-}_{(\varepsilon)}$, the continuously differentiable scalar field $f(\boldsymbol{r}) = (\tilde{\boldsymbol{r}} - \boldsymbol{r}) \cdot (\tilde{\boldsymbol{r}} - \boldsymbol{r})$ indeed achieves local minimum at any point satisfying $\boldsymbol{r} = \tilde{\boldsymbol{r}} + \varepsilon \boldsymbol{n} \in \partial V$. This can be done by evaluating the Hessian matrix $\hat{\mathcal{H}}_f(\boldsymbol{r})$ of $f(\boldsymbol{r})$ at $\boldsymbol{r} = \tilde{\boldsymbol{r}} + \varepsilon \boldsymbol{n}$:

$$\hat{\mathcal{H}}_f(\boldsymbol{r}) := \left.\begin{pmatrix} \frac{\partial^2 f(\boldsymbol{r})}{\partial \xi^2} & \frac{\partial^2 f(\boldsymbol{r})}{\partial \xi \partial \eta} \\ \frac{\partial^2 f(\boldsymbol{r})}{\partial \xi \partial \eta} & \frac{\partial^2 f(\boldsymbol{r})}{\partial \eta^2} \end{pmatrix}\right|_{\boldsymbol{r}=\tilde{\boldsymbol{r}}+\varepsilon\boldsymbol{n}} = 2 \begin{pmatrix} \varepsilon \boldsymbol{n} \cdot \frac{\partial^2 \boldsymbol{r}}{\partial \xi^2} + \frac{\partial \boldsymbol{r}}{\partial \xi} \cdot \frac{\partial \boldsymbol{r}}{\partial \xi} & \varepsilon \boldsymbol{n} \cdot \frac{\partial^2 \boldsymbol{r}}{\partial \xi \partial \eta} + \frac{\partial \boldsymbol{r}}{\partial \xi} \cdot \frac{\partial \boldsymbol{r}}{\partial \eta} \\ \varepsilon \boldsymbol{n} \cdot \frac{\partial^2 \boldsymbol{r}}{\partial \xi \partial \eta} + \frac{\partial \boldsymbol{r}}{\partial \xi} \cdot \frac{\partial \boldsymbol{r}}{\partial \eta} & \varepsilon \boldsymbol{n} \cdot \frac{\partial^2 \boldsymbol{r}}{\partial \eta^2} + \frac{\partial \boldsymbol{r}}{\partial \eta} \cdot \frac{\partial \boldsymbol{r}}{\partial \eta} \end{pmatrix}.$$

Using the notation $\tilde{\boldsymbol{r}}^\sharp = \boldsymbol{r} - \varepsilon \boldsymbol{n}(\boldsymbol{r})/2 \in S_{j,\varepsilon/2}$, and $\tilde{\boldsymbol{n}}^\sharp(\tilde{\boldsymbol{r}}^\sharp) = \boldsymbol{n}(\boldsymbol{r})$, we can rewrite the Hessian matrix $\hat{\mathcal{H}}_f(\boldsymbol{r})$ as

$$\hat{\mathcal{H}}_f(\boldsymbol{r}) = 2 \left[\begin{pmatrix} \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \xi} & \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \eta} \\ \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \eta} & \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \eta} \cdot \frac{\partial \tilde{\boldsymbol{r}}^\sharp}{\partial \eta} \end{pmatrix} + \frac{\varepsilon^2}{4} \begin{pmatrix} \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \xi} & \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \eta} \\ \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \xi} \cdot \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \eta} & \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \eta} \cdot \frac{\partial \tilde{\boldsymbol{n}}^\sharp}{\partial \eta} \end{pmatrix} \right].$$

Without loss of generality, we may choose (ξ,η) to be the principal curvature coordinates on the surface $S^-_{(\varepsilon/2)}$ that diagonalize the Weingarten transformation, so that the Hessian matrix $\hat{\mathcal{H}}_f(\boldsymbol{r})$ simplifies to

$$\hat{\mathcal{H}}_f(\boldsymbol{r})=2\left[\begin{pmatrix}1&0\\0&1\end{pmatrix}+\frac{\varepsilon^2}{4}\begin{pmatrix}\kappa_1^2(\tilde{\boldsymbol{r}}^\sharp)&0\\0&\kappa_2^2(\tilde{\boldsymbol{r}}^\sharp)\end{pmatrix}\right],$$

where the principal curvatures $\kappa_1(\tilde{\boldsymbol{r}}^\sharp)$ and $\kappa_2(\tilde{\boldsymbol{r}}^\sharp)$ satisfy (Ref. [81], p. 212)

$$\frac{\kappa_1(\tilde{\boldsymbol{r}}^\sharp)+\kappa_2(\tilde{\boldsymbol{r}}^\sharp)}{2}=H(\tilde{\boldsymbol{r}}^\sharp)=\frac{H(\boldsymbol{r})+\frac{\varepsilon}{2}K(\boldsymbol{r})}{1+\varepsilon H(\boldsymbol{r})+\frac{\varepsilon^2}{4}K(\boldsymbol{r})},\quad \kappa_1(\tilde{\boldsymbol{r}}^\sharp)\kappa_2(\tilde{\boldsymbol{r}}^\sharp)=K(\tilde{\boldsymbol{r}}^\sharp)=\frac{K(\boldsymbol{r})}{1+\varepsilon H(\boldsymbol{r})+\frac{\varepsilon^2}{4}K(\boldsymbol{r})},$$

and

$$\left|\frac{\kappa_1(\tilde{\boldsymbol{r}}^\sharp)-\kappa_2(\tilde{\boldsymbol{r}}^\sharp)}{2}\right|=\left|\frac{\sqrt{H^2(\boldsymbol{r})-K(\boldsymbol{r})}}{1+\varepsilon H(\boldsymbol{r})+\frac{\varepsilon^2}{4}K(\boldsymbol{r})}\right|.$$

It is then evident that there exists a positive constant $\varepsilon''\in(0,\varepsilon']$ such that the Hessian matrix $\hat{\mathcal{H}}_f(\boldsymbol{r})$ is positive-definite for all $\varepsilon\in(0,\varepsilon'')$ and $\boldsymbol{r}\in\partial V$. Therefore, for every point $\tilde{\boldsymbol{r}}\in S^-_{(\varepsilon)},0<\varepsilon<\varepsilon''$, we have $\operatorname{dist}(\tilde{\boldsymbol{r}},\partial V)=\min_{\boldsymbol{r}\in\partial V}|\tilde{\boldsymbol{r}}-\boldsymbol{r}|=\varepsilon$, thereby confirming that $S^-_{(\varepsilon)}\subset\Sigma^-_{(\varepsilon)},\forall\varepsilon<\varepsilon''$. In other words, for sufficiently small $\varepsilon<\varepsilon''$, we have the identity

$$\Sigma^-_{(\varepsilon)}=S^-_{(\varepsilon)}:=\left\{\tilde{\boldsymbol{r}}\in\mathbb{R}^3\,|\exists\boldsymbol{r}\in\partial V\text{ such that }\tilde{\boldsymbol{r}}=\boldsymbol{r}-\varepsilon\boldsymbol{n}(\boldsymbol{r})\right\}.$$

Proof of (iii): We claim that there exists a positive constant $\varepsilon'''\in(0,\varepsilon'')$ such that for every fixed $\varepsilon\in(0,\varepsilon''']$, the equation $\tilde{\boldsymbol{r}}=\boldsymbol{r}-\varepsilon\boldsymbol{n}(\boldsymbol{r})$ admits only one solution of $\boldsymbol{r}\in\partial V$ for each given $\tilde{\boldsymbol{r}}\in S^-_{(\varepsilon)}$. If the opposite is true, we may construct a sequence of points $\{\tilde{\boldsymbol{r}}_s=\boldsymbol{r}_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}_s)|\boldsymbol{r}_s\in\partial V,0<\varepsilon(s)<\varepsilon''/s,s=1,2,\dots\}$ and another sequence of points $\{\boldsymbol{r}'_s\in\partial V|s=1,2,\dots\}$, such that $\boldsymbol{r}_s\neq\boldsymbol{r}'_s$ but $\tilde{\boldsymbol{r}}_s=\boldsymbol{r}_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}_s)=\boldsymbol{r}'_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}'_s)$. Now that $\partial V\subset\mathbb{R}^3$ is compact, the sequence $\{\boldsymbol{r}_s\in\partial V|s=1,2,\dots\}$ must contain a convergent subsequence that converges to a point $\boldsymbol{r}_\infty\in\partial V$, according to the Bolzano-Weierstraß theorem. Consequently, a corresponding subsequence of $\{\tilde{\boldsymbol{r}}_s=\boldsymbol{r}_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}_s)|s=1,2,\dots\}$ will also converge to the same limit $\boldsymbol{r}_\infty$. In an open neighborhood $O_\infty\subset\mathbb{R}^3$ of $\boldsymbol{r}_\infty$, the local curvilinear coordinate system $\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)=\boldsymbol{r}(\xi,\eta)-\varepsilon\boldsymbol{n}(\boldsymbol{r}(\xi,\eta))$ is well-defined, as the Jacobian determinant does not vanish:

$$\left[\frac{\partial\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial\xi}\times\frac{\partial\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial\eta}\right]\cdot\frac{\partial\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial\varepsilon}\neq0$$

according to the proof of (i). Thus, for each $\tilde{\boldsymbol{r}}\in O_\infty$, the equation $\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)=\boldsymbol{r}(\xi,\eta)-\varepsilon\boldsymbol{n}(\boldsymbol{r}(\xi,\eta))$ admits only one solution of $\boldsymbol{r}\in\partial V$, according to the implicit function theorem. On the other hand, by our assumption of the non-existence of the constant $\varepsilon'''>0$, the neighborhood O_∞ contains an infinite subsequence of $\{\tilde{\boldsymbol{r}}_s=\boldsymbol{r}_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}_s)=\boldsymbol{r}'_s-\varepsilon(s)\boldsymbol{n}(\boldsymbol{r}'_s)|\boldsymbol{r}_s,\boldsymbol{r}'_s\in\partial V,\boldsymbol{r}_s\neq\boldsymbol{r}'_s,s=1,2,\dots\}$ where the solution to $\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)=\boldsymbol{r}(\xi,\eta)-\varepsilon\boldsymbol{n}(\boldsymbol{r}(\xi,\eta))$ is not unique. This contradiction shows that the positive constant $\varepsilon'''\in(0,\varepsilon'')$ must exist. Therefore, for $\varepsilon\leq\varepsilon'''$, the mapping $\boldsymbol{r}\in\partial V\mapsto\tilde{\boldsymbol{r}}=\boldsymbol{r}-\varepsilon\boldsymbol{n}\in S^-_{(\varepsilon)}$ is a continuous bijection from a compact set ∂V to a compact set $S^-_{(\varepsilon)}$, and its inverse mapping must also be continuous. Accordingly, the surface $S^-_{(\varepsilon)}=\Sigma^-_{(\varepsilon)},0\leq\varepsilon\leq\varepsilon'''$ is topologically equivalent to ∂V, and is a smooth surface of class C^∞ in its own right.[31] In particular, if ∂V is connected, then so must be the surface $S^-_{(\varepsilon)}=\Sigma^-_{(\varepsilon)}$ whenever $0\leq\varepsilon\leq\varepsilon'''$.

As a consequence of the results proved in steps (i)-(iii), we see that the spatial region formed by a stack of parallel surfaces $\bigcup_{0\leq\varepsilon\leq\varepsilon'''}S^-_{(\varepsilon)}=\bigcup_{0\leq\varepsilon\leq\varepsilon'''}\Sigma^-_{(\varepsilon)}$ is topologically equivalent to the Cartesian product $\partial V\times[0,\varepsilon''']$: a sequence of points $\{\tilde{\boldsymbol{r}}_s|s=1,2,\dots\}\subset\bigcup_{0\leq\varepsilon\leq\varepsilon'''}S^-_{(\varepsilon)}$ converges to a point $\tilde{\boldsymbol{r}}\in\bigcup_{0\leq\varepsilon\leq\varepsilon'''}S^-_{(\varepsilon)}$ if and only if the respective local curvilinear coordinates converge $\lim_{j\to\infty}\sqrt{(\xi_s-\xi)^2+(\eta_s-\eta)^2+(\varepsilon_s-\varepsilon)^2}=0$. The three-dimensional curvilinear coordinate system $\tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)=\boldsymbol{r}(\xi,\eta)-\varepsilon\boldsymbol{n}(\boldsymbol{r}(\xi,\eta))$ is hence well-behaved in the following three-dimensional open neighborhood of ∂V (the "tubular neighborhood"[32])

$$\partial V\cup\left[\bigcup_{0<\varepsilon_*<2\varepsilon}S^-_{(\varepsilon_*)}\right]\cup\left[\bigcup_{0<\varepsilon_*<2\varepsilon}S^+_{(\varepsilon_*)}\right]\tag{2.30}$$

---

[31] If we start with a surface ∂V of class C^m where $m<\infty$, then the parallel surfaces are of class C^{m-1}.

[32] The term "tubular" may sound more appropriate if we replace the smooth surface ∂V by a smooth curve. It is merely a convention to name certain surroundings of a smooth manifold in any dimension (smooth curves, smooth surfaces, etc.) as a "tubular neighborhood".

and the local curvilinear coordinate system $\tilde{\boldsymbol{r}} : (\xi, \eta, \varepsilon) \mapsto V^-_{(\varepsilon''')}$ defines a smooth vector field of type $C^\infty(U_\star \times (-\varepsilon''', \varepsilon'''); \mathbb{R}^3)$. As a result, the unit normal vectors $\boldsymbol{n}(\tilde{\boldsymbol{r}}) = \boldsymbol{n}(\xi, \eta, \varepsilon)$ also forms a smooth vector field of type $C^\infty(U_\star \times (-\varepsilon''', \varepsilon'''); \mathbb{R}^3)$. ▮

Remark Here, in the proof of Proposition 2.6, the constraint "sufficiently small $\varepsilon > 0$" cannot be dispensed, in either the proof of (i), (ii) or (iii). In fact, suppose that $V = O(\mathbf{0}, R)$ is an open-ball neighborhood of the origin with radius R, then $\partial(V \setminus V_{(R)}) = V \setminus V_{(R)} = \{\mathbf{0}\}$ fails to be a smooth surface. One may also pick certain volumes V of non-convex shape, so that for some finitely large ε, the set $\Sigma^-_{(\varepsilon)}$ is not connected while $S^-_{(\varepsilon)}$ is connected, and at a certain point $\tilde{\boldsymbol{r}} \in S^-_{(\varepsilon)}$, the function $f(\boldsymbol{r}) = (\tilde{\boldsymbol{r}} - \boldsymbol{r}) \cdot (\tilde{\boldsymbol{r}} - \boldsymbol{r}), \boldsymbol{r} \in \partial V$ attains local maximum (as opposed to minimum). When V is not convex, it is also possible that either $\Sigma^-_{(\varepsilon)}$ or $S^-_{(\varepsilon)}$ is self-intersecting for some finitely large ε.

The expository proof presented here is no more than a modest extension of the geometric arguments in pp. 109-114 and p. 212 of Ref. [81], as well as a specific case of the abstract "tubular neighborhood theorem" in pp. 109-118 of Ref. [85]. We have adapted the standard geometric and topological arguments in Proposition 2.6, so as to elicit a self-contained discussion herein about the geometric and physical consequences of a smooth dielectric boundary. ▮

Intuitively speaking, one may expect that a surface does not occupy a positive volume in $\mathbb{R}^3$, and a bounded surface has finite area. However, in Ref. [86], A. S. Besicovitch has constructed a surface $S_\times \subset \mathbb{R}^3$ that is topologically equivalent to the boundary of an open ball $\partial O(\mathbf{0}, R) \subset \mathbb{R}^3, R > 0$, with finite area and positive volume $\iiint_{S_\times} \mathrm{d}^3\boldsymbol{r} > 0$. Meanwhile, fractal structures like the Menger sponge may exhibit infinite surface area despite being a compact set in the three dimensional space $\mathbb{R}^3$.

The following corollary of the tubular neighborhood theorem shows that a compact smooth surface will not have a counter-intuitively positive volume as in the Besicovitch construction, neither will it have infinite surface area as a Menger sponge.

Corollary 2.3 *If $\Sigma \Subset \mathbb{R}^3$ is a compact and smooth surface of class C^∞, then the volume of Σ vanishes: $\iiint_\Sigma \mathrm{d}^3\tilde{\boldsymbol{r}} = 0$, and its surface area is finite $\oiint_\Sigma \mathrm{d}S < +\infty$.*

Proof First, we consider a surface neighborhood $O_\star \cap \Sigma$ of a point $\boldsymbol{r}_\star \in \Sigma$ on the compact and smooth surface $\Sigma \Subset \mathbb{R}^3$ that corresponds to a local coordinate chart $U_\star \subset \mathbb{R}^2$. With the well-behaved local curvilinear coordinate system $\tilde{\boldsymbol{r}}(\xi, \eta, \varepsilon) = \boldsymbol{r}(\xi, \eta) - \varepsilon \boldsymbol{n}(\boldsymbol{r}(\xi, \eta))$ derived from the tubular neighborhood theorem, we may compute the volume of $O_\star \cap \Sigma$ as

$$\iiint_{O_\star \cap \Sigma} \mathrm{d}^3\tilde{\boldsymbol{r}} = \iiint_{(\xi,\eta)\in U_\star, \varepsilon=0} \left| \left[\frac{\partial \tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial \xi} \times \frac{\partial \tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial \eta} \right] \cdot \frac{\partial \tilde{\boldsymbol{r}}(\xi,\eta,\varepsilon)}{\partial \varepsilon} \right| \mathrm{d}\xi \, \mathrm{d}\eta \, \mathrm{d}\varepsilon = 0,$$

because the domain of integration $U_\star \times \{0\} = \{(\xi, \eta, \varepsilon) \in \mathbb{R}^3 | (\xi, \eta) \in U_\star, \varepsilon = 0\}$ has zero Lebesgue measure: $\iiint_{U_\star \times \{0\}} \mathrm{d}\xi \, \mathrm{d}\eta \, \mathrm{d}\varepsilon = 0$. Then, we note that $\bigcup_{\boldsymbol{r}_\star \in \Sigma} O_\star \supset \Sigma$ forms an open cover of the compact set $\Sigma \Subset \mathbb{R}^3$. According to the Heine-Borel theorem, we may pick a finite subcover, i.e. there exists a subset $F = \{\boldsymbol{r}_1, \boldsymbol{r}_2, \dots, \boldsymbol{r}_N\} \subset \Sigma$ consisting of finitely many points, such that $\bigcup_{\boldsymbol{r}_\star \in F} O_\star = \bigcup_{j=1}^N O_j \supset \Sigma$ and $\bigcup_{j=1}^N (O_j \cap \Sigma) = \Sigma$. The volume of Σ thus will not exceed the sum of the volumes of $N < +\infty$ subsets $O_j \cap \Sigma$:

$$\iiint_\Sigma \mathrm{d}^3\tilde{\boldsymbol{r}} \le \sum_{j=1}^N \iiint_{O_j \cap \Sigma} \mathrm{d}^3\tilde{\boldsymbol{r}} = 0.$$

Here, we have used the facts that each $O_j \cap \Sigma$ has zero volume, and the sum of finitely many zeros is zero.

Now, consider a tubular neighborhood of Σ that is topologically equivalent to $\Sigma \times [-\delta, \delta]$, $\delta > 0$, parametrized by well-behaved curvilinear coordinates $\tilde{\boldsymbol{r}} = \boldsymbol{r} + \varepsilon \boldsymbol{n}(\boldsymbol{r}), \boldsymbol{r} \in \Sigma, \varepsilon \in [-\delta, \delta]$. Being topologically equivalent to the Cartesian product of compact sets Σ and $[-\delta, \delta]$, the tubular neighborhood in question is a compact set with finite volume:

$$\int_{-\delta}^{\delta} \left\{ \oiint_\Sigma [1 - 2\varepsilon H(\boldsymbol{r}) + \varepsilon^2 K(\boldsymbol{r})] \, \mathrm{d}S \right\} \mathrm{d}\varepsilon = \iiint_{\tilde{\boldsymbol{r}} = \boldsymbol{r} + \varepsilon \boldsymbol{n}(\boldsymbol{r}), \boldsymbol{r} \in \Sigma, \varepsilon \in [-\delta, \delta]} \mathrm{d}^3\tilde{\boldsymbol{r}} < +\infty.$$

One may always pick $\delta > 0$ sufficiently small so that $1/2 \le 1 - 2\varepsilon H(\boldsymbol{r}) + \varepsilon^2 K(\boldsymbol{r}), \forall \varepsilon \in [-\delta, \delta]$, which shows that

$$\oiint_\Sigma \mathrm{d}S \le \frac{1}{\delta} \iiint_{\tilde{\boldsymbol{r}} = \boldsymbol{r} + \varepsilon \boldsymbol{n}(\boldsymbol{r}), \boldsymbol{r} \in \Sigma, \varepsilon \in [-\delta, \delta]} \mathrm{d}^3\tilde{\boldsymbol{r}} < +\infty. \quad ▮$$

As another consequence of the tubular neighborhood theorem, we claim that if V is also connected, then it remains connected under the perturbation by parallel surfaces, as shown below.

Corollary 2.4 *For a connected, open and bounded set* V *with smooth boundary (which may or may not be connected), one may pick sufficiently small* $\varepsilon \in (0, \varepsilon'''/2)$, *so that the following two sets are connected*

$$V \setminus \left[\bigcup_{0<\varepsilon_* \le \varepsilon} S^-_{(\varepsilon_*)} \right], V \cup \left[\bigcup_{0 \le \varepsilon_* < 2\varepsilon} S^+_{(\varepsilon_*)} \right].$$

Proof Consider the mapping $h_- : V \longrightarrow V \setminus \left[\bigcup_{0<\varepsilon_* \le \varepsilon} S^-_{(\varepsilon_*)} \right]$ as

$$h_-(\boldsymbol{r}) = \begin{cases} \boldsymbol{r} + (\varepsilon + \frac{1}{2}\varepsilon')\boldsymbol{n}(\boldsymbol{r}'), & \boldsymbol{r}' = \boldsymbol{r} + \varepsilon'\boldsymbol{n}(\boldsymbol{r}') \in \partial V, 0 < \varepsilon' < 2\varepsilon \\ \boldsymbol{r}, & \text{otherwise.} \end{cases}$$

From the proof of (iii) in Proposition 2.6, we know that the mapping h_- is continuous. As the image of a connected set V under a continuous mapping h_-, the set $V \setminus \left[\bigcup_{0<\varepsilon_* \le \varepsilon} S^-_{(\varepsilon_*)} \right] = h_-(V)$ is connected. Similarly, we may construct a continuous mapping $h_+ : V \longrightarrow V \cup \left[\bigcup_{0 \le \varepsilon_* < 2\varepsilon} S^+_{(\varepsilon_*)} \right]$ as

$$h_+(\boldsymbol{r}) = \begin{cases} \boldsymbol{r} + (2\varepsilon + 4\varepsilon')\boldsymbol{n}(\boldsymbol{r}'), & \boldsymbol{r}' = \boldsymbol{r} + \varepsilon'\boldsymbol{n}(\boldsymbol{r}') \in \partial V, 0 < \varepsilon' < 2\varepsilon \\ \boldsymbol{r}, & \text{otherwise,} \end{cases}$$

to show that $V \cup \left[\bigcup_{0 \le \varepsilon_* < 2\varepsilon} S^+_{(\varepsilon_*)} \right] = h_+(V)$ is connected. ▮

S2.3.2. Connectedness in the Perspectives of de Rham and Hodge

For an effective solution to the light scattering problem, we usually require a connected exterior volume $\mathbb{R}^3 \setminus (V \cup \partial V)$. The following lemma shows that this topological constraint is closely related to the requirement that each connected component of V has a connected boundary.

Lemma 2.3 (Alexander Relation) *Assume that a bounded open set* $V = \bigcup_{j=1}^{N_V} V_j \Subset O(\mathbf{0}, R/2)$ *has finitely many connected components* $V_1, \dots, V_{N_V}$, *and each connected component has a smooth boundary surface* $\partial V_j, j = 1, \dots, N_V$ *that does not intersect the boundary of a distinct component* $\partial V_j \cap \partial V_{j'} = \emptyset, 1 \le j < j' \le N_V$. *The exterior volume* $\mathbb{R}^3 \setminus (V \cup \partial V)$ *is connected, if and only if each component* $V_j, j = 1, \dots, N_V$ *has a connected boundary* ∂V_j.

Proof We will prove the conclusion *a fortiori*, namely, we will use the Mayer-Vietoris sequence in de Rham cohomology (see Ref. [27], p. 408-411) to provide an algebraic identity

$$\dim \mathcal{H}^0(V \cup \partial V; \mathbb{R}) + \dim \mathcal{H}^0(\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V; \mathbb{R}) - \dim \mathcal{H}^0(\partial V; \mathbb{R}) = 1, \tag{2.31}$$

where $\dim \mathcal{H}^0(\overline{M}; \mathbb{R})$ counts the number of connected components in a compact manifold $\overline{M}$. Here, a compact set $\overline{M} \Subset \mathbb{R}^3$ qualifies as a d-dimensional compact manifold if for every point $\boldsymbol{r}_\star \in \overline{M}$, there exists an open set $O_\star \subset \mathbb{R}^3$, such that $\boldsymbol{r}_\star \in O_\star$ and the neighborhood $O_\star \cap \overline{M}$ is topologically equivalent to either $\mathbb{R}^d$ or $\mathbb{R}^d_+ := \mathbb{R}^{d-1} \times [0, +\infty)$. In particular, both $V \cup \partial V$ and $\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V$ are 3-dimensional compact manifolds, and ∂V is a 2-dimensional compact manifold.

It can be recognized that $\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V$ is connected if and only if $\mathbb{R}^3 \setminus (V \cup \partial V)$ is connected.[33] Thus, according to Eq. 2.31, the exterior volume is connected $1 = \dim \mathcal{H}^0(\mathbb{R}^3 \setminus (V \cup \partial V), \mathbb{R}) = \dim \mathcal{H}^0(\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V, \mathbb{R})$ if and only if $V \cup \partial V$ has the same number of connected components as that of ∂V, i.e. $\dim \mathcal{H}^0(V \cup \partial V, \mathbb{R}) = \dim \mathcal{H}^0(\partial V, \mathbb{R})$.

---

[33] First we point out that as a compact manifold with smooth boundary, the set $\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V \subset \mathbb{R}^3$ is connected if and only if its interior $O(\mathbf{0}, R) \setminus (V \cup \partial V)$ is, as can be seen from the isomorphisms $\mathcal{H}^0(\mathrm{Cl}\,(O(\mathbf{0}, R)) \setminus V; \mathbb{R}) \cong \mathcal{H}^3_c(O(\mathbf{0}, R) \setminus (V \cup \partial V); \mathbb{R}) \cong \mathcal{H}^0(O(\mathbf{0}, R) \setminus (V \cup \partial V); \mathbb{R})$. There is a continuous transformation from $\mathbb{R}^3 \setminus (V \cup \partial V)$ to $O(\mathbf{0}, R) \setminus (V \cup \partial V)$ that keeps any point $\boldsymbol{r} \in O(\mathbf{0}, 2R/3)$ intact, and transforms a point $\boldsymbol{r} \notin O(\mathbf{0}, 2R/3)$ to $3R\boldsymbol{r}/(3|\boldsymbol{r}| + R)$. The inverse transformation that keeps any point $\boldsymbol{r} \in O(\mathbf{0}, 2R/3)$ intact, and transforms a point $\boldsymbol{r} \notin O(\mathbf{0}, 2R/3)$ to $R\boldsymbol{r}/(3R - 3|\boldsymbol{r}|)$, is also continuous. Hence, the open set $O(\mathbf{0}, R) \setminus (V \cup \partial V)$ is topologically equivalent to $\mathbb{R}^3 \setminus (V \cup \partial V)$, so that the connectedness of one implies that of the other.

Now we prove Eq. 2.31. As the smooth boundary surface $\partial V_j, j=1,\dots,N_V<+\infty$ does not intersect the boundary of a distinct component $\partial V_j\cap\partial V_{j'}=\emptyset, 1\le j<j'\le N_V$, the union of all boundary surfaces $\partial V=\bigcup_{j=1}^{N_V}\partial V_j$ is a 2-dimensional smooth and compact manifold in its own right. Picking two compact manifolds with boundary $\overline{M}_1=\mathrm{Cl}\,(O(\mathbf{0},R))\setminus V$ and $\overline{M}_2=V\cup\partial V$, we have the Mayer-Vietoris sequence

$$\{\mathbf{0}\}\longrightarrow\mathcal{H}^0(\overline{M}_1\cup\overline{M}_2;\mathbb{R})\longrightarrow\mathcal{H}^0(\overline{M}_1;\mathbb{R})\oplus\mathcal{H}^0(\overline{M}_2;\mathbb{R})\longrightarrow\mathcal{H}^0(\overline{M}_1\cap\overline{M}_2;\mathbb{R})\longrightarrow\mathcal{H}^1(\overline{M}_1\cup\overline{M}_2;\mathbb{R})=\{\mathbf{0}\}.$$

Here, $\overline{M}_1\cup\overline{M}_2=\mathrm{Cl}\,(O(\mathbf{0},R))$ can be contracted to a single point, so it has trivial de Rham cohomology groups $\mathcal{H}^m(\overline{M}_1\cup\overline{M}_2;\mathbb{R})=\{\mathbf{0}\}$ for positive integers m. It is clear that $\overline{M}_1\cup\overline{M}_2=\mathrm{Cl}\,(O(\mathbf{0},R))$ is connected, so $\dim\mathcal{H}^0(\overline{M}_1\cup\overline{M}_2;\mathbb{R})=1$. Meanwhile, $\dim\mathcal{H}^0(\overline{M}_2;\mathbb{R})=\dim\mathcal{H}^0(V\cup\partial V;\mathbb{R})=N_V$ counts the number of connected components in $V\cup\partial V$.

As the Mayer-Vietoris sequence is exact, we know that the alternating sum of the dimensions of the vector spaces in the sequence must vanish:

$$\begin{aligned}0&=\dim\mathcal{H}^0(\overline{M}_1\cup\overline{M}_2;\mathbb{R})-\dim[\mathcal{H}^0(\overline{M}_1;\mathbb{R})\oplus\mathcal{H}^0(\overline{M}_2;\mathbb{R})]+\dim\mathcal{H}^0(\overline{M}_1\cap\overline{M}_2;\mathbb{R})\\&=1-[\dim\mathcal{H}^0(\mathrm{Cl}\,(O(\mathbf{0},R))\setminus V;\mathbb{R})+\dim\mathcal{H}^0(V\cup\partial V;\mathbb{R})]+\dim\mathcal{H}^0(\partial V;\mathbb{R}).\end{aligned}$$

This establishes Eq. 2.31 as claimed. ■

Remark In the literature, an equivalent form of Eq. 2.31, written as "number of connected components of V" + "number of *bounded* connected components of $\mathbb{R}^3\setminus(V\cup\partial V)$" = "number of connected components of ∂V", has been known as the Alexander relation (see Ref. [65], pp. 384-387 for an analytic proof of the Alexander relation using potential theory). ■

Although the rotation of a vector field is necessarily divergence-free, i.e. $\mathrm{ran}\,(\nabla\times)\subset\ker(\nabla\cdot)$ holds for smooth vector fields defined on any open set $V\subset\mathbb{R}^3$, the reverse implication $\ker(\nabla\cdot)\subset\mathrm{ran}\,(\nabla\times)$ only holds when V satisfies some specific topological properties. In parallel, the gradient of a compactly supported smooth scalar field is necessarily an irrotational and compactly supported vector field, i.e. $\{\nabla f|f\in C_0^\infty(V;\mathbb{R}^3)\}\subset\ker(\nabla\times)\cap C_0^\infty(V;\mathbb{R}^3)$ for any open set $V\subset\mathbb{R}^3$, we still need additional topological constraints on V before we can assert that any compactly supported smooth irrotational field is the gradient of a compactly supported scalar field.

In the following, without loss of generality, we may momentarily confine our discussion to real-valued vector/scalar fields defined on 3-dimensional compact and connected manifolds imbedded in $\mathbb{R}^3$.

Proposition 2.7 *Let $M\subset\mathbb{R}^3$ be a bounded, open and connected set with a smooth boundary surface ∂M. The relation* $\ker(\nabla\cdot)=\mathrm{ran}\,(\nabla\times)$ *holds for smooth vector fields defined on $M\cup\partial M$, if and only if the boundary ∂M is connected.*

Proof We recall the fact that if a bounded, open and connected set $M\subset\mathbb{R}^3$ with smooth boundary ∂M has a vanishing second Betti number $b_2(M\cup\partial M)=0$, then every divergence-free vector field $\boldsymbol{F}\in\ker(\nabla\cdot)$ on $M\cup\partial M$ can be represented as the rotation of another vector field $\boldsymbol{F}=\nabla\times\boldsymbol{G}\in\mathrm{ran}\,(\nabla\times)$ (see Ref. [27], p. 369), and vice versa. Here, the second Betti number $b_2(M\cup\partial M):=\dim\mathcal{H}^2(M\cup\partial M;\mathbb{R})$ is the dimensionality of $\mathcal{H}^2(M\cup\partial M;\mathbb{R}):=\ker(\nabla\cdot)/\mathrm{ran}\,(\nabla\times)$, the second de Rham cohomology group of $M\cup\partial M$. We will show that the boundary ∂M is connected if and only if $b_2(M\cup\partial M)=0$.

According to the Hodge theory (see Ref. [27], p. 369), the vector space $\mathcal{H}^2(M\cup\partial M;\mathbb{R})$ is isomorphic to $\mathcal{H}_2^A(M\cup\partial M;\mathbb{R})$, the totality of solutions to the vector harmonic equation subject to certain boundary conditions:

$$\mathcal{H}_2^A(M\cup\partial M;\mathbb{R})=\{\boldsymbol{U}\in C^\infty(M\cup\partial M;\mathbb{R}^3)|\nabla^2\boldsymbol{U}(\boldsymbol{r})=\mathbf{0},\forall\boldsymbol{r}\in M\cup\partial M;\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})=\mathbf{0},\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in\partial M\}.$$

Here, the relation $\nabla^2\boldsymbol{U}=-\nabla\times\nabla\times\boldsymbol{U}-\nabla(\nabla\cdot\boldsymbol{U})$ is consistent with the definition of the Hodge-Laplace operator on differential 2-forms.

We first claim that every member $\boldsymbol{U}\in\mathcal{H}_2^A(M\cup\partial M;\mathbb{R})$ must be divergence-free, i.e. $\mathcal{H}_2^A(M\cup\partial M;\mathbb{R})\subset\ker(\nabla\cdot)$. This is because the scalar Laplace equation

$$0=\nabla\cdot\nabla^2\boldsymbol{U}(\boldsymbol{r})=-\nabla^2[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})],\quad\forall\boldsymbol{r}\in M\cup\partial M$$

and the vanishing boundary condition $\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in\partial M$ together imply that

$$0=\oint\!\!\!\oint_{\partial M}[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})](\boldsymbol{n}\cdot\nabla)[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]\,\mathrm{d}S=\iiint_M\{[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]\nabla^2[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]+|\nabla[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]|^2\}\,\mathrm{d}^3\boldsymbol{r}=\iiint_M|\nabla[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]|^2\,\mathrm{d}^3\boldsymbol{r}.$$

Here, the gradient $\nabla[\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]$ vanishes identically. We thus have $\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in M$ owing to the vanishing boundary condition and the vanishing gradient. The transversality result $\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in M$ is not totally unexpected, because the isomorphism $\mathcal{H}^2(M\cup\partial M;\mathbb{R})\cong\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$ in Hodge theory gives rise to a representation of every divergence-free field $\boldsymbol{F}\in\ker(\nabla\cdot)\cap C^\infty(M\cup\partial M;\mathbb{R})$ as $\boldsymbol{F}=\nabla\times\boldsymbol{G}+\boldsymbol{U},\boldsymbol{G}\in C^\infty(M\cup\partial M;\mathbb{R}),\boldsymbol{U}\in\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$, i.e. the rotation of a smooth vector field plus a member in $\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$. This necessarily implies that $\boldsymbol{U}\in\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$ is divergence-free: $\nabla\cdot\boldsymbol{U}=\nabla\cdot\boldsymbol{F}-\nabla\cdot\nabla\times\boldsymbol{G}=0$.

We then claim that every member $\boldsymbol{U}\in\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$ must also be irrotational, i.e. $\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})\subset\ker(\nabla\times)$. To show this, we use the Green identity for vector fields (see Ref. [14], p. 155 or Ref. [27], p. 361) to deduce

$$\iiint_M[\boldsymbol{U}(\boldsymbol{r})\cdot\nabla^2\boldsymbol{U}(\boldsymbol{r})+|\nabla\times\boldsymbol{U}(\boldsymbol{r})|^2+|\nabla\cdot\boldsymbol{U}(\boldsymbol{r})|^2]\,\mathrm{d}^3\boldsymbol{r}=\oiint_{\partial M}\{[\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})]\cdot[\nabla\times\boldsymbol{U}(\boldsymbol{r})]+[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})][\nabla\cdot\boldsymbol{U}(\boldsymbol{r})]\}\,\mathrm{d}S,$$

which now implies

$$\iiint_M|\nabla\times\boldsymbol{U}(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}=0,\quad\text{i.e. }\nabla\times\boldsymbol{U}(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in M\cup\partial M.$$

Therefore, for any $\boldsymbol{U}\in\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$ satisfying the vector harmonic equation subject to the boundary conditions $\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})=\boldsymbol{0},\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in\partial M$, the Green identity

$$\boldsymbol{U}(\boldsymbol{r})=\oiint_{\partial M}\left[\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}(\boldsymbol{n}'\cdot\nabla')\boldsymbol{U}(\boldsymbol{r}')-\boldsymbol{U}(\boldsymbol{r}')(\boldsymbol{n}'\cdot\nabla')\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\mathrm{d}S',\quad\forall\boldsymbol{r}\in M$$

may be modified into the following "Kirchhoff diffraction integral" (see the argument in Ref. [9], p. 483):

$$\begin{aligned}\boldsymbol{U}(\boldsymbol{r})&=-\oiint_{\partial M}\left\{\frac{\boldsymbol{n}'\times[\nabla'\times\boldsymbol{U}(\boldsymbol{r}')]}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}+[\boldsymbol{n}'\times\boldsymbol{U}(\boldsymbol{r}')]\times\nabla'\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}+[\boldsymbol{n}'\cdot\boldsymbol{U}(\boldsymbol{r}')]\nabla'\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right\}\mathrm{d}S'\\&=-\oiint_{\partial M}[\boldsymbol{n}'\cdot\boldsymbol{U}(\boldsymbol{r}')]\nabla'\frac{1}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}S',\quad\forall\boldsymbol{r}\in M,\end{aligned}$$

where $\boldsymbol{n}$ is the outward unit normal vector on the boundary ∂M.

Now, we see that $\boldsymbol{U}(\boldsymbol{r})=\nabla u(\boldsymbol{r})$ is a gradient field, where

$$u(\boldsymbol{r})=\oiint_{\partial M}\frac{\boldsymbol{n}'\cdot\boldsymbol{U}(\boldsymbol{r}')}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}S',\quad\forall\boldsymbol{r}\in M.$$

The boundary condition $\boldsymbol{n}\times\boldsymbol{U}(\boldsymbol{r})=\boldsymbol{n}\times\nabla u(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in\partial M$ entails that $u(\boldsymbol{r})\equiv u_0,\forall\boldsymbol{r}\in\partial M$ is a constant at the boundary surface, so long as the boundary ∂M is connected. Therefore, with the scalar Laplace equation $\nabla^2u(\boldsymbol{r})=\nabla\cdot\boldsymbol{U}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in M\cup\partial M$, we obtain

$$0=\oiint_{\partial M}[u(\boldsymbol{r})-u_0](\boldsymbol{n}\cdot\nabla)u(\boldsymbol{r})\,\mathrm{d}S=\iiint_M\{[u(\boldsymbol{r})-u_0]\nabla^2u(\boldsymbol{r})+|\nabla u(\boldsymbol{r})|^2\}\,\mathrm{d}^3\boldsymbol{r}=\iiint_M|\nabla u(\boldsymbol{r})|^2\,\mathrm{d}^3\boldsymbol{r}.$$

This eventually shows that $\boldsymbol{U}(\boldsymbol{r})=\nabla u(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in M\cup\partial M$, for whatever $\boldsymbol{U}\in\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$. Hence, the second Betti number $b_2(M\cup\partial M):=\dim\mathcal{H}^2(M\cup\partial M;\mathbb{R})=\dim\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})=0$ has to vanish when the boundary ∂M is connected.

Conversely, if ∂M is not connected, we can always pick a scalar field $f(\boldsymbol{r}),\boldsymbol{r}\in M\cup\partial M$ that satisfies the Laplace equation $\nabla^2f(\boldsymbol{r})=0,\forall\boldsymbol{r}\in M$ and assumes a distinct constant value on each connected component of the boundary ∂M. It can be verified that with thus prescribed boundary conditions on a smooth boundary, the harmonic function f is uniquely determined (see Theorem 3.2.2 in Ref. [30] or Corollary 11.13 in Ref. [74] for the solvability of the corresponding Dirichlet problem), and $f\in C^\infty(M\cup\partial M;\mathbb{R})$ because the boundary condition is of type $C^\infty(\partial M;\mathbb{R})$ (see Ref. [30], p. 252, or Ref. [27], p. 307 for the smoothness of the solution to the Dirichlet problem with smooth boundary condition). Now that $f(\boldsymbol{r}),\boldsymbol{r}\in M\cup\partial M$ is a non-constant scalar field, and $\boldsymbol{F}=\nabla f$ satisfies all the qualifications as a member of $\mathcal{H}^A_2(M\cup\partial M;\mathbb{R})$, we see that the second de Rham cohomology group $\mathcal{H}^2(M\cup\partial M;\mathbb{R})$ will be non-trivial when the boundary ∂M is not connected. $\blacksquare$

Remark Here, we point out several observations related to the proof of this proposition.

(1) The proof of this proposition immediately gives a heuristic physical picture behind the statement in Proposition 2.4. Suppose that the exterior volume $\mathbb{R}^3\setminus(V\cup\partial V)$ is connected, and $\boldsymbol{E}\in C^\infty(V\cup\partial V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$ is a solution to the homogeneous Born equation $(\hat{I}-\chi\hat{\mathcal{G}})\boldsymbol{E}=(\hat{I}+\hat{\mathcal{G}})\boldsymbol{E}=\boldsymbol{0}$ for $\chi=-1$, so that the external field vanishes identically (see Theorem 1.2):

$$\boldsymbol{E}(\boldsymbol{r}):=-\nabla\times\nabla\times\iiint_V\frac{\boldsymbol{E}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\equiv\boldsymbol{0},\quad\forall\boldsymbol{r}\in\mathbb{R}^3\setminus(V\cup\partial V).$$

Recalling the continuity of tangential components in the electric field across a dielectric boundary (see Ref. [9], p. 18), we have the boundary behavior of the internal field as $\boldsymbol{n}\times\boldsymbol{E}(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in\partial V$. Meanwhile, the internal field must also satisfy the differential equation $\nabla^2\boldsymbol{E}(\boldsymbol{r})+(1+\chi)k^2\boldsymbol{E}(\boldsymbol{r})=\nabla^2\boldsymbol{E}(\boldsymbol{r})=\boldsymbol{0},\forall\boldsymbol{r}\in V\cup\partial V$. These conditions altogether qualify $\boldsymbol{E}$ as a member of the Hodge space $\mathcal{H}^A_2(V\cup\partial V;\mathbb{R})=\{\boldsymbol{0}\}$, thus proving that

$$\ker(\hat{I}+\hat{\mathcal{G}})\cap C^\infty(V\cup\partial V;\mathbb{C}^3)\cap\ker(\nabla\cdot)=\{\boldsymbol{0}\}. \tag{2.32}$$

In other words, so long as $b_2(V\cup\partial V):=\dim\mathcal{H}^2(V\cup\partial V;\mathbb{R})=\dim\mathcal{H}^A_2(V\cup\partial V;\mathbb{R})=0$, the vanishing tangential components $\boldsymbol{n}\times\boldsymbol{E}(\boldsymbol{r})\equiv\boldsymbol{0}$ on the boundary surface ∂M for the field $\boldsymbol{E}\in C^\infty(V\cup\partial V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$ leads to an identically vanishing internal field $\boldsymbol{E}(\boldsymbol{r})\equiv\boldsymbol{0},\forall\boldsymbol{r}\in V$. Proposition 2.4 extends the conclusion equation (Eq. 2.32) in this physical picture by replacing the function space $C^\infty(V\cup\partial V;\mathbb{C}^3)$ with a larger one $C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)$.

(2) We note that for a d-dimensional compact manifold $M\cup\partial M$ with boundary ∂M and interior M, we have the following Poincaré duality relation (see Eqs. 9.66 and 9.73 in Ref. [27], Chap. 5):

$$\mathcal{H}^m(M\cup\partial M;\mathbb{R})\cong\mathcal{H}^{d-m}_c(M;\mathbb{R}),m=0,1,\dots,d. \tag{2.33}$$

Here, for the specific case $d=3,m=2$, the Alexander-Spanier cohomology group on the right-hand side of the above formula is defined by $\mathcal{H}^1_c(M;\mathbb{R})=[\ker(\nabla\times)\cap C^\infty_0(M;\mathbb{R}^3)]/\{\nabla f|f\in C^\infty_0(M;\mathbb{R}^3)\}$. If the second Betti number vanishes $\dim\mathcal{H}^2(M\cup\partial M;\mathbb{R})=0$, then we have $\dim\mathcal{H}^1_c(M;\mathbb{R})=0$, which means that every irrotational and compactly supported vector field $\boldsymbol{F}\in\ker(\nabla\times)\cap C^\infty_0(M;\mathbb{R}^3)$ can be represented as the gradient of a scalar field with compact support $\boldsymbol{F}=\nabla f,f\in C^\infty_0(M;\mathbb{R}^3)$. This is indeed the crucial topological tool employed in the rigorous proof of Proposition 2.4.

(3) We recall that the quantity $b_0(\partial M):=\dim\mathcal{H}^0(\partial M;\mathbb{R})$ counts the number of connected components in the compact manifold ∂M. In this proposition, we have provided a detailed proof of the equivalence $b_2(M\cup\partial M)=0\iff b_0(\partial M)=1$ for the compact and connected manifold $M\cup\partial M\subset\mathbb{R}^3$ with smooth boundary ∂M, using techniques similar to those employed in the uniqueness theorem of electrostatics (see Ref. [9], p. 37).

We point out that this result has a direct topological interpretation, which we will sketch below. From the long exact sequence in de Rham cohomology (see Ref. [27], p. 370 and p. 374) for a connected manifold $M\Subset\mathbb{R}^3$ with smooth boundary ∂M, which reads

$$\begin{aligned}&\{\boldsymbol{0}\}\longrightarrow\mathcal{H}^0(M\cup\partial M;\mathbb{R})\longrightarrow\mathcal{H}^0(\partial M;\mathbb{R})\longrightarrow\mathcal{H}^1_c(M;\mathbb{R})\longrightarrow\mathcal{H}^1(M\cup\partial M;\mathbb{R})\longrightarrow\mathcal{H}^1(\partial M;\mathbb{R})\longrightarrow\\&\longrightarrow\mathcal{H}^2_c(M;\mathbb{R})\longrightarrow\mathcal{H}^2(M\cup\partial M;\mathbb{R})\longrightarrow\mathcal{H}^2(\partial M;\mathbb{R})\longrightarrow\mathcal{H}^3_c(M;\mathbb{R})\longrightarrow\{\boldsymbol{0}\},\end{aligned}$$

we know that the alternating sum of the dimensions of the vector spaces appearing in the sequence amounts to zero:

$$\begin{aligned}0&=\dim\mathcal{H}^0(M\cup\partial M;\mathbb{R})-\dim\mathcal{H}^0(\partial M;\mathbb{R})+\dim\mathcal{H}^1_c(M;\mathbb{R})-\dim\mathcal{H}^1(M\cup\partial M;\mathbb{R})+\dim\mathcal{H}^1(\partial M;\mathbb{R})\\&\quad-\dim\mathcal{H}^2_c(M;\mathbb{R})+\dim\mathcal{H}^2(M\cup\partial M;\mathbb{R})-\dim\mathcal{H}^2(\partial M;\mathbb{R})+\dim\mathcal{H}^3_c(M;\mathbb{R})\\&=b_0(M\cup\partial M)-b_0(\partial M)+b_2(M\cup\partial M)-b_1(M\cup\partial M)+b_1(\partial M)\\&\quad-b_1(M\cup\partial M)+b_2(M\cup\partial M)-b_0(\partial M)+b_0(M\cup\partial M).\end{aligned} \tag{2.34}$$

In the above computation, we have exploited the Poincaré duality relation as expressed by Eq. 2.33 as well as the Poincaré duality relation for 2-dimensional compact and oriented manifolds, which leads to $b_0(\partial M)=b_2(\partial M)$.

We may further argue that Eq. 2.34 can be simplified by the relation $b_1(\partial M)=2b_1(M\cup\partial M)$ if the compact set $M\cup\partial M\Subset\mathbb{R}^3$ has a smooth and connected boundary surface $\partial M\Subset\mathbb{R}^3$, with $b_0(\partial M)=1$. The uniformization theorem allows us to topologically classify the surface ∂M according to its first Betti number $b_1(\partial M)=2g$. Here, a surface with $g=0$ is topologically equivalent to a spherical surface, and a positive integer $g>0$ counts the number of holes in a pretzel-shaped surface. Meanwhile, the first Betti number $b_1(M\cup\partial M)$ counts the maximum number of cuts that $M\cup\partial M$ can suffer without being divided into two pieces. By exhausting all the topological classes of ∂M, we can check that $b_1(\partial M)=2g$ if and only if $b_1(M\cup\partial M)=g$. Therefore, if ∂M is a smooth and connected surface, with $b_0(\partial M)=1$, then Eq. 2.34 leads us to

$$b_2(M\cup\partial M)=b_0(\partial M)-b_0(M\cup\partial M), \tag{2.35}$$

the right-hand side of which further simplifies to $b_0(\partial M)-1=0$. In other words, $b_0(\partial M)=1$ implies $b_2(M\cup\partial M)=0$. Conversely, if we know that $b_2(M\cup\partial M)=\dim\mathcal{H}^1_c(M;\mathbb{R})=0$, then the long exact sequence can be truncated as

$$\{\boldsymbol{0}\}\longrightarrow\mathcal{H}^0(M\cup\partial M;\mathbb{R})\longrightarrow\mathcal{H}^0(\partial M;\mathbb{R})\longrightarrow\mathcal{H}^1_c(M;\mathbb{R})=\{\boldsymbol{0}\},$$

which also implies that $b_0(M\cup\partial M)-b_0(\partial M)=0=b_2(M\cup\partial M)$, and $b_0(\partial M)=1$.

Thus, we see that the equivalence relation $b_2(M \cup \partial M) = 0 \iff b_0(\partial M) = 1$ is intrinsic to the topology of the compact manifold $M \cup \partial M \Subset \mathbb{R}^3$ with smooth boundary ∂M.

Finally, we point out that the statement in Eq. 2.35 concerning the second Betti number can be rephrased for a bounded open set $V \Subset \mathbb{R}^3$ as $b_2(V) = \dim \mathcal{H}^2(V;\mathbb{R}) = \dim \mathcal{H}^0(\partial V;\mathbb{R}) - \dim \mathcal{H}^0(V;\mathbb{R}) = \dim \mathcal{H}^0(V \cup \partial V;\mathbb{R}) - 1$ (see Ref. [67], p. 229 for an analytic proof without using the exact sequence). ∎

S2.3.3. Transverse Vector Fields in Star-Shaped Dielectrics

Here, we state and prove an analog of Proposition 2.4 for star-shaped dielectrics. We call a region $M \subset \mathbb{R}^3$ star-shaped, if there exists a point $\boldsymbol{r}_0 \in M$ such that the line segment $L_{\boldsymbol{r}_0,\boldsymbol{r}} := \{(1-\beta)\boldsymbol{r}_0 + \beta\boldsymbol{r} | 0 \le \beta \le 1\}$ joining $\boldsymbol{r}_0$ to any point $\boldsymbol{r} \in M$ also lies in M, i.e. $L_{\boldsymbol{r}_0,\boldsymbol{r}} \subset M, \forall \boldsymbol{r} \in M$.

Proposition 2.8 *If the exterior volume $\mathbb{R}^3 \setminus (V \cup \partial V)$ is connected and the boundary has zero volume $\iint_{\partial V} \mathrm{d}^3\boldsymbol{r} = 0$, the dielectric volume $V = \bigcup_{j=1}^{N_V} V_j$ has connected components $V_1, \dots, V_{N_V}$ where $N_V < +\infty$, and each connected component V_j is star-shaped, then $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F} = \boldsymbol{0}$ holds for any transverse vector field $\boldsymbol{F} \in T(V;\mathbb{C}^3)$.*

Proof Without loss of generality, we may assume that $N_V = 1$ (i.e. V is connected), and $\boldsymbol{r}_0 = \boldsymbol{0} \in V$ (i.e. $L_{\boldsymbol{0},\boldsymbol{r}} \subset V, \forall \boldsymbol{r} \in V$). In this case, we may outline the proof as follows. For any $\boldsymbol{E} \in \ker(\hat{\mathcal{G}} + \hat{I})$, we will construct a family of vector fields $\{\boldsymbol{E}^{(\varepsilon)} \in C_0^\infty(V;\mathbb{C}^3) \cap \ker(\hat{\mathcal{G}} + \hat{I}) | 0 < \varepsilon < 1\}$, which are all irrotational $\nabla \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in V$ and approximate $\boldsymbol{E}$ in the L^2-norm: $\lim_{\varepsilon \to 0^+} \|\boldsymbol{E}^{(\varepsilon)} - \boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)} = 0$. Consequently, we will show that $\langle \boldsymbol{F}, \boldsymbol{E}^{(\varepsilon)} \rangle_V = 0$ for any physically admissible vector field $\boldsymbol{F} \in T(V;\mathbb{C}^3)$. In passage to the limit $\varepsilon \to 0^+$, we may thus confirm that the orthogonality relation $\langle \boldsymbol{F}, \boldsymbol{E} \rangle_V = 0, \forall \boldsymbol{E} \in \ker(\hat{\mathcal{G}} + \hat{I})$ holds for all $\boldsymbol{F} \in T(V;\mathbb{C}^3)$, which entails the claimed relation $\hat{P}_{\ker(\hat{\mathcal{G}}+\hat{I})}\boldsymbol{F} = \boldsymbol{0}$.

The construction of $\boldsymbol{E}^{(\varepsilon)}$ can be accomplished in two steps. Firstly, we consider the spatially rescaled field in the star-shaped region V:

$$\boldsymbol{E}^{[\varepsilon]}(\boldsymbol{r}) := (\hat{S}_\varepsilon \boldsymbol{E})(\boldsymbol{r}) = \begin{cases} \boldsymbol{E}((1-\varepsilon)^{-1}\boldsymbol{r}), & \text{if } (1-\varepsilon)^{-1}\boldsymbol{r} \in V; \\ \boldsymbol{0}, & \text{otherwise.} \end{cases}$$

It is thus clear that such a vector field is compactly supported in V: $\operatorname{supp} \boldsymbol{E}^{[\varepsilon]} \subset \{\boldsymbol{r} \in V | p_V(\boldsymbol{r}) \le 1 - \varepsilon\} \Subset V$, where the Minkowski functional is defined by $p_V(\boldsymbol{r}) = \inf\{\alpha > 0 | \alpha^{-1}\boldsymbol{r} \in V\}$. The Fourier transform of $\boldsymbol{E}^{[\varepsilon]}$ is given by

$$\tilde{\boldsymbol{E}}^{[\varepsilon]}(\boldsymbol{q}) = (1-\varepsilon)^3 \iiint_V \boldsymbol{E}(\boldsymbol{r}) e^{i(1-\varepsilon)\boldsymbol{q}\cdot\boldsymbol{r}} \,\mathrm{d}^3\boldsymbol{r} = (1-\varepsilon)^3 \tilde{\boldsymbol{E}}((1-\varepsilon)\boldsymbol{q}).$$

Therefore, we have $\boldsymbol{q} \times \tilde{\boldsymbol{E}}^{[\varepsilon]}(\boldsymbol{q}) = \boldsymbol{0}, \forall \boldsymbol{q} \in \mathbb{R}^3$, so $\boldsymbol{E}^{[\varepsilon]} \in \ker(\hat{\mathcal{G}} + \hat{I})$. Secondly, we smooth out $\tilde{\boldsymbol{E}}^{[\varepsilon]}$ by a convolution with the mollifier κ_δ (Eq. 2.4) to obtain a vector field $\boldsymbol{E}^{(\varepsilon)} \in C_0^\infty(V;\mathbb{C}^3) \cap \ker(\hat{\mathcal{G}} + \hat{I})$, as $\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) = \iiint_V \boldsymbol{E}^{[\varepsilon]}(\boldsymbol{r}')\varkappa_\delta(\boldsymbol{r} - \boldsymbol{r}') \,\mathrm{d}^3\boldsymbol{r}'$. We may pick $\delta = \delta_\varepsilon \in (0, \frac{1}{2} \min_{\boldsymbol{r} \in \partial V, \boldsymbol{r}' \in \operatorname{supp} \boldsymbol{E}^{[\varepsilon]}} |\boldsymbol{r} - \boldsymbol{r}'|) \subset (0, \frac{1}{2}\varepsilon)$ so as to guarantee the relation $\operatorname{supp} \boldsymbol{E}^{(\varepsilon)} \Subset V$. As the convolution amounts to multiplication by a scalar field in the Fourier transforms, we have $\boldsymbol{q} \times \tilde{\boldsymbol{E}}^{(\varepsilon)}(\boldsymbol{q}) = \boldsymbol{0}, \forall \boldsymbol{q} \in \mathbb{R}^3$, so $\boldsymbol{E}^{(\varepsilon)} \in \ker(\hat{\mathcal{G}} + \hat{I})$, as well.

We now verify that the family of vector fields $\{\boldsymbol{E}^{(\varepsilon)} \in C_0^\infty(V;\mathbb{C}^3) \cap \ker(\hat{\mathcal{G}} + \hat{I}) | 0 < \varepsilon < 1\}$ thus constructed are indeed irrotational and approximate $\boldsymbol{E}$ in the L^2-norm. Here, the vector identity $\nabla \times [\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) e^{i\boldsymbol{q}\cdot\boldsymbol{r}}] = e^{i\boldsymbol{q}\cdot\boldsymbol{r}} \nabla \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) + i\boldsymbol{q} \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) e^{i\boldsymbol{q}\cdot\boldsymbol{r}}$ leads to

$$\boldsymbol{0} = \iiint_V \nabla \times [\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) e^{i\boldsymbol{q}\cdot\boldsymbol{r}}] \,\mathrm{d}^3\boldsymbol{r} = \iiint_V e^{i\boldsymbol{q}\cdot\boldsymbol{r}} \nabla \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) \,\mathrm{d}^3\boldsymbol{r} + i\boldsymbol{q} \times \tilde{\boldsymbol{E}}^{(\varepsilon)}(\boldsymbol{q}) = \iiint_V e^{i\boldsymbol{q}\cdot\boldsymbol{r}} \nabla \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) \,\mathrm{d}^3\boldsymbol{r}, \quad \forall \boldsymbol{q} \in \mathbb{R}^3,$$

which entails irrotationality $\nabla \times \boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in V$. Furthermore, it is clear that

$$\begin{aligned} \lim_{\varepsilon \to 0^+} \|\boldsymbol{E}^{(\varepsilon)} - \boldsymbol{E}^{[\varepsilon]}\|_{L^2(V;\mathbb{C}^3)} &= \lim_{\varepsilon \to 0^+} \frac{1}{(2\pi)^{3/2}} \|\tilde{\boldsymbol{E}}^{[\varepsilon]} \widetilde{\varkappa_{\delta_\varepsilon}} - \tilde{\boldsymbol{E}}^{[\varepsilon]}\|_{L^2(\mathbb{R}^3;\mathbb{C}^3)} \\ &= \lim_{\varepsilon \to 0^+} \frac{(1-\varepsilon)^3}{(2\pi)^{3/2}} \sqrt{\iiint_{\mathbb{R}^3} |\tilde{\boldsymbol{E}}((1-\varepsilon)\boldsymbol{q})|^2 |\widetilde{\varkappa_{\delta_\varepsilon}}(\boldsymbol{q}) - 1|^2 \,\mathrm{d}^3\boldsymbol{q}} \\ &= \lim_{\varepsilon \to 0^+} \left(\frac{1-\varepsilon}{2\pi}\right)^{3/2} \sqrt{\iiint_{\mathbb{R}^3} |\tilde{\boldsymbol{E}}(\boldsymbol{q})|^2 |\widetilde{\varkappa_{\frac{\delta_\varepsilon}{(1-\varepsilon)}}}(\boldsymbol{q}) - 1|^2 \,\mathrm{d}^3\boldsymbol{q}} = 0, \end{aligned}$$

as the convolution kernel $\varkappa_\delta$ is an approximate identity in the limit procedure $\delta \to 0^+$ (Theorem 2.16 in Ref. [29]). The relation $\lim_{\varepsilon\to0^+}\|\boldsymbol{E}-\boldsymbol{E}^{[\varepsilon]}\|_{L^2(V;\mathbb{C}^3)}=\lim_{\varepsilon\to0^+}\|\boldsymbol{E}-\hat{S}_\varepsilon\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$ can be readily verified by the dominated convergence theorem (Theorem 1.8 in Ref. [29]), so long as $\boldsymbol{E}\in C_0^\infty(V;\mathbb{C}^3)$, where $C_0^\infty(V;\mathbb{C}^3)$ is a dense subset of $L^2(V;\mathbb{C}^3)$ (Theorem 2.16 in Ref. [29]). For a generic $\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$, we may pick a family of infinitely differentiable and compactly supported vector fields $\{\boldsymbol{E}^{\{\varepsilon'\}}\in C_0^\infty(V;\mathbb{C}^3)|\varepsilon'>0\}$ such that $\lim_{\varepsilon'\to0^+}\|\boldsymbol{E}^{\{\varepsilon'\}}-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$. With the inequality

$$\|\boldsymbol{E}-\hat{S}_\varepsilon\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\leq\|\boldsymbol{E}^{\{\varepsilon'\}}-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}+\|\boldsymbol{E}^{\{\varepsilon'\}}-\hat{S}_\varepsilon\boldsymbol{E}^{\{\varepsilon'\}}\|_{L^2(V;\mathbb{C}^3)}+\|\hat{S}_\varepsilon\boldsymbol{E}-\hat{S}_\varepsilon\boldsymbol{E}^{\{\varepsilon'\}}\|_{L^2(V;\mathbb{C}^3)}$$

and the fact that $\|\hat{S}_\varepsilon\|_{L^2(V;\mathbb{C}^3)}=(1-\varepsilon)^3$, we arrive at

$$\varlimsup_{\varepsilon\to0^+}\|\boldsymbol{E}-\hat{S}_\varepsilon\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}\leq2\|\boldsymbol{E}^{\{\varepsilon'\}}-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}+\lim_{\varepsilon\to0^+}\|\boldsymbol{E}^{\{\varepsilon'\}}-\hat{S}_\varepsilon\boldsymbol{E}^{\{\varepsilon'\}}\|_{L^2(V;\mathbb{C}^3)}=2\|\boldsymbol{E}^{\{\varepsilon'\}}-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}.$$

Taking the limit $\varepsilon'\to0^+$, we obtain $\varlimsup_{\varepsilon\to0^+}\|\boldsymbol{E}-\hat{S}_\varepsilon\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0,\forall\boldsymbol{E}\in L^2(V;\mathbb{C}^3)$. This eventually confirms that $\lim_{\varepsilon\to0^+}\|\boldsymbol{E}^{(\varepsilon)}-\boldsymbol{E}\|_{L^2(V;\mathbb{C}^3)}=0$.

Now that V is star-shaped, we claim that the irrotational field $\boldsymbol{E}^{(\varepsilon)}$ can be represented as the gradient of a scalar field

$$\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r})=\nabla f(\boldsymbol{r}),\quad\text{where }f(\boldsymbol{r})=\int_0^1\boldsymbol{E}^{(\varepsilon)}(\beta\boldsymbol{r})\cdot\boldsymbol{r}\,\mathrm{d}\,\beta-\int_0^1\boldsymbol{E}^{(\varepsilon)}(\beta\boldsymbol{r}_*)\cdot\boldsymbol{r}_*\,\mathrm{d}\,\beta,\quad\boldsymbol{r}_*\in\partial V.$$

We may also check that $f\in C_0^\infty(V;\mathbb{C})$, in particular, we have the set inclusion relation $\operatorname{supp}f\subset\{\boldsymbol{r}^\sharp\in V|p_V(\boldsymbol{r}^\sharp)\leq\min_{\boldsymbol{r}'\in\operatorname{supp}\boldsymbol{E}^{(\varepsilon)}}p_V(\boldsymbol{r}')\}$. This can be shown by considering a large open ball $O(\mathbf{0},R)\supset V\cup\partial V$, zero-padding $\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r})$ for $\boldsymbol{r}\in O(\mathbf{0},2R)\setminus V$, then converting the representation of $f(\boldsymbol{r})$ for $\boldsymbol{r}\in\{\boldsymbol{r}^\sharp\in V|p_V(\boldsymbol{r}^\sharp)\leq\min_{\boldsymbol{r}'\in\operatorname{supp}\boldsymbol{E}^{(\varepsilon)}}p_V(\boldsymbol{r}')\}$ into a line integral of an irrotational vector field over a loop:

$$\begin{aligned}f(\boldsymbol{r})\;=\;&\left[\int_0^1\boldsymbol{E}^{(\varepsilon)}(\beta\boldsymbol{r})\cdot\boldsymbol{r}\,\mathrm{d}\,\beta+\int_0^1\boldsymbol{E}^{(\varepsilon)}\left((1-\beta)\boldsymbol{r}+\beta R\frac{\boldsymbol{r}}{|\boldsymbol{r}|}\right)\cdot\boldsymbol{r}\,\mathrm{d}\,\beta\right]+\int_{C_{R\boldsymbol{r}/|\boldsymbol{r}|,R\boldsymbol{r}_*/|\boldsymbol{r}_*|}}\boldsymbol{E}^{(\varepsilon)}(\boldsymbol{r}')\cdot\,\mathrm{d}\,\boldsymbol{r}'\\&-\left[\int_0^1\boldsymbol{E}^{(\varepsilon)}(\beta\boldsymbol{r}_*)\cdot\boldsymbol{r}_*\,\mathrm{d}\,\beta+\int_0^1\boldsymbol{E}^{(\varepsilon)}\left((1-\beta)\boldsymbol{r}_*+\beta R\frac{\boldsymbol{r}_*}{|\boldsymbol{r}_*|}\right)\cdot\boldsymbol{r}_*\,\mathrm{d}\,\beta\right]=0.\end{aligned}$$

Here, the arc $C_{R\boldsymbol{r}/|\boldsymbol{r}|,R\boldsymbol{r}_*/|\boldsymbol{r}_*|}\subset\partial O(\mathbf{0},R)$ is a smooth curve that starts from $R\boldsymbol{r}/|\boldsymbol{r}|$ and ends at $R\boldsymbol{r}_*/|\boldsymbol{r}_*|$. Therefore, with $\boldsymbol{F}\in T(V;\mathbb{C}^3):=L^2(V;\mathbb{C}^3)\cap C^\infty(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)$, we may deduce

$$\langle\boldsymbol{F},\boldsymbol{E}^{(\varepsilon)}\rangle_V=\iiint_V\boldsymbol{F}^*(\boldsymbol{r})\cdot\nabla f(\boldsymbol{r})\,\mathrm{d}^3\boldsymbol{r}=\iiint_V\nabla\cdot[\boldsymbol{F}^*(\boldsymbol{r})f(\boldsymbol{r})]\,\mathrm{d}^3\boldsymbol{r}=0,\quad0<\varepsilon<1\tag{2.36}$$

by quoting the divergence-free condition $\nabla\cdot\boldsymbol{F}(\boldsymbol{r})=0,\forall\boldsymbol{r}\in V$, and using the Gauss theorem in vector calculus to convert a volume integral into a surface integral with vanishing integrand. As the scalar field $f\in C_0^\infty(V;\mathbb{C})$ is compactly supported in V, the Gauss theorem is indeed applicable, irrespective of the smoothness of the boundary ∂V. Therefore, Eq. 2.36 leads us to $\langle\boldsymbol{F},\boldsymbol{E}\rangle_V=\lim_{\varepsilon\to0^+}\langle\boldsymbol{F},\boldsymbol{E}^{(\varepsilon)}\rangle_V=0,\forall\boldsymbol{F}\in T(V;\mathbb{C}^3)$. ▮

Remark A star-shaped open set $V\subset\mathbb{R}^d$ ($d\geq1$) has vanishing Betti numbers $b_m(V)=\dim\mathcal{H}^m(V;\mathbb{R})=0$ for all positive integers m, according to the Poincaré lemma (see Ref. [87], pp. 33-35, also Ref. [88], p. 23). In particular, as seen in the proof of Proposition 2.8 for a star-shaped, bounded, and open set $V\subset\mathbb{R}^3$, it is possible to represent a compactly supported irrotational field $\boldsymbol{E}^{(\varepsilon)}\in C_0^\infty(V;\mathbb{R}^3)\cap\ker(\nabla\times)$ as the gradient of a scalar field with compact support. This property is necessarily guaranteed by the Poincaré duality relation for open manifolds: $\dim\mathcal{H}_c^1(V;\mathbb{R})=\dim\mathcal{H}^2(V;\mathbb{R})=0$ (see Ref. [87], p. 46, also Ref. [88], p. 127). In other words, we have seen that a unifying topological theme in Propositions 2.4 and 2.8 is the trivialization of cohomology group $\mathcal{H}^2\cong\mathcal{H}_c^1$.

Convex polyhedra are among the many examples of star-shaped region without a smooth boundary. Hence Proposition 2.8 gives an extension of Propositions 2.4 to a certain class of non-smooth dielectrics that might be of practical interest. ▮

S2.3.4. Surface Geometry and Optical Resonance

Here, we supply some detailed derivations in differential geometry that is instrumental in the establishment of the optical resonance theorem (Theorem 2.1) for dielectric media with smooth boundary surfaces.

In the following three lemmata, $\Sigma = \partial M$ will stand for the smooth boundary surface of a bounded open set $M \Subset \mathbb{R}^3$, with local curvilinear coordinates (ξ, η), normal vector $\boldsymbol{n}$, surface element $\mathrm{d}\,\tilde{S}$, mean curvature H, Gaussian curvature K. The coordinate argument $\boldsymbol{r}$ of a scalar or vector field will be suppressed for the sake of brevity, so long as no confusion arises. Smooth vector fields written in Greek letters ($\boldsymbol{\alpha}, \boldsymbol{\beta}$ etc.) are tangent to the surface Σ (i.e. with vanishing normal component $\boldsymbol{n} \cdot \boldsymbol{\alpha} = 0, \boldsymbol{n} \cdot \boldsymbol{\beta} = 0$ etc.), whereas smooth vector fields written in Latin letters ($\boldsymbol{a}, \boldsymbol{b}$ etc.) may or may not be tangent to the surface Σ. The Weingarten transform to an arbitrary smooth vector field $\boldsymbol{f}$ is defined as $\hat{W}(\boldsymbol{f}) = -(\boldsymbol{f} \cdot \nabla)\boldsymbol{n}$. Here, whenever we talk about a "smooth field", we mean that it is infinitely differentiable in a certain tubular neighborhood of the smooth surface Σ.

Lemma 2.4 *Let $\boldsymbol{n}(\xi, \eta, \varepsilon)$ be a smooth vector field defined in a tubular neighborhood of Σ via the procedure in Proposition 2.6, then the following identities hold:*[34]

$$(\boldsymbol{n} \cdot \nabla)\boldsymbol{n} = \boldsymbol{0}, \quad \boldsymbol{n} \cdot [(\boldsymbol{a} \cdot \nabla)\boldsymbol{n}] = 0, \quad \nabla \times \boldsymbol{n} = \boldsymbol{0}, \quad \nabla \cdot \boldsymbol{n} = -2H, \tag{2.37}$$

for any smooth vector field $\boldsymbol{a}$ and smooth scalar field u. For any two smooth vector fields $\boldsymbol{a}$ and $\boldsymbol{b}$, we have

$$\boldsymbol{a} \cdot \{\boldsymbol{n} \times [(\boldsymbol{b} \cdot \nabla)\boldsymbol{n}]\} + \boldsymbol{a} \cdot \{[(\boldsymbol{n} \times \boldsymbol{b}) \cdot \nabla]\boldsymbol{n}\} = 2H\boldsymbol{n} \cdot (\boldsymbol{a} \times \boldsymbol{b}). \tag{2.38}$$

Proof From Eq. 2.29, we know that the normal vector field $\boldsymbol{n}(\xi, \eta, \varepsilon) = \boldsymbol{n}(\xi, \eta, 0)$ does not explicitly depend on the variable ε, hence

$$(\boldsymbol{n} \cdot \nabla)\boldsymbol{n} = -\frac{\partial \boldsymbol{n}(\xi, \eta, \varepsilon)}{\partial \varepsilon} = \boldsymbol{0}.$$

Now that $\partial \boldsymbol{n}/\partial \varepsilon = \boldsymbol{0}$, and the differential operator $\boldsymbol{a} \cdot \nabla$ is a linear combination of $\partial/\partial\xi, \partial/\partial\eta$ and $\partial/\partial\varepsilon$, we know that $(\boldsymbol{a} \cdot \nabla)\boldsymbol{n}$ is a linear combination of $\partial \boldsymbol{n}/\partial\xi$ and $\partial \boldsymbol{n}/\partial\eta$, which in turn, can be expressed as a linear combination of tangent vectors $\partial \tilde{\boldsymbol{r}}/\partial\xi, \partial \tilde{\boldsymbol{r}}/\partial\eta$, according to the Weingarten transformation (Eq. 2.28). This shows that the vector $(\boldsymbol{a} \cdot \nabla)\boldsymbol{n}$ is orthogonal to $\boldsymbol{n}$, i.e. $\boldsymbol{n} \cdot [(\boldsymbol{a} \cdot \nabla)\boldsymbol{n}] = 0$.

Expansion of the vanishing gradient $\nabla(\boldsymbol{n} \cdot \boldsymbol{n}) = \boldsymbol{0}$ results in

$$\boldsymbol{0} = \nabla(\boldsymbol{n} \cdot \boldsymbol{n}) = 2(\boldsymbol{n} \cdot \nabla)\boldsymbol{n} + 2\boldsymbol{n} \times (\nabla \times \boldsymbol{n}) = 2\boldsymbol{n} \times (\nabla \times \boldsymbol{n}),$$

hence the tangential components of $\nabla \times \boldsymbol{n}$ must vanish. Meanwhile, we may use the Stokes theorem to perform the following computations for sufficiently small $\delta > 0$:

$$\frac{1}{\pi\delta^2} \iint_{\{\boldsymbol{r}(\xi,\eta)|(\xi-\xi_\star)^2+(\eta-\eta_\star)^2<\delta^2\}} \boldsymbol{n} \cdot \nabla \times \boldsymbol{n} \,\mathrm{d}\,\tilde{S} = \frac{1}{\pi\delta^2} \oint_{\{\boldsymbol{r}(\xi,\eta)|(\xi-\xi_\star)^2+(\eta-\eta_\star)^2=\delta^2\}} \boldsymbol{n} \cdot \mathrm{d}\,\tilde{\boldsymbol{l}} = 0,$$

where the line element $\mathrm{d}\,\tilde{\boldsymbol{l}}$ lies in the tangent plane. In the limit of $\delta \to 0^+$, we may check that $\boldsymbol{n} \cdot \nabla \times \boldsymbol{n}$ must vanish identically. Therefore, the vector $\nabla \times \boldsymbol{n}$ has vanishing components in both the tangential and normal directions.

To demonstrate the relation $\nabla \cdot \boldsymbol{n} = -2H$ for $\boldsymbol{r} \in \Sigma$, we apply the Gauss theorem of vector calculus to a tubular neighborhood $\tilde{\boldsymbol{r}}(U_{\star,\delta} \times (-\varepsilon', \varepsilon')) := \{\tilde{\boldsymbol{r}}(\xi, \eta, \varepsilon)|(\xi - \xi_\star)^2 + (\eta - \eta_\star)^2 < \delta^2, |\varepsilon| < \varepsilon'\}$ as follows:

$$\begin{aligned}
\nabla \cdot \boldsymbol{n}(\xi_\star, \eta_\star, 0) &= \lim_{\substack{\delta\to 0^+\\ \varepsilon'\to 0^+}} \frac{\iiint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times(-\varepsilon',\varepsilon'))} \nabla \cdot \boldsymbol{n} \,\mathrm{d}^3\tilde{\boldsymbol{r}}}{\iiint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times(-\varepsilon',\varepsilon'))} \mathrm{d}^3\tilde{\boldsymbol{r}}} = \lim_{\substack{\delta\to 0^+\\ \varepsilon'\to 0^+}} \frac{\iint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times\{-\varepsilon'\})} \mathrm{d}\,\tilde{S} - \iint_{\tilde{\boldsymbol{r}}(U_{\star}\times\{\varepsilon'\})} \mathrm{d}\,\tilde{S}}{\iiint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times(-\varepsilon',\varepsilon'))} \mathrm{d}^3\tilde{\boldsymbol{r}}} \\
&= \lim_{\substack{\delta\to 0^+\\ \varepsilon'\to 0^+}} \frac{\iint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times\{0\})} [-2\varepsilon' H - 2\varepsilon' H + O(\varepsilon'^2)] \,\mathrm{d}\,\tilde{S}}{2\varepsilon' \iint_{\tilde{\boldsymbol{r}}(U_{\star,\delta}\times\{0\})} \mathrm{d}\,\tilde{S}} = -2H(\xi_\star, \eta_\star, 0).
\end{aligned}$$

This concludes the proof of all the identities listed in Eq. 2.37.

As both sides of Eq. 2.38 are independent of the choice of local coordinates, we may verify the identity by choosing a special curvilinear coordinate system for the sake of convenience. In particular, we may locally parameterize the smooth surface Σ by the principal curvature coordinates (ξ, η) that diagonalize the Weingarten transformation $\hat{W}$ (see Eq. 2.28), with the tangent vectors $\partial \boldsymbol{r}/\partial\xi = \boldsymbol{e}_1, \partial \boldsymbol{r}/\partial\eta = \boldsymbol{e}_2$ being both unit vectors at a point of interest $\boldsymbol{r}_\star \in \Sigma$, satisfying $\boldsymbol{e}_1\cdot\boldsymbol{e}_1=\boldsymbol{e}_2\cdot\boldsymbol{e}_2=1,\boldsymbol{e}_1\cdot\boldsymbol{e}_2=0,\boldsymbol{e}_1\times\boldsymbol{e}_2=\boldsymbol{n}:=\boldsymbol{e}_3$, as well as $\partial\boldsymbol{n}/\partial\xi=-\kappa_1\boldsymbol{e}_1,\partial\boldsymbol{n}/\partial\eta=-\kappa_2\boldsymbol{e}_2$. Here, κ_1 and κ_2 represent the two principal curvatures (i.e. eigenvalues of the Weingarten transformation $\hat{W}$). At the point $\boldsymbol{r}_\star$, we may consider the decompositions $\boldsymbol{a}=a_1\boldsymbol{e}_1+a_2\boldsymbol{e}_2+a_3\boldsymbol{e}_3$ and $\boldsymbol{b}=b_1\boldsymbol{e}_1+b_2\boldsymbol{e}_2+b_3\boldsymbol{e}_3$. Direct computation shows that $\boldsymbol{n}\times[(\boldsymbol{b}\cdot\nabla)\boldsymbol{n}]=\boldsymbol{e}_3\times[-\kappa_1b_1\boldsymbol{e}_1-\kappa_2b_2\boldsymbol{e}_2]=-\kappa_1b_1\boldsymbol{e}_2+\kappa_2b_2\boldsymbol{e}_1$, so $\boldsymbol{a}\cdot\{\boldsymbol{n}\times[(\boldsymbol{b}\cdot\nabla)\boldsymbol{n}]\}=-\kappa_1b_1a_2+\kappa_2b_2a_1$. Similarly, we have $\boldsymbol{a}\cdot\{[(\boldsymbol{n}\times\boldsymbol{b})\cdot\nabla]\boldsymbol{n}\}=-\kappa_2b_1a_2+\kappa_1b_2a_1$. As $2H=\operatorname{Tr}(\hat{W})=\kappa_1+\kappa_2$ by definition, the truthfulness of Eq. 2.38 is now evident. ∎

---

[34] In some literature, the mean curvature H may be defined with a sign convention opposite to the one presented in S2.3.1, thus one might obtain the identity $\nabla \cdot \boldsymbol{n} = +2H$ instead.

Remark Using the extended definition of the Weingarten transform $\hat{W}(\boldsymbol{f})=-(\boldsymbol{f}\cdot\nabla)\boldsymbol{n}$, we may write Eq. 2.38 as $\boldsymbol{a}\cdot(\boldsymbol{n}\times\hat{W}(\boldsymbol{b}))+\boldsymbol{a}\cdot[\hat{W}(\boldsymbol{n}\times\boldsymbol{b})]=2H\boldsymbol{a}\cdot(\boldsymbol{n}\times\boldsymbol{b})$. Furthermore, as the vector field $\boldsymbol{a}$ is arbitrary in Eq. 2.38, we obtain the following result:

$$\boldsymbol{n}\times\hat{W}(\boldsymbol{b})+\hat{W}(\boldsymbol{n}\times\boldsymbol{b})=2H\boldsymbol{n}\times\boldsymbol{b}. \tag{2.39}$$

It is also easy to recognize that $\boldsymbol{a}\cdot\hat{W}(\boldsymbol{b})=\boldsymbol{b}\cdot\hat{W}(\boldsymbol{a})$ holds for any two smooth vector fields $\boldsymbol{a}$ and $\boldsymbol{b}$. ∎

Hereafter, we will denote the tangential projection of a vector field $\boldsymbol{a}$ as $\boldsymbol{a}_{\boldsymbol{t}}=(\boldsymbol{n}\times\boldsymbol{a})\times\boldsymbol{n}$, and name $\boldsymbol{a}_{\boldsymbol{n}}=(\boldsymbol{n}\cdot\boldsymbol{a})\boldsymbol{n}$ as the normal projection.

Lemma 2.5 *Let $\boldsymbol{\alpha}$ be a smooth tangential field and $\boldsymbol{f}$ be a smooth vector field, then we have the identities*

$$\begin{aligned}\nabla\times\boldsymbol{\alpha} &= \hat{W}(\boldsymbol{\alpha})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)(\boldsymbol{\alpha}\times\boldsymbol{n})-\boldsymbol{n}[\nabla\cdot(\boldsymbol{n}\times\boldsymbol{\alpha})], &&(2.40)\\ \nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\boldsymbol{f})] &= -\hat{W}((\boldsymbol{n}\cdot\nabla)\boldsymbol{f})-\boldsymbol{n}\{\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]\}\\ &\quad -2\nabla H\times\boldsymbol{f}_{\boldsymbol{t}}-2H\{\hat{W}(\boldsymbol{f})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)(\boldsymbol{f}\times\boldsymbol{n})+\boldsymbol{n}[\boldsymbol{n}\cdot(\nabla\times\boldsymbol{f})]\} &&(2.41)\end{aligned}$$

Proof On one hand, from the identity

$$\boldsymbol{n}\times(\nabla\times\boldsymbol{\alpha})=\nabla(\boldsymbol{n}\cdot\boldsymbol{\alpha})-(\boldsymbol{\alpha}\cdot\nabla)\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)\boldsymbol{\alpha}-\boldsymbol{\alpha}\times(\nabla\times\boldsymbol{n})=\hat{W}(\boldsymbol{\alpha})-(\boldsymbol{n}\cdot\nabla)\boldsymbol{\alpha},$$

we may deduce

$$(\nabla\times\boldsymbol{\alpha})_{\boldsymbol{t}}=\hat{W}(\boldsymbol{\alpha})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)(\boldsymbol{\alpha}\times\boldsymbol{n}). \tag{2.42}$$

On the other hand, the vector identity $\nabla\cdot(\boldsymbol{n}\times\boldsymbol{\alpha})=\boldsymbol{\alpha}\cdot(\nabla\times\boldsymbol{n})-\boldsymbol{n}\cdot(\nabla\times\boldsymbol{\alpha})=-\boldsymbol{n}\cdot(\nabla\times\boldsymbol{\alpha})$ reveals that $(\nabla\times\boldsymbol{\alpha})_{\boldsymbol{n}}=-\boldsymbol{n}[\nabla\cdot(\boldsymbol{n}\times\boldsymbol{\alpha})]$. Combining the tangential and normal projections of $\nabla\times\boldsymbol{\alpha}$, we arrive at Eq. 2.40.

Using Eq. 2.39, we may split $\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\boldsymbol{f})]$ into two parts as $\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\boldsymbol{f})]=\nabla\times\hat{W}(\boldsymbol{f}_{\boldsymbol{t}})-2\nabla\times(H\boldsymbol{f}_{\boldsymbol{t}})$. Exploiting Eq. 2.40, we may evaluate

$$\nabla\times\hat{W}(\boldsymbol{f}_{\boldsymbol{t}}) \;=\; \hat{W}^2(\boldsymbol{f})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)[\hat{W}(\boldsymbol{f})\times\boldsymbol{n}]-\boldsymbol{n}\{\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]\}$$

using the principal curvature coordinates (ξ,η) as in the proof of Eq. 2.38. We start the simplification of $\nabla\times\hat{W}(\boldsymbol{f}_{\boldsymbol{t}})$ by writing $\boldsymbol{f}_{\boldsymbol{t}}=f_1\boldsymbol{e}_1+f_2\boldsymbol{e}_2$ and computing

$$\hat{W}^2(\boldsymbol{f})+\hat{W}((\boldsymbol{n}\cdot\nabla)\boldsymbol{f})-(\boldsymbol{n}\cdot\nabla)\hat{W}(\boldsymbol{f})=\kappa_1^2f_1\boldsymbol{e}_1+\kappa_2^2f_2\boldsymbol{e}_2-[(\boldsymbol{n}\cdot\nabla)\kappa_1]f_1\boldsymbol{e}_1-[(\boldsymbol{n}\cdot\nabla)\kappa_2]f_2\boldsymbol{e}_2.$$

We note that once the local curvilinear coordinates (ξ,η) depict a principal curvature coordinate system on Σ, they will remain so on the parallel surfaces of Σ. As a consequence, the values of principal curvatures at the corresponding points will be obey the relation

$$\kappa_1(\boldsymbol{r}-\varepsilon\boldsymbol{n})=\frac{\kappa_1(\boldsymbol{r})}{1+\varepsilon\kappa_1(\boldsymbol{r})},\quad \kappa_2(\boldsymbol{r}-\varepsilon\boldsymbol{n})=\frac{\kappa_2(\boldsymbol{r})}{1+\varepsilon\kappa_2(\boldsymbol{r})},$$

so we have $(\boldsymbol{n}\cdot\nabla)\kappa_1=\kappa_1^2,(\boldsymbol{n}\cdot\nabla)\kappa_2=\kappa_2^2$, and consequently $\hat{W}^2(\boldsymbol{f})+\hat{W}((\boldsymbol{n}\cdot\nabla)\boldsymbol{f})-(\boldsymbol{n}\cdot\nabla)\hat{W}(\boldsymbol{f})=\boldsymbol{0}$,

$$\nabla\times\hat{W}(\boldsymbol{f}_{\boldsymbol{t}})=-\hat{W}((\boldsymbol{n}\cdot\nabla)\boldsymbol{f})-\boldsymbol{n}\{\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]\}.$$

Furthermore, we may quote Eq. 2.40 to derive

$$\begin{aligned}-2\nabla\times(H\boldsymbol{f}_{\boldsymbol{t}}) &= -2\nabla H\times\boldsymbol{f}_{\boldsymbol{t}}-2H\nabla\times\boldsymbol{f}_{\boldsymbol{t}}\\ &= -2\nabla H\times\boldsymbol{f}_{\boldsymbol{t}}-2H\{\hat{W}(\boldsymbol{f}_{\boldsymbol{t}})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)(\boldsymbol{f}_{\boldsymbol{t}}\times\boldsymbol{n})-\boldsymbol{n}[\nabla\cdot(\boldsymbol{n}\times\boldsymbol{f})]\}\\ &= -2\nabla H\times\boldsymbol{f}_{\boldsymbol{t}}-2H\{\hat{W}(\boldsymbol{f})\times\boldsymbol{n}-(\boldsymbol{n}\cdot\nabla)(\boldsymbol{f}\times\boldsymbol{n})+\boldsymbol{n}[\boldsymbol{n}\cdot(\nabla\times\boldsymbol{f})]\},\end{aligned}$$

which eventually lead to Eq. 2.41. ∎

Lemma 2.6 *For a smooth vector field $\boldsymbol{f}$, we have the inequality*

$$\begin{aligned}|\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\boldsymbol{f})]| \;\le\;& \left(\sqrt{4H^2-2K}+2|H|\right)|(\boldsymbol{n}\cdot\nabla)\boldsymbol{f}|+2|\nabla H||\boldsymbol{f}|+2|H|\left(\sqrt{4H^2-2K}|\boldsymbol{f}|+2|\nabla\times\boldsymbol{f}|\right)\\ &+2\sqrt{4H^2-2K}\,[|\nabla(\boldsymbol{e}_x\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_y\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_z\cdot\boldsymbol{f})|]\end{aligned}\tag{2.43}$$

Proof Using the inequality $|\hat{W}(\boldsymbol{a})|^2\le\max(\kappa_1^2,\kappa_2^2)|\boldsymbol{a}|^2\le(\kappa_1^2+\kappa_2^2)|\boldsymbol{a}|^2=(4H^2-2K)|\boldsymbol{a}|^2$, it is easy to deduce from Eq. 2.41 the following estimate:

$$\begin{aligned}|\nabla\times[\boldsymbol{n}\times\hat{W}(\boldsymbol{n}\times\boldsymbol{f})]| \;\le\;& \sqrt{4H^2-2K}|(\boldsymbol{n}\cdot\nabla)\boldsymbol{f}|+|\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]|\\ &+2|\nabla H||\boldsymbol{f}|+2|H|\left(\sqrt{4H^2-2K}|\boldsymbol{f}|+|(\boldsymbol{n}\cdot\nabla)\boldsymbol{f}|+|\nabla\times\boldsymbol{f}|\right).\end{aligned}$$

Thus, we only need to verify

$$|\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]|\le2\sqrt{4H^2-2K}\,[|\nabla(\boldsymbol{e}_x\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_y\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_z\cdot\boldsymbol{f})|]+2|H\nabla\times\boldsymbol{f}|\tag{2.44}$$

in order to arrive at Eq. 2.43. Using the vector identities

$$\begin{aligned}\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})] &= -\boldsymbol{n}\cdot\nabla\times\hat{W}(\boldsymbol{f}),\\ \nabla(\boldsymbol{f}\cdot\boldsymbol{n}) &= (\boldsymbol{n}\cdot\nabla)\boldsymbol{f}-\hat{W}(\boldsymbol{f})+\boldsymbol{n}\times(\nabla\times\boldsymbol{f}),\\ \boldsymbol{n}\cdot\nabla\times[\boldsymbol{n}\times(\nabla\times\boldsymbol{f})] &= 2H\boldsymbol{n}\cdot\nabla\times\boldsymbol{f}-\boldsymbol{n}\cdot[(\boldsymbol{n}\cdot\nabla)\nabla\times\boldsymbol{f}],\end{aligned}$$

that are direct consequences of Eq. 2.37, we may bound the left-hand side of Eq. 2.44 as

$$|\nabla\cdot[\boldsymbol{n}\times\hat{W}(\boldsymbol{f})]|\le2|H\nabla\times\boldsymbol{f}|+|\nabla\times(\boldsymbol{n}\cdot\nabla)\boldsymbol{f}-(\boldsymbol{n}\cdot\nabla)\nabla\times\boldsymbol{f}|\tag{2.45}$$

Using the Cartesian decomposition $\boldsymbol{f}=(\boldsymbol{e}_x\cdot\boldsymbol{f})\boldsymbol{e}_x+(\boldsymbol{e}_y\cdot\boldsymbol{f})\boldsymbol{e}_y+(\boldsymbol{e}_z\cdot\boldsymbol{f})\boldsymbol{e}_z$, we may expand the last term on the right-hand side of Eq. 2.45. A typical term of such an expansion looks like

$$\boldsymbol{e}_z\left[\frac{\partial}{\partial x}(\boldsymbol{n}\cdot\nabla)(\boldsymbol{e}_y\cdot\boldsymbol{f})-(\boldsymbol{n}\cdot\nabla)\frac{\partial}{\partial x}(\boldsymbol{e}_y\cdot\boldsymbol{f})\right]=\boldsymbol{e}_z\left(\frac{\partial\boldsymbol{n}}{\partial x}\cdot\nabla\right)(\boldsymbol{e}_y\cdot\boldsymbol{f})=-[\hat{W}(\boldsymbol{e}_x)\cdot\nabla(\boldsymbol{e}_y\cdot\boldsymbol{f})]\boldsymbol{e}_z,$$

which is bounded by $\sqrt{4H^2-2K}\,|\nabla(\boldsymbol{e}_y\cdot\boldsymbol{f})|$. There are six such terms in the Cartesian decomposition of the expression $|\nabla\times(\boldsymbol{n}\cdot\nabla)\boldsymbol{f}-(\boldsymbol{n}\cdot\nabla)\nabla\times\boldsymbol{f}|$, eventually leading to an upper bound as $2\sqrt{4H^2-2K}\,[|\nabla(\boldsymbol{e}_x\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_y\cdot\boldsymbol{f})|+|\nabla(\boldsymbol{e}_z\cdot\boldsymbol{f})|]$. ∎

In what follows, the dielectric volume $V=\bigcup_{j=1}^{N_V}V_j\Subset\mathbb{R}^3$ will be a bounded open set with finitely many connected components $V_1,\dots,V_{N_V}$, each of which has a smooth boundary surface $\partial V_j, j=1,\dots,N_V$. The dielectric boundary $\partial V=\bigsqcup_{j=1}^{N_V}\partial V_j$ consists of mutually disjoint subsets $\partial V_1,\dots,\partial V_{N_V}$, satisfying

$$\delta_0:=\min_{\substack{\boldsymbol{r}\in\partial V_j,\boldsymbol{r}'\in\partial V_{j'}\\ 1\le j<j'\le N_V}}|\boldsymbol{r}-\boldsymbol{r}'|>0.$$

The "pinched volume" is defined by $V^-_{(\varepsilon)}:=V\setminus\{\tilde{\boldsymbol{r}}=\boldsymbol{r}^\sharp-\varepsilon'\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp\in\partial V,\varepsilon'\in(0,\varepsilon]\}$ while the "swollen volume" is defined by $V^+_{(\delta)}:=V\cup\{\tilde{\boldsymbol{r}}=\boldsymbol{r}^\sharp+\delta'\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp\in\partial V,\delta'\in[0,\delta)\}$. The parallel surface $\partial V^-_{(\varepsilon)}:=\{\tilde{\boldsymbol{r}}=\boldsymbol{r}^\sharp-\varepsilon\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp\in\partial V\}$ is assumed to be smooth for $\varepsilon\in(0,2\varepsilon_V]$, while the parallel surface $\partial V^+_{(\delta)}:=\{\tilde{\boldsymbol{r}}=\boldsymbol{r}^\sharp+\delta\boldsymbol{n}^\sharp(\boldsymbol{r}^\sharp)|\boldsymbol{r}^\sharp\in\partial V\}$ is assumed to be smooth for $\delta\in(0,\delta_V]$ where $\delta_V<\delta_0/2$.

Lemma 2.7 *There is a smooth scalar field $\varsigma\in C^\infty(\mathbb{R}^3;\mathbb{R})$ satisfying*

$$\varsigma(\boldsymbol{r})=1,\forall\boldsymbol{r}\in V\setminus V^-_{(3\varepsilon_V/2)},\qquad\varsigma(\boldsymbol{r})=0,\forall\boldsymbol{r}\in(\mathbb{R}^3\setminus V^+_{(\delta_V)})\cup V^-_{(2\varepsilon_V)},\tag{2.46}$$

and a Lipschitz estimate

$$|\varsigma(\boldsymbol{r})-\varsigma(\boldsymbol{r}')|\le C_1|\boldsymbol{r}-\boldsymbol{r}'|,\quad\forall\boldsymbol{r},\boldsymbol{r}'\in\mathbb{R}^3,\tag{2.47}$$

for some finite constant $C_1<+\infty$. Meanwhile, there is a finite constant $C_2<+\infty$ that fulfills a Lipschitz estimate

$$|D^{\boldsymbol{\mu}}(\boldsymbol{n}'-\boldsymbol{n})|:=|D^{\boldsymbol{\mu}}\boldsymbol{n}(\boldsymbol{r}')-D^{\boldsymbol{\mu}}\boldsymbol{n}(\boldsymbol{r})|\le C_2|\boldsymbol{r}-\boldsymbol{r}'|,\quad|\boldsymbol{\mu}|\le2\tag{2.48}$$

whenever $\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')\ne0$. For any given $\delta_V<\delta_0/2$, the finite constants C_1 and C_2 are solely determined by the geometry of V and the choice of the parameter ε_V.

Proof Picking a positive constant $\delta > 0$ satisfying $3\delta/2 < \delta_V - \delta$, $3(\varepsilon_V + \delta)/2 < 2\varepsilon_V - \delta$, we may first construct a scalar field

$$\varsigma^\sharp(\boldsymbol r) = \begin{cases} 1, & \boldsymbol r \in V^+_{(3\delta/2)} \setminus V^-_{(3(\varepsilon_V+\delta)/2)} \\ 0, & \text{otherwise} \end{cases}$$

and then obtain $\varsigma \in C^\infty(\mathbb R^3;\mathbb R)$ as a convolution $\varsigma(\boldsymbol r) = \iiint_{\mathbb R^3} \varsigma^\sharp(\boldsymbol r')\varkappa_\delta(\boldsymbol r - \boldsymbol r')\,\mathrm d^3\boldsymbol r'$. Here, the convolution kernel $\varkappa_\delta$ is given by Eq. 2.4. Using the properties of parallel surfaces, one can directly verify that the scalar field $\varsigma \in C^\infty(\mathbb R^3;\mathbb R)$ thus constructed satisfies the requirements in Eq. 2.46. Now that the gradient field $\nabla\varsigma \in C^0(\operatorname{Cl}(\mathbb R^3 \setminus V^+_{(\delta_V)});\mathbb R^3) \cap C^\infty_0(\mathbb R^3 \setminus V^+_{(\delta_V)};\mathbb R^3)$ is continuous on a compact set $\operatorname{Cl}(\mathbb R^3 \setminus V^+_{(\delta_V)})$, and vanishes outside $\operatorname{Cl}(\mathbb R^3 \setminus V^+_{(\delta_V)})$, it must be bounded in the entire space. In other words, there is a finite constant $C_1 < +\infty$ such that $|\nabla\varsigma(\boldsymbol r)| \le C_1, \forall \boldsymbol r \in \mathbb R^3$. From the mean-value theorem in calculus, it immediately follows that $|\varsigma(\boldsymbol r) - \varsigma(\boldsymbol r')| \le C_1|\boldsymbol r - \boldsymbol r'|, \forall \boldsymbol r, \boldsymbol r' \in \mathbb R^3$, as claimed in Eq. 2.47.

Noting that $\boldsymbol n(\boldsymbol r) := -\nabla \operatorname{dist}(\boldsymbol r, \partial V^+_{(\delta_V)})/|\nabla \operatorname{dist}(\boldsymbol r, \partial V^+_{(\delta_V)})|$ defines a smooth vector field $\boldsymbol n \in C^\infty(\operatorname{Cl}(V^+_{(5\delta/2)} \setminus V^-_{((3\varepsilon_V+5\delta)/2)});\mathbb R^3)$ in the compact set $\operatorname{Cl}(V^+_{(5\delta/2)} \setminus V^-_{((3\varepsilon_V+5\delta)/2)})$ with smooth boundary $\partial V^+_{(5\delta/2)} \cup \partial V^-_{((3\varepsilon_V+5\delta)/2)}$, we have the Lipschitz estimate

$$|D^{\boldsymbol\mu}(\boldsymbol n' - \boldsymbol n)| := |D^{\boldsymbol\mu}\boldsymbol n(\boldsymbol r') - D^{\boldsymbol\mu}\boldsymbol n(\boldsymbol r)| \le C_2|\boldsymbol r - \boldsymbol r'|, \quad |\boldsymbol\mu| \le 2, \forall \boldsymbol r, \boldsymbol r' \in \operatorname{Cl}(V^+_{(5\delta/2)} \setminus V^-_{((3\varepsilon_V+5\delta)/2)})$$

according to the Sobolev imbeddings into Lipschitz spaces (see Ref. [66], p. 99). By our construction of $\varsigma \in C^\infty(\mathbb R^3;\mathbb R)$, we have $\varsigma(\boldsymbol r)\varsigma(\boldsymbol r') = 0$ if either $\boldsymbol r$ or $\boldsymbol r'$ falls outside $\operatorname{Cl}(V^+_{(5\delta/2)} \setminus V^-_{((3\varepsilon_V+5\delta)/2)})$. Therefore, we have confirmed the Lipschitz estimate in Eq. 2.48 for all cases where $\varsigma(\boldsymbol r)\varsigma(\boldsymbol r') \ne 0$. ▮

Lemma 2.8 *For $\varepsilon \in (0, \varepsilon_V]$, define a scalar field*

$$g_{(\varepsilon)}(\boldsymbol r) := \oiint_{\partial V^-_{(\varepsilon)}} \frac{(\boldsymbol n' - \boldsymbol n)\cdot \boldsymbol p(\boldsymbol r')}{4\pi|\boldsymbol r - \boldsymbol r'|}\,\mathrm d\tilde S'$$

for a smooth vector field $\boldsymbol p \in C^\infty(V;\mathbb C^3) \cap L^2(V;\mathbb C^3) \cap \ker(\nabla\cdot) \cap \ker(\nabla\times) \subset \mathfrak h^2(V;\mathbb C^3)$ that is square-integrable, divergence-free and irrotational. There is a finite constant $g_{\varepsilon_V;V} < +\infty$ such that

$$\|g_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb C)} \le g_{\varepsilon_V;V}\|\boldsymbol p\|_{L^2(V;\mathbb C^3)}$$

is satisfied by all $\varepsilon \in (0, \varepsilon_V]$.

Proof We decompose $\boldsymbol p = \boldsymbol p_{\boldsymbol t} + \boldsymbol p_{\boldsymbol n}$ into tangential and normal projections $\boldsymbol p_{\boldsymbol t}(\boldsymbol r') = [\boldsymbol n' \times \boldsymbol p(\boldsymbol r')] \times \boldsymbol n', \boldsymbol p_{\boldsymbol n}(\boldsymbol r') = [\boldsymbol n' \cdot \boldsymbol p(\boldsymbol r')]\boldsymbol n'$, and apply the Gauss theorem in vector calculus to convert surface integrals into volume integrals in the following computations:

$$\begin{aligned} g_{(\varepsilon)}(\boldsymbol r) &= \oiint_{\partial V^-_{(\varepsilon)}} \frac{[\boldsymbol n' \cdot \boldsymbol p(\boldsymbol r')][(\boldsymbol n' - \boldsymbol n)\cdot \boldsymbol n']}{4\pi|\boldsymbol r - \boldsymbol r'|}\,\mathrm d\tilde S' + \oiint_{\partial V^-_{(\varepsilon)}} \frac{\boldsymbol n' \cdot \{\boldsymbol p(\boldsymbol r') \times [\boldsymbol n' \times (\boldsymbol n' - \boldsymbol n)]\}}{4\pi|\boldsymbol r - \boldsymbol r'|}\,\mathrm d\tilde S' \\ &= \iiint_{V^-_{(\varepsilon)}\setminus V^-_{(3\varepsilon_V/2)}} \boldsymbol p(\boldsymbol r')\cdot\nabla'\left[\frac{(\boldsymbol n' - \boldsymbol n)\cdot \boldsymbol n'}{4\pi|\boldsymbol r - \boldsymbol r'|}\right]\mathrm d^3\boldsymbol r' + \iiint_{V^-_{(\varepsilon)}\setminus V^-_{(3\varepsilon_V/2)}} \boldsymbol p(\boldsymbol r')\cdot\left\{\nabla' \times \left[\frac{(\boldsymbol n' - \boldsymbol n)\times \boldsymbol n'}{4\pi|\boldsymbol r - \boldsymbol r'|}\right]\right\}\mathrm d^3\boldsymbol r' \\ &\quad + \oiint_{\partial V^-_{(3\varepsilon_V/2)}} \frac{(\boldsymbol n' - \boldsymbol n)\cdot \boldsymbol p(\boldsymbol r')}{4\pi|\boldsymbol r - \boldsymbol r'|}\,\mathrm d\tilde S' \\ &= \iiint_{V^-_{(\varepsilon)}\setminus V^-_{(3\varepsilon_V/2)}} \boldsymbol p(\boldsymbol r')\cdot \boldsymbol g(\boldsymbol r, \boldsymbol r')\,\mathrm d^3\boldsymbol r' + \oiint_{\partial V^-_{(3\varepsilon_V/2)}} \frac{(\boldsymbol n' - \boldsymbol n)\cdot \boldsymbol p(\boldsymbol r')}{4\pi|\boldsymbol r - \boldsymbol r'|}\,\mathrm d\tilde S'. \end{aligned} \tag{2.49}$$

In the derivation of Eq. 2.49, we have already exploited the conditions that $\nabla' \cdot \boldsymbol p(\boldsymbol r') = 0, \nabla' \times \boldsymbol p(\boldsymbol r') = \boldsymbol 0, \forall \boldsymbol r' \in V$.

We may abbreviate Eq. 2.49 as $g_{(\varepsilon)} = g^\sharp_{(\varepsilon)} + g^{\sharp\sharp}_{(\varepsilon)}$, where $g^\sharp_{(\varepsilon)}$ is the contribution from the volume integral in $V^-_{(\varepsilon)} \setminus V^-_{(3\varepsilon_V/2)}$ and $g^{\sharp\sharp}_{(\varepsilon)}$ is the contribution from the surface integral on $\partial V^-_{(3\varepsilon_V/2)}$. We will briefly derive two bound estimates $\|g^\sharp_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb C)} \le g^{\sharp,0}_{\varepsilon_V;V}\|\boldsymbol p\|_{L^2(V;\mathbb C^3)}$ and $\|g^{\sharp\sharp}_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb C)} \le g^{\sharp\sharp}_{\varepsilon_V;V}\|\boldsymbol p\|_{L^2(V;\mathbb C^3)}$, before probing into the upper bounds of $\|g^\sharp_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb C)}$.

For $\boldsymbol{r} \in V \setminus V^-_{(\varepsilon_V)}$ and $\boldsymbol{r}' \in V \setminus V^-_{(3\varepsilon_V/2)}$, we have $\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}') = 1$ according to Eq. 2.46. Consequently, the Lipschitz estimate in Eq. 2.48 reveals that the integral kernel $\boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')$ satisfies

$$|\boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')| < \frac{g_1}{|\boldsymbol{r} - \boldsymbol{r}'|}, \quad \boldsymbol{r} \in V \setminus V^-_{(\varepsilon_V)}, \boldsymbol{r}' \in V \setminus V^-_{(3\varepsilon_V/2)}$$

for some constant $g_1 < +\infty$. Thus, the square integrability condition

$$g^{\sharp,0}_{\varepsilon_V;V} := \sqrt{\iiint_{V \setminus V^-_{(\varepsilon_V)}} \left(\iiint_{V \setminus V^-_{(3\varepsilon_V/2)}} |\boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')|^2 \,\mathrm{d}^3 \boldsymbol{r}' \right) \mathrm{d}^3 \boldsymbol{r}} < +\infty$$

leads to an estimate $\|g^\sharp_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le g^{\sharp,0}_{\varepsilon_V;V} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$ for a finite constant $g^{\sharp,0}_{\varepsilon_V;V}$ applicable to all $\varepsilon \in (0, \varepsilon_V]$.

Using the Bergman space inequality (Eq. 2.19), we can argue that there is a finite constant $g^\sharp_{\varepsilon_V;V} < +\infty$ satisfying

$$\|g^{\sharp\sharp}_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le \sqrt{\frac{4\pi}{3} \left(\frac{3\varepsilon_V}{2} \right)^3} g^{\sharp\sharp}_{\varepsilon_V;V} \max_{\boldsymbol{r}' \in \partial V^-_{(3\varepsilon_V/2)}} |\boldsymbol{p}(\boldsymbol{r}')| \le g^{\sharp\sharp}_{\varepsilon_V;V} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}.$$

Here, with the Lipschitz estimate in Eq. 2.48, we may set a finite upper bound for $g^{\sharp\sharp}_{\varepsilon_V;V}$ using the bounded surface area $\oiint_{\partial V^-_{(3\varepsilon_V/2)}} \mathrm{d}\,\tilde{S}' < +\infty$, as well as the maximum values of

$$\left| \frac{(\boldsymbol{n}' - \boldsymbol{n})}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \right|, \quad \left| (\boldsymbol{u} \cdot \nabla) \frac{(\boldsymbol{n}' - \boldsymbol{n})}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \right|, \quad \left| (\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \frac{(\boldsymbol{n}' - \boldsymbol{n})}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \right|, \qquad \boldsymbol{u}, \boldsymbol{v} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$$

for the points $\boldsymbol{r} \in V \setminus V^-_{(\varepsilon_V)}, \boldsymbol{r}' \in \partial V^-_{(3\varepsilon_V/2)}$ satisfying $\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}') = 1$.

Now, we will use the Calderón-Zygmund theory in harmonic analysis to derive an estimate

$$\sum_{|\mu|=2} \|D^{\boldsymbol{\mu}} g^\sharp_{(\varepsilon)}\|^2_{L^2(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le (g^{\sharp,2}_{\varepsilon_V;V} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)})^2 \quad \text{with} \quad g^{\sharp,2}_{\varepsilon_V;V} < +\infty.$$

First, for the smooth vector field $\boldsymbol{p} \in C^\infty(V;\mathbb{C}^3)$, we may differentiate under the integral sign as

$$\begin{aligned}
&(\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \iiint_{V^-_{(\varepsilon)} \setminus V^-_{(3\varepsilon_V/2)}} \boldsymbol{p}(\boldsymbol{r}') \cdot \boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}') \,\mathrm{d}^3 \boldsymbol{r}' \\
&= \boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r}) \cdot \boldsymbol{p}(\boldsymbol{r}) + \lim_{\varepsilon' \to 0^+} \iiint_{V^-_{(\varepsilon)} \setminus [V^-_{(3\varepsilon_V/2)} \cup O(\boldsymbol{r}, \varepsilon')]} \boldsymbol{p}(\boldsymbol{r}') \cdot [(\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')] \,\mathrm{d}^3 \boldsymbol{r}', \quad \forall \boldsymbol{r} \in V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)},
\end{aligned}$$

for $\boldsymbol{u}, \boldsymbol{v} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$. Here, the vector $\boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r})$ is determined by the local shape of the parallel surface at $\boldsymbol{r}$, via the relation

$$\boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r}) := \lim_{\varepsilon' \to 0^+} \varepsilon'^2 \oiint_{\boldsymbol{r}' \in \partial O(\boldsymbol{r}, \varepsilon')} \left\{ \frac{\boldsymbol{r}' - \boldsymbol{r}}{|\boldsymbol{r} - \boldsymbol{r}'|} \left[(\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \frac{(\boldsymbol{n}' - \boldsymbol{n}) \cdot \boldsymbol{n}'}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \right] + \frac{\boldsymbol{r}' - \boldsymbol{r}}{|\boldsymbol{r} - \boldsymbol{r}'|} \times \left[(\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \frac{(\boldsymbol{n}' - \boldsymbol{n}) \times \boldsymbol{n}'}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \right] \right\} \mathrm{d}\,\Omega'$$

where $\mathrm{d}\,\Omega'$ is the infinitesimal solid angle on the spherical surface $\partial O(\boldsymbol{r}, \varepsilon')$. With the Lipschitz estimate in Eq. 2.48, we can see that $|\boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r})|$ is bounded for $\boldsymbol{r} \in V \setminus V^-_{(\varepsilon_V)}$, hence we have $\sum_{\boldsymbol{u},\boldsymbol{v} \in \{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}} \|\boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}} \cdot \boldsymbol{p}\|^2_{L^2(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le C_3 \|\boldsymbol{p}\|^2_{L^2(V;\mathbb{C}^3)}, \forall \varepsilon \in (0, \varepsilon_V]$, for some finite constant $C_3 < +\infty$ associated with the geometry of V. Then, it is clear that the L^2-estimate for $D^{\boldsymbol{\mu}} g^\sharp_{(\varepsilon)}, |\boldsymbol{\mu}| = 2$ can be reduced to the study of the linear transformations $\hat{g}_{\boldsymbol{u},\boldsymbol{v}}$ for $\boldsymbol{u}, \boldsymbol{v} \in \{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$, where

$$(\hat{g}_{\boldsymbol{u},\boldsymbol{v}} \boldsymbol{p})(\boldsymbol{r}) := \lim_{\varepsilon' \to 0^+} \iiint_{V^-_{(\varepsilon)} \setminus [V^-_{(3\varepsilon_V/2)} \cup O(\boldsymbol{r}, \varepsilon')]} \boldsymbol{p}(\boldsymbol{r}') \cdot [(\boldsymbol{u} \cdot \nabla)(\boldsymbol{v} \cdot \nabla) \boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')] \,\mathrm{d}^3 \boldsymbol{r}'.$$

We will establish the boundedness of the linear operator $\hat{g}_{\boldsymbol{u},\boldsymbol{v}} : L^2(V \setminus V^-_{(3\varepsilon_V/2)};\mathbb{C}^3) \longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ in two steps: (i) Verify that the second-order derivatives of $\boldsymbol{g}(\boldsymbol{r}, \boldsymbol{r}')$ induces a Calderón-Zygmund kernel; (ii) Verify that a modified version of the cancellation condition is satisfied by the Calderón-Zygmund kernel in question.

(i) We claim that every Cartesian component of

$$\boldsymbol{g}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r},\boldsymbol{r}') := \varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\boldsymbol{g}(\boldsymbol{r},\boldsymbol{r}')$$

defines a Calderón-Zygmund kernel satisfying the three criteria outlined in Eq. 1.27, where the smooth scalar field $\varsigma \in C^\infty(\mathbb{R}^3;\mathbb{R})$ satisfies Eq. 2.46. To qualify $\boldsymbol{g}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r},\boldsymbol{r}')$ as a Calderón-Zygmund kernel, it would suffice to decompose the gradient and rotation into Cartesian components and investigate the behavior of third-order partial derivatives

$$\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)(\boldsymbol{w}\cdot\nabla')\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|} \quad\text{and}\quad \varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)(\boldsymbol{w}\cdot\nabla')\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}, \qquad \forall \boldsymbol{u},\boldsymbol{v},\boldsymbol{w}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}.$$

It is then evident that all the inequalities in Eq. 1.27 hold by virtue of Eqs. 2.47 and 2.48.

(ii) Following Ref. [49], pp. 305-306, we define two integrals

$$I_{\varepsilon,N}(\boldsymbol{r}) := \iiint_{\varepsilon<|\boldsymbol{r}-\boldsymbol{r}'|<N} K(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r}' \quad\text{and}\quad J_{\varepsilon,N}(\boldsymbol{r}') := \iiint_{\varepsilon<|\boldsymbol{r}-\boldsymbol{r}'|<N} K(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^3\boldsymbol{r},$$

where the kernel $K(\boldsymbol{r},\boldsymbol{r}')$ stands for any Cartesian component of $\boldsymbol{g}_{\boldsymbol{u},\boldsymbol{v}}(\boldsymbol{r},\boldsymbol{r}')$. We then claim that there is a finite constant $A<+\infty$ fulfilling a modified version of the cancellation condition (see Ref. [49], p. 306 and p. 324) for all $0<\varepsilon<N<+\infty$:

$$|I_{\varepsilon,N}(\boldsymbol{r})|\leq A, \quad \forall\boldsymbol{r}\in\mathbb{R}^3; \qquad |J_{\varepsilon,N}(\boldsymbol{r}')|\leq A, \quad \forall\boldsymbol{r}'\in\mathbb{R}^3.$$

Once this "cancellation condition" can be confirmed for the integral kernel $\boldsymbol{g}_{\boldsymbol{u},\boldsymbol{v}}$, we may reveal the induced linear mapping $\hat{g}_{\boldsymbol{u},\boldsymbol{v}}: L^2(V\setminus V^-_{(3\varepsilon_V/2)};\mathbb{C}^3)\longrightarrow L^2(\mathbb{R}^3;\mathbb{C}^3)$ as a bounded operator.

Momentarily assuming that $\boldsymbol{r}\in V\setminus V^-_{(3\varepsilon_V/2)}, N<\operatorname{dist}(\boldsymbol{r},\partial V^+_{(\delta_V)}\cup\partial V^-_{(2\varepsilon_V)})$, we may evaluate the contribution to $I_{\varepsilon,N}(\boldsymbol{r})$ by converting it to a surface integral with the help of the Gauss theorem in the following fashion:

$$\begin{aligned}
&\iiint_{O(\boldsymbol{r},N)\setminus O(\boldsymbol{r},\varepsilon)}\left\{\nabla'\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]+\nabla'\times\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\right\}\mathrm{d}^3\boldsymbol{r}'\\
={}& N^2\oiint_{\partial O(\boldsymbol{r},N)}\left\{\frac{\boldsymbol{r}'-\boldsymbol{r}}{|\boldsymbol{r}-\boldsymbol{r}'|}\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]+\frac{\boldsymbol{r}'-\boldsymbol{r}}{|\boldsymbol{r}-\boldsymbol{r}'|}\times\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\right\}\mathrm{d}\,\Omega'\\
&-\varepsilon^2\oiint_{\partial O(\boldsymbol{r},\varepsilon)}\left\{\frac{\boldsymbol{r}'-\boldsymbol{r}}{|\boldsymbol{r}-\boldsymbol{r}'|}\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]+\frac{\boldsymbol{r}'-\boldsymbol{r}}{|\boldsymbol{r}-\boldsymbol{r}'|}\times\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\right\}\mathrm{d}\,\Omega'
\end{aligned}\tag{2.50}$$

where $\mathrm{d}\,\Omega'$ represents the infinitesimal solid angle on the spherical boundary surfaces $\partial O(\boldsymbol{r},N)$ and $\partial O(\boldsymbol{r},\varepsilon)$. Quoting the Lipschitz estimate in Eq. 2.48, we can verify that Eq. 2.50 is uniformly bounded as $\varepsilon\to0^+$. This shows the boundedness of $|I_{\varepsilon,N}(\boldsymbol{r})|$ under the constraint that $\boldsymbol{r}\in V\setminus V^-_{(3\varepsilon_V/2)}, N<\operatorname{dist}(\boldsymbol{r},\partial V^+_{(\delta_V)}\cup\partial V^-_{(2\varepsilon_V)})$. Now if we relax this constraint and bring the smooth factors $\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')$ into picture, we can see that $|I_{\varepsilon,N}(\boldsymbol{r})|$ is also uniformly bounded for any given $\boldsymbol{r}\in\mathbb{R}^3$ in passage to the limit $N\to+\infty$.

The bound estimate of $|J_{\varepsilon,N}(\boldsymbol{r})|$ can be built upon a similar strategy. We may take notice of the fact that once integrated upon the volume $O(\boldsymbol{r}',N)\setminus O(\boldsymbol{r}',\varepsilon)$, both terms

$$\begin{aligned}
(\nabla'+\nabla)\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right] &= O\left(\frac{1}{|\boldsymbol{r}-\boldsymbol{r}'|^2}\right)\\
(\nabla'+\nabla)\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right] &= O\left(\frac{1}{|\boldsymbol{r}-\boldsymbol{r}'|^2}\right)
\end{aligned}$$

will yield uniformly bounded results that hold for $\boldsymbol{r}'\in V\setminus V^-_{(3\varepsilon_V/2)}, N<\operatorname{dist}(\boldsymbol{r}',\partial V^+_{(\delta_V)}\cup\partial V^-_{(2\varepsilon_V)})$. In the meantime, the volume integral

$$\iiint_{O(\boldsymbol{r}',N)\setminus O(\boldsymbol{r}',\varepsilon)}\left\{\nabla\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\cdot\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]+\nabla\times\left[(\boldsymbol{u}\cdot\nabla)(\boldsymbol{v}\cdot\nabla)\frac{(\boldsymbol{n}'-\boldsymbol{n})\times\boldsymbol{n}'}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right]\right\}\mathrm{d}^3\boldsymbol{r}$$

may be recast into surface integrals on $\partial O(\boldsymbol{r}',N)$ and $\partial O(\boldsymbol{r}',\varepsilon)$, in the same vein as Eq. 2.50. Eventually, when the smooth factors $\varsigma(\boldsymbol{r})\varsigma(\boldsymbol{r}')$ come into play for large values of N, the boundedness of $|J_{\varepsilon,N}(\boldsymbol{r})|$ also becomes evident.

Combining the conclusions in (i) and (ii) with the former result $\sum_{\boldsymbol{u},\boldsymbol{v}\in\{\boldsymbol{e}_x,\boldsymbol{e}_y,\boldsymbol{e}_z\}}\|\boldsymbol{s}_{\boldsymbol{u},\boldsymbol{v}}\cdot\boldsymbol{p}\|^2_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})}\leq C_3\|\boldsymbol{p}\|^2_{L^2(V;\mathbb{C}^3)}$, we have $\sum_{|\boldsymbol{\mu}|=2}\|D^{\boldsymbol{\mu}}g^\sharp_{(\varepsilon)}\|^2_{L^2(\mathbb{R}^3;\mathbb{C})}\leq g^{\sharp,2}_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(3\varepsilon_V/2)};\mathbb{C}^3)}\leq g^{\sharp,2}_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$ for some finite constant $g^{\sharp,2}_{\varepsilon_V;V} < +\infty$. Using the Gagliardo-Nirenberg-Sobolev inequality, we may derive a similar bound for $\|\nabla g^{\sharp}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C}^3)} \le g^{\sharp,1}_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$.

Recalling the formerly deduced inequality $\|g^{\sharp}_{(\varepsilon)}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le g^{\sharp,0}_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$ and $\|g^{\sharp\sharp}_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le g^{\sharp\sharp}_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$, we arrive at the bound estimate

$$\|g_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le \left[\sqrt{(g^{\sharp,0}_{\varepsilon_V;V})^2 + \frac{(g^{\sharp,1}_{\varepsilon_V;V})^2}{k^2} + \frac{(g^{\sharp,2}_{\varepsilon_V;V})^2}{k^4}} + g^{\sharp\sharp}_{\varepsilon_V;V}\right]\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}, \quad \forall\varepsilon\in(0,\varepsilon_V]$$

as claimed. ▮

Corollary 2.5 *Consider two smooth vector fields* $\boldsymbol{U},\boldsymbol{p}\in C^\infty(V;\mathbb{C}^3)\cap L^2(V;\mathbb{C}^3)\cap\ker(\nabla\cdot)\cap\ker(\nabla\times)\subset\mathfrak{b}^2(V;\mathbb{C}^3)$ *that are square-integrable, divergence-free and irrotational, and a scalar field* $g_{(\varepsilon)}$ *as defined in Lemma 2.8. Define a functional of* $\varepsilon;\boldsymbol{U},\boldsymbol{p}$ *as*

$$\psi^\sharp[\varepsilon;\boldsymbol{U},\boldsymbol{p}] = \oiint_{\partial V^-_{(\varepsilon+0)}} 2H(\boldsymbol{r})[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})]g_{(\varepsilon)}(\boldsymbol{r})\,\mathrm{d}\tilde{S} - \oiint_{\partial V^-_{(\varepsilon+0)}} \boldsymbol{U}(\boldsymbol{r})\cdot\nabla g_{(\varepsilon)}(\boldsymbol{r})\,\mathrm{d}\tilde{S},$$

then we have a finite constant $\Psi^\sharp_{\varepsilon_V;V} < +\infty$ *fulfilling the inequality*

$$|\psi^\sharp[\varepsilon;\boldsymbol{U},\boldsymbol{p}]| \le \Psi^\sharp_{\varepsilon_V;V}\|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}, \quad \forall\varepsilon\in(0,\varepsilon_V].$$

Proof We may use the Gauss theorem to derive

$$\begin{aligned}\psi^\sharp[\varepsilon;\boldsymbol{U},\boldsymbol{p}] &= \iiint_{V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r})\cdot\nabla[2H(\boldsymbol{r})g_{(\varepsilon)}(\boldsymbol{r})]\,\mathrm{d}^3\boldsymbol{r}\\ &\quad-\iiint_{V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r})\cdot\{\nabla[\boldsymbol{n}\cdot\nabla g_{(\varepsilon)}(\boldsymbol{r})]-\nabla\times[\boldsymbol{n}\times\nabla g_{(\varepsilon)}(\boldsymbol{r})]\}\,\mathrm{d}^3\boldsymbol{r}\\ &\quad+\oiint_{\partial V^-_{(\varepsilon_V)}} 2H(\boldsymbol{r})[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})]g_{(\varepsilon)}(\boldsymbol{r})\,\mathrm{d}\tilde{S} - \oiint_{\partial V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r})\cdot\nabla g_{(\varepsilon)}(\boldsymbol{r})\,\mathrm{d}\tilde{S}. \end{aligned} \tag{2.51}$$

After applying the Cauchy-Schwarz inequality to the first two lines on the right-hand side of Eq. 2.51, we can immediately realize that these volume integrals will be bounded by some constant multiple of $\|\boldsymbol{U}\|_{L^2(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)},\mathbb{C}^3)}\|g_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le \|\boldsymbol{U}\|_{L^2(V,\mathbb{C}^3)}\|g_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})}$, which in turn, must be bounded by $g_{\varepsilon_V;V}\|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$, according to Lemma 2.8.

Next, we note that the Bergman space estimate (Eq. 2.19) will yield two inequalities

$$\oiint_{\partial V^-_{(\varepsilon_V)}} |2H(\boldsymbol{r})[\boldsymbol{n}\cdot\boldsymbol{U}(\boldsymbol{r})]|^2\,\mathrm{d}\tilde{S} \le C_4\|\boldsymbol{U}\|^2_{L^2(V,\mathbb{C}^3)}, \quad \oiint_{\partial V^-_{(\varepsilon_V)}} |\boldsymbol{U}(\boldsymbol{r})|^2\,\mathrm{d}\tilde{S} \le C_5\|\boldsymbol{U}\|^2_{L^2(V,\mathbb{C}^3)},$$

where the finite constants $C_4, C_5 < +\infty$ are of purely geometrical origin. Meanwhile, quoting the boundary trace imbedding theorem (see Ref. [66], p. 164) for the Sobolev spaces $W^{m,p}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C}^{d'})\subset L^{p'}(\partial V^-_{(\varepsilon)}\cup\partial V^-_{(\varepsilon_V)};\mathbb{C}^{d'})$ with $p\le p'\le(d-1)p/(d-mp), d=3, d'=1,3$, we obtain two inequalities

$$\begin{aligned}\oiint_{\partial V^-_{(\varepsilon_V)}} |g_{(\varepsilon)}(\boldsymbol{r})|^2\,\mathrm{d}\tilde{S} &\le C_6\|g_{(\varepsilon)}\|^2_{W^{1,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C})} \le C_6(g_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)})^2,\\ \oiint_{\partial V^-_{(\varepsilon_V)}} |\nabla g_{(\varepsilon)}(\boldsymbol{r})|^2\,\mathrm{d}\tilde{S} &\le C_6\|\nabla g_{(\varepsilon)}\|^2_{W^{1,2}(V^-_{(\varepsilon)}\setminus V^-_{(\varepsilon_V)};\mathbb{C}^3)} \le C_6(g_{\varepsilon_V;V}\|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)})^2,\end{aligned}$$

where the constant $C_6 < +\infty$ can be chosen as uniformly bounded for all $\varepsilon\in(0,\varepsilon_V]$. Therefore, the bound estimates for the surface integrals in the last line of Eq. 2.51 can also be accomplished with the Cauchy-Schwarz inequality, and the finite constants C_4, C_5, C_6. ▮

Corollary 2.6 *Consider two smooth vector fields* $\boldsymbol{U}, \boldsymbol{p} \in C^\infty(V; \mathbb{C}^3) \cap L^2(V; \mathbb{C}^3) \cap \ker(\nabla\cdot) \cap \ker(\nabla\times) \subset \mathfrak{b}^2(V; \mathbb{C}^3)$*. For* $\varepsilon \in (0, \varepsilon_V]$*, define a vector field*

$$\boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}) := \oiint_{\partial V^-_{(\varepsilon)}} \frac{(\boldsymbol{n}' - \boldsymbol{n}) \times \boldsymbol{p}(\boldsymbol{r}')}{4\pi|\boldsymbol{r} - \boldsymbol{r}'|} \,\mathrm{d}\,\tilde{S}'.$$

There is a finite constant $G_{\varepsilon_V;V} < +\infty$ *such that*

$$\|\boldsymbol{G}_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)}; \mathbb{C}^3)} \le G_{\varepsilon_V;V} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$$

is satisfied by all $\varepsilon \in (0, \varepsilon_V]$*. Furthermore, the functional*

$$\psi^{\sharp\sharp}[\varepsilon; \boldsymbol{U}, \boldsymbol{p}] = \oiint_{\partial V^-_{(\varepsilon+0)}} 2H(\boldsymbol{r})[\boldsymbol{n} \times \boldsymbol{U}(\boldsymbol{r})] \cdot \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}) \,\mathrm{d}\,\tilde{S} + \oiint_{\partial V^-_{(\varepsilon+0)}} \boldsymbol{U}(\boldsymbol{r}) \cdot [2\hat{W}(\boldsymbol{n} \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})) - \nabla \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})] \,\mathrm{d}\,\tilde{S}$$

fulfills a bound estimate

$$|\psi^{\sharp\sharp}[\varepsilon; \boldsymbol{U}, \boldsymbol{p}]| \le \Psi^{\sharp\sharp}_{\varepsilon_V;V} \|\boldsymbol{U}\|_{L^2(V;\mathbb{C}^3)} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}, \quad \forall \varepsilon \in (0, \varepsilon_V].$$

Proof After projecting the vector field $\boldsymbol{G}_{(\varepsilon)}$ to every unit vector $\boldsymbol{u}$ in the Cartesian basis set $\{\boldsymbol{e}_x, \boldsymbol{e}_y, \boldsymbol{e}_z\}$, we can establish the inequality $\|\boldsymbol{G}_{(\varepsilon)}\|_{W^{2,2}(V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)}; \mathbb{C}^3)} \le G_{\varepsilon_V;V} \|\boldsymbol{p}\|_{L^2(V;\mathbb{C}^3)}$ following essentially the same line of thoughts as in Lemma 2.8. The adaptation of the proof only amounts to a replacement of all the occurrence of $(\boldsymbol{n}' - \boldsymbol{n})$ with $\boldsymbol{u} \times (\boldsymbol{n}' - \boldsymbol{n})$, while all the other reasonings in Lemma 2.8 remain valid.

Applying the Gauss theorem to $\psi^{\sharp\sharp}[\varepsilon; \boldsymbol{U}, \boldsymbol{p}]$, we obtain

$$\begin{aligned} \psi^{\sharp\sharp}[\varepsilon; \boldsymbol{U}, \boldsymbol{p}] &= -\iiint_{V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r}) \cdot \nabla \times [2H(\boldsymbol{r})\boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})] \,\mathrm{d}^3\boldsymbol{r} - \iiint_{V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r}) \cdot \nabla \times [\boldsymbol{n} \times \hat{W}(\boldsymbol{n} \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}))] \,\mathrm{d}^3\boldsymbol{r} \\ &\quad - \iiint_{V^-_{(\varepsilon)} \setminus V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r}) \cdot \{\nabla[\boldsymbol{n} \cdot \nabla \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})] - \nabla \times [\boldsymbol{n} \times \nabla \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})]\} \,\mathrm{d}^3\boldsymbol{r} \\ &\quad + \oiint_{\partial V^-_{(\varepsilon_V)}} 2H(\boldsymbol{r})[\boldsymbol{n} \times \boldsymbol{U}(\boldsymbol{r})] \cdot \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}) \,\mathrm{d}\,\tilde{S} + \oiint_{\partial V^-_{(\varepsilon_V)}} \boldsymbol{U}(\boldsymbol{r}) \cdot [2\hat{W}(\boldsymbol{n} \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})) - \nabla \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})] \,\mathrm{d}\,\tilde{S}. \end{aligned} \tag{2.52}$$

Recalling the bound estimate of $\nabla \times [\boldsymbol{n} \times \hat{W}(\boldsymbol{n} \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}))]$ from Eq. 2.43 as well as the geometric relation $|\hat{W}(\boldsymbol{n} \times \boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r}))|^2 \le [4H^2(\boldsymbol{r}) - 2K(\boldsymbol{r})]|\boldsymbol{G}_{(\varepsilon)}(\boldsymbol{r})|^2$, we can bound all the volume integrals and the surface integrals in Eq. 2.52 using the methodology that has worked for Eq. 2.51. ∎

So far, using the geometric information of the dielectric volume V, we have established bound estimates for all the terms in Eq. 2.26, which are crucial for the proof of the "optical resonance theorem" (Theorem 2.1).

S2.3.5. Optical Resonance in Mie Scattering

Here, we verify the implications of the "optical resonance theorem" (Theorem 2.1) in the specific case of light scattering on spherical dielectric (Mie scattering). With the help of spherical harmonic functions $Y_{\ell m}$, spherical Bessel functions j_ℓ and spherical Hankel functions $h^{(2)}_\ell$, we will give an independent and conceptually simpler derivation of the "optical resonance theorem" in Mie scattering.

Proposition 2.9 (Optical Resonance in Mie Scattering) *If the dielectric volume* $V = O(\boldsymbol{0}, R)$*,* $0 < R < +\infty$ *is spherical, then* $\hat{\mathcal{G}}(\hat{I} + 2\hat{\mathcal{G}}) : \Phi(O(\boldsymbol{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\boldsymbol{0}, R); \mathbb{C}^3)$ *is a compact operator, and we have* $\sigma^\Phi(\hat{\mathcal{G}}) \setminus \sigma^\Phi_p(\hat{\mathcal{G}}) = \{0, -1/2\}$*. Meanwhile, the spectrum* $\sigma^\Phi(\hat{\mathcal{G}})$ *is a countable set, with* $\{0, -1/2\}$ *being the only possible accumulation points. In other words, we can enumerate* $\sigma^\Phi(\hat{\mathcal{G}}) = \{\lambda_s | s = 1, 2, \ldots\}$ *as a sequence, and there may exist non-trivial subsequences*[35] *in* $\sigma^\Phi(\hat{\mathcal{G}})$ *that converge to either* 0 *or* $-1/2$*, but not any other complex numbers as their limit.*

---

[35] We call a subsequence non-trivial if all its members are distinct.

Proof We will give a constructive proof that the operator $(\hat{\mathcal{G}}-\hat{\gamma})(\hat{I}+2(\hat{\mathcal{G}}-\hat{\gamma})):\Phi(O(\mathbf{0},R);\mathbb{C}^3)\longrightarrow\Phi(O(\mathbf{0},R);\mathbb{C}^3)$ is compact, which entails the compactness of $\hat{\mathcal{G}}(\hat{I}+2\hat{\mathcal{G}}):\Phi(O(\mathbf{0},R);\mathbb{C}^3)\longrightarrow\Phi(O(\mathbf{0},R);\mathbb{C}^3)$ as well as most of the other claims in the proposition.

We begin the constructive proof by recalling the eigenequation $(\hat{\gamma}-\hat{\mathcal{G}})\boldsymbol{F}_{\ell m}=(\hat{I}-\hat{\mathcal{D}})\boldsymbol{F}_{\ell m}=\ell\boldsymbol{F}_{\ell m}/(2\ell+1)$ for $\boldsymbol{F}_{\ell m}(|\boldsymbol{r}|,\theta,\phi)=\nabla[|\boldsymbol{r}|^\ell Y_{\ell m}(\theta,\phi)],\ell\in\mathbb{Z}\cap[1,+\infty);m\in\mathbb{Z}\cap[-\ell,\ell]$, as pointed out in the Remark to Proposition 1.3. With the normalized eigenvector $\boldsymbol{f}_{\ell m}=\boldsymbol{F}_{\ell m}/\|\boldsymbol{F}_{\ell m}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}$, we will construct a spectral decomposition for the Hermitian operator $\hat{\gamma}-\hat{\mathcal{G}}=\hat{I}-\hat{\mathcal{D}}:\Phi(O(\mathbf{0},R);\mathbb{C}^3)\longrightarrow\Phi(O(\mathbf{0},R);\mathbb{C}^3)$ as

$$(\hat{I}-\hat{\mathcal{D}})\boldsymbol{F}=\sum_{\ell=1}^{\infty}\left[\sum_{m=-\ell}^{\ell}\frac{\ell\langle\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}\boldsymbol{f}_{\ell m}}{2\ell+1}\right],\quad\forall\boldsymbol{F}\in\Phi(O(\mathbf{0},R);\mathbb{C}^3).\tag{2.53}$$

Here, the right-hand side of Eq. 2.53 is interpreted as an infinite series in ℓ that converges in the L^2-norm of $\Phi(O(\mathbf{0},R);\mathbb{C}^3)$.

We embark on the verification of Eq. 2.53 by picking a Hermitian operator $\hat{P}:\Phi(O(\mathbf{0},R);\mathbb{C}^3)\longrightarrow\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))$ as the orthogonal projection to the Hilbert subspace $\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}})):=\operatorname{Cl}\{(\hat{I}-\hat{\mathcal{D}})\boldsymbol{F}|\boldsymbol{F}\in\Phi(O(\mathbf{0},R);\mathbb{C}^3)\}\subset\Phi(O(\mathbf{0},R);\mathbb{C}^3)\cap\mathfrak{b}^2(O(\mathbf{0},R);\mathbb{C}^3)$, which satisfies $(\hat{I}-\hat{\mathcal{D}})=\hat{P}(\hat{I}-\hat{\mathcal{D}})=(\hat{I}-\hat{\mathcal{D}})\hat{P}$. From the identity $(\hat{I}-\hat{\mathcal{D}})(\hat{I}-\hat{P})\boldsymbol{F}=\mathbf{0}$ and the proper inclusion $\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))\subsetneqq\Phi(O(\mathbf{0},R);\mathbb{C}^3)$, we see that $0\in\sigma_p^{\Phi}(\hat{I}-\hat{\mathcal{D}})$. As $\boldsymbol{f}_{\ell m}=(\hat{I}-\hat{\mathcal{D}})((2\ell+1)\boldsymbol{f}_{\ell m}/\ell)=\hat{P}\boldsymbol{f}_{\ell m}\in\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}})$, it is also clear that $\langle\boldsymbol{f}_{\ell m},(\hat{I}-\hat{P})\boldsymbol{F}\rangle_{O(\mathbf{0},R)}=\langle(\hat{I}-\hat{P})\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}=0,\forall\ell m$ holds for every $\boldsymbol{F}\in\Phi(O(\mathbf{0},R);\mathbb{C}^3)$. Now that both sides of Eq. 2.53 vanish when $\boldsymbol{F}$ is orthogonal to $\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))$ (i.e. when $\boldsymbol{F}=(\hat{I}-\hat{P})\boldsymbol{F}$), it would suffice to prove Eq. 2.53 for cases where $\boldsymbol{F}=\hat{P}\boldsymbol{F}\in\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))$. Using the properties of spherical harmonics, one can verify that the Hilbert subspace $\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))$ satisfies

$$\begin{aligned}\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))&\subset\operatorname{Cl}\{\nabla f|f\in\mathfrak{b}^2(O(\mathbf{0},R);\mathbb{C})\cap W^{1,2}(O(\mathbf{0},R);\mathbb{C})\}\\&=\operatorname{Cl}\left\{\nabla f\left|f(\boldsymbol{r})=\sum_{\ell=0}^{\infty}\left[\sum_{m=-\ell}^{\ell}\frac{a_{\ell m}|\boldsymbol{r}|^\ell Y_{\ell m}(\theta,\phi)}{\sqrt{R^{2\ell+3}/(2\ell+3)}}\right],\sum_{\ell=0}^{\infty}\left[\sum_{m=-\ell}^{\ell}|a_{\ell m}|^2(\ell+1)^2\right]<+\infty\right.\right\}\\&=\left\{\nabla f\left|f(\boldsymbol{r})=\sum_{\ell=0}^{\infty}\left[\sum_{m=-\ell}^{\ell}\frac{a_{\ell m}|\boldsymbol{r}|^\ell Y_{\ell m}(\theta,\phi)}{\sqrt{R^{2\ell+3}/(2\ell+3)}}\right],\sum_{\ell=0}^{\infty}\left[\sum_{m=-\ell}^{\ell}|a_{\ell m}|^2(\ell+1)^2\right]<+\infty\right.\right\}\\&=\left\{\nabla f\left|\nabla f(\boldsymbol{r})=\sum_{\ell=1}^{\infty}\left[\sum_{m=-\ell}^{\ell}\frac{\sqrt{\ell(2\ell+3)}a_{\ell m}\boldsymbol{f}_{\ell m}}{R}\right],\sum_{\ell=1}^{\infty}\left[\sum_{m=-\ell}^{\ell}|a_{\ell m}|^2(\ell+1)^2\right]<+\infty\right.\right\}.\end{aligned}\tag{2.54}$$

Thus, we have a convergent spherical harmonic expansion $\boldsymbol{F}=\sum_{\ell=1}^{\infty}\sum_{m=-\ell}^{\ell}\langle\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}\boldsymbol{f}_{\ell m},\forall\boldsymbol{F}\in\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))$ according to the last line of Eq. 2.54. Using the convergence of the spherical harmonic expansion and the bound on the operator norm $\|\hat{I}-\hat{\mathcal{D}}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}\leq1$, we may conclude that

$$\begin{aligned}\left\|(\hat{I}-\hat{\mathcal{D}})\boldsymbol{F}-\sum_{\ell=0}^{L}\left[\sum_{m=-\ell}^{\ell}\frac{\ell\langle\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}\boldsymbol{f}_{\ell m}}{2\ell+1}\right]\right\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}&=\left\|(\hat{I}-\hat{\mathcal{D}})\sum_{\ell=L+1}^{\infty}\left[\sum_{m=-\ell}^{\ell}\langle\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}\boldsymbol{f}_{\ell m}\right]\right\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}\\&\leq\left\|\sum_{\ell=L+1}^{\infty}\left[\sum_{m=-\ell}^{\ell}\langle\boldsymbol{f}_{\ell m},\boldsymbol{F}\rangle_{O(\mathbf{0},R)}\boldsymbol{f}_{\ell m}\right]\right\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}\\&\to0,\quad\text{as }L\to\infty.\end{aligned}$$

Hence, we have verified Eq. 2.53 as claimed. In conjunction with this, we may also justify the relation $\sum_{\ell m}b_{\ell m}\boldsymbol{f}_{\ell m}=(\hat{I}-\hat{\mathcal{D}})\sum_{\ell m}(2\ell+1)b_{\ell m}\boldsymbol{f}_{\ell m}/\ell\in\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}})$ so long as $\sum_{\ell m}|b_{\ell m}|^2<+\infty$, which, in turn, strengthens Eq. 2.54 as

$$\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))\subset\left\{\sum_{\ell m}b_{\ell m}\boldsymbol{f}_{\ell m}\left|\sum_{\ell m}|b_{\ell m}|^2<+\infty\right.\right\}\subset\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}).$$

This shows that $\{\boldsymbol{f}_{\ell m}|\ell\in\mathbb{Z}\cap[1,+\infty);m\in\mathbb{Z}\cap[-\ell,\ell]\}$ forms a complete orthonormal basis for $\operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}}))=\operatorname{ran}^{\Phi}(\hat{I}-\hat{\mathcal{D}})$. Now, comparing Eq. 2.53 with the standard spectral decomposition of a Hermitian operator (see Ref. [44], p. 319), and using the fact that $0\in\sigma_p^{\Phi}(\hat{I}-\hat{\mathcal{D}})$, we may deduce the structure of the spectrum as

$$\sigma_p^{\Phi}(\hat{I}-\hat{\mathcal{D}})=\{\ell/(2\ell+1)|\ell=0,1,2,\dots\},\quad\sigma_c^{\Phi}(\hat{I}-\hat{\mathcal{D}})=\{1/2\},\quad\sigma^{\Phi}(\hat{I}-\hat{\mathcal{D}})=\sigma_p^{\Phi}(\hat{I}-\hat{\mathcal{D}})\cup\sigma_c^{\Phi}(\hat{I}-\hat{\mathcal{D}}).$$

Using Eq. 2.53, we may establish the spectral decomposition the Hermitian operator $\hat{H} := (\hat{\mathcal{G}} - \hat{\gamma})(\hat{I} + 2(\hat{\mathcal{G}} - \hat{\gamma})) : \Phi(O(\mathbf{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$ as below:

$$\hat{H}\boldsymbol{F} = \lim_{L\to\infty} \hat{H}_L \boldsymbol{F}, \quad \text{where } \hat{H}_L\boldsymbol{F} = -\sum_{\ell=1}^{L}\left[\sum_{m=-\ell}^{\ell} \frac{\ell\langle \boldsymbol{f}_{\ell m}, \boldsymbol{F}\rangle_{O(\mathbf{0},R)} \boldsymbol{f}_{\ell m}}{(2\ell+1)^2}\right], \quad \forall \boldsymbol{F} \in \Phi(O(\mathbf{0}, R); \mathbb{C}^3).$$

Here, every member in the family of bounded operators $\{\hat{H}_L : \Phi(O(\mathbf{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0}, R); \mathbb{C}^3) | L = 1, 2, \dots\}$ has finite-dimensional range, hence is a compact operator. Meanwhile, we have the uniform convergence

$$\lim_{L\to\infty}\left[\sup_{\boldsymbol{F}\in\Phi(O(\mathbf{0},R);\mathbb{C}^3)\setminus\{\mathbf{0}\}} \frac{\|\hat{H}\boldsymbol{F} - \hat{H}_L\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}}{\|\boldsymbol{F}\|_{L^2(V;\mathbb{C}^3)}}\right] = \lim_{L\to\infty} \frac{L+1}{(2L+3)^2} = 0,$$

which reveals $\hat{H} : \Phi(O(\mathbf{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$ as a compact operator. As the difference $\hat{\mathcal{G}}(\hat{I} + 2\hat{\mathcal{G}}) - \hat{H} = \hat{\gamma} + 2\hat{\mathcal{G}}\hat{\gamma} + 2\hat{\gamma}\hat{\mathcal{G}} - 2\hat{\gamma}^2$ is a compact operator, the operator $\hat{\mathcal{G}}(\hat{I} + 2\hat{\mathcal{G}}) : \Phi(O(\mathbf{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$ is also compact.

It is also true that $\{0, -1/2\} \subset \sigma_c^{\Phi}(\hat{\mathcal{G}})$. Otherwise, the assumption of $0 \in \rho^{\Phi}(\hat{\mathcal{G}})$ (resp. $-1/2 \in \rho^{\Phi}(\hat{\mathcal{G}})$) would lead to the conclusion that $(\hat{\mathcal{G}} - \hat{\gamma})$ (resp. $\hat{I}/2 + (\hat{\mathcal{G}} - \hat{\gamma})$) is a Fredholm operator, which contradicts the facts that $\dim\ker^{\Phi}(\hat{\mathcal{G}} - \hat{\gamma}) = +\infty$ and $\operatorname{ran}^{\Phi}(\hat{I}/2 + (\hat{\mathcal{G}} - \hat{\gamma})) \neq \operatorname{Cl}(\operatorname{ran}^{\Phi}(\hat{I}/2 + (\hat{\mathcal{G}} - \hat{\gamma})))$. This proves that the continuous spectrum consists of exactly two points $\sigma_c^{\Phi}(\hat{\mathcal{G}}) = \{0, -1/2\}$. ▮

Remark We note that unlike the conclusion $\sigma_c(\hat{\mathcal{G}}) \subset [-1, 0]$ for merely energetically admissible (square-integrable) inputs $\boldsymbol{E}_{\text{inc}} \in L^2(O(\mathbf{0}, R); \mathbb{C}^3)$, we now have $\sigma_c^{\Phi}(\hat{\mathcal{G}}) = \{0, -1/2\}$ for physically admissible inputs $\boldsymbol{E}_{\text{inc}} \in \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$, along with the conclusion that $\sigma_p^{\Phi}(\hat{\mathcal{G}})$ is countable. In other words, whenever the incident field is physically admissible $\boldsymbol{E}_{\text{inc}} \in \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$, the solution to the Born equation $(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}}$ can only have countably many singularities for $\chi \in \mathbb{C}$, without encountering any branch cut. Owing to some subtle differences between the Hilbert spaces $L^2(V; \mathbb{C}^3)$ and $\Phi(V; \mathbb{C}^3)$, the conclusion in this proposition does not contradict the claim in recent literature [16, 37] that the spectrum of light scattering on a sphere (in the Hilbert space $L^2(V; \mathbb{C}^3)$) may contain a line segment as its subset.

Such an absence of branch cut does not contradict the multi-valued square-root function for the refractive index $n = \sqrt{1+\chi}$, as can be *partly* illustrated by the following explicit expression of the Mie series (see Ref. [6], p. 122 or Ref. [16]) for light scattering on a dielectric sphere:

$$\begin{aligned}\boldsymbol{E}(\boldsymbol{r}) &= ((\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\text{inc}})(\boldsymbol{r})\\ &= \sum_{\ell=1}^{\infty}\sum_{m=-\ell}^{\ell}\left\{\frac{\alpha_{\ell m}\nabla\times[\boldsymbol{r}j_\ell(\sqrt{1+\chi}k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)]}{\sqrt{\ell(\ell+1)}kR\left[j_\ell(\sqrt{1+\chi}kR)\left.\frac{\mathrm{d}\,[xh_\ell^{(2)}(x)]}{\mathrm{d}\,x}\right|_{x=kR} - h_\ell^{(2)}(kR)\left.\frac{\mathrm{d}\,[yj_\ell(y)]}{\mathrm{d}\,y}\right|_{y=\sqrt{1+\chi}kR}\right]}\right.\\ &\quad\left.+\frac{i\beta_{\ell m}\nabla\times\nabla\times[\boldsymbol{r}j_\ell(\sqrt{1+\chi}k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)]}{\sqrt{\ell(\ell+1)}k^2R\left[(1+\chi)j_\ell(\sqrt{1+\chi}kR)\left.\frac{\mathrm{d}\,[xh_\ell^{(2)}(x)]}{\mathrm{d}\,x}\right|_{x=kR} - h_\ell^{(2)}(kR)\left.\frac{\mathrm{d}\,[yj_\ell(y)]}{\mathrm{d}\,y}\right|_{y=\sqrt{1+\chi}kR}\right]}\right\}, \quad |\boldsymbol{r}| < R \qquad (2.55)\end{aligned}$$

with

$$\begin{aligned}\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) &= \sum_{\ell=1}^{\infty}\sum_{m=-\ell}^{\ell}\left\{\frac{i\alpha_{\ell m}\nabla\times[\boldsymbol{r}j_\ell(k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)]}{\sqrt{\ell(\ell+1)}} - \frac{\beta_{\ell m}\nabla\times\nabla\times[\boldsymbol{r}j_\ell(k|\boldsymbol{r}|)Y_{\ell m}(\theta,\phi)]}{\sqrt{\ell(\ell+1)}k}\right\}, \quad |\boldsymbol{r}| < R,\\ j_\ell(z) &= (-z)^\ell\left(\frac{\mathrm{d}}{z\,\mathrm{d}\,z}\right)^\ell\left(\frac{\sin z}{z}\right) = z^\ell\sum_{m=0}^{\infty}\frac{(-z^2)^m}{2^m m!(2\ell+2m+1)!!}, \quad h_\ell^{(2)}(z) = (-z)^\ell\left(\frac{\mathrm{d}}{z\,\mathrm{d}\,z}\right)^\ell\left(\frac{ie^{-iz}}{z}\right).\end{aligned}$$

In each summand on the right-hand side of Eq. 2.55 with index ℓ, the common factors $[(1+\chi)k^2R^2]^{\ell/2}$ in the numerator and the denominator exactly cancel each other, leaving only a quotient of two power series in $1+\chi$ without any branch cut, thereby making the Mie series effectively single-valued. We call this a partial illustration for the absence of the branch cut, because in the Mie series representation, the incident field $\boldsymbol{E}_{\text{inc}}$ is not only taken to be transverse, but also a solution to the Helmholtz equation $(\nabla^2 + k^2)\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in O(\boldsymbol{0}, R)$. The conclusion of Proposition 2.9 shows that the transversality alone is sufficient for the elimination of the branch cut, and the Helmholtz equation is not essential for this purpose. ▮

It is customary (see for example, Ref. [9], p. 430) to classify multipole electromagnetic radiation into TE waves (*t*ransverse *e*lectric waves satisfying $\boldsymbol{r} \cdot \boldsymbol{E} = 0$) and TM waves (*t*ransverse *m*agnetic waves satisfying $\boldsymbol{r} \cdot (\nabla \times \boldsymbol{E}) = 0$). Let $g(|\boldsymbol{r}|)$ denote either the function $j_\ell(k|\boldsymbol{r}|)$ or $j_\ell(\sqrt{1+\chi}k|\boldsymbol{r}|)$, then we note that the relations

$$\boldsymbol{r} \cdot \{\nabla \times [\boldsymbol{r} g(|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)]\} = 0, \quad \boldsymbol{r} \cdot \{\nabla \times \nabla \times \nabla \times [\boldsymbol{r} g(|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)]\} = 0$$

are satisfied, leading to a natural decomposition of the corresponding Mie series for $\boldsymbol{E}(\boldsymbol{r}) = \boldsymbol{E}^{\text{TE}}(\boldsymbol{r}) + \boldsymbol{E}^{\text{TM}}(\boldsymbol{r})$ and $\boldsymbol{E}_{\text{inc}}(\boldsymbol{r}) = \boldsymbol{E}^{\text{TE}}_{\text{inc}}(\boldsymbol{r}) + \boldsymbol{E}^{\text{TM}}_{\text{inc}}(\boldsymbol{r})$ into TE waves and TM waves. It is evident from Eq. 2.55 that the TE and TM waves are decoupled in the Mie scattering solution as $\boldsymbol{E}^{\text{TE}} = (\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}^{\text{TE}}_{\text{inc}}$ and $\boldsymbol{E}^{\text{TM}} = (\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}^{\text{TM}}_{\text{inc}}$.

To facilitate further discussion, we define two sequences of analytic functions $\{f^{\text{TE}}_\ell(\chi) | \ell = 1, 2, \ldots\}$ and $\{f^{\text{TM}}_\ell(\chi) | \ell = 1, 2, \ldots\}$ in the complex-variable $\chi \in \mathbb{C}$ related to the denominators in the Mie series for TE and TM waves:

$$f^{\text{TE}}_\ell(\chi) := \frac{1}{[(1+\chi)k^2R^2]^{\ell/2}} \left[j_\ell(\sqrt{1+\chi}kR) \left.\frac{\mathrm{d}\,[x h^{(2)}_\ell(x)]}{\mathrm{d}\,x}\right|_{x=kR} - h^{(2)}_\ell(kR) \left.\frac{\mathrm{d}\,[y j_\ell(y)]}{\mathrm{d}\,y}\right|_{y=\sqrt{1+\chi}kR} \right], \tag{2.56a}$$

$$f^{\text{TM}}_\ell(\chi) := \frac{1}{[(1+\chi)k^2R^2]^{\ell/2}} \left[(1+\chi) j_\ell(\sqrt{1+\chi}kR) \left.\frac{\mathrm{d}\,[x h^{(2)}_\ell(x)]}{\mathrm{d}\,x}\right|_{x=kR} - h^{(2)}_\ell(kR) \left.\frac{\mathrm{d}\,[y j_\ell(y)]}{\mathrm{d}\,y}\right|_{y=\sqrt{1+\chi}kR} \right] \tag{2.56b}$$

Proposition 2.10 (Analytic Properties of Mie Coefficients) *For the light scattering problem on spherical dielectric* $(\hat{I} - \chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}} \in \Phi(O(\boldsymbol{0}, R), \mathbb{C}^3)$*, the eigenvalues are enumerated by* $\sigma^{\Phi}_p(\hat{\mathcal{G}}) = \sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) \cup \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$*, where*

$$\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) = \{\lambda \in \mathbb{C} \setminus \{0\} | f^{\text{TE}}_\ell(1/\lambda) = 0, \ell = 1, 2, \ldots\}, \quad \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}}) = \{\lambda \in \mathbb{C} \setminus \{0\} | f^{\text{TM}}_\ell(1/\lambda) = 0, \ell = 1, 2, \ldots\}. \tag{2.57}$$

Neither $\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}})$ *nor* $\sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$ *contains the point* $\{-1\}$*. Moreover,* $\sigma^{\Phi}_p(\hat{\mathcal{G}})$ *is a countable set, and* $\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}})$ *has no accumulation points other than* $\{0\}$*.*

Proof From the boundary matching condition employed in the construction of the Mie series (see Ref. [6], p. 123 and p. 154), we have the eigenequation $\hat{\mathcal{G}}\boldsymbol{E}^{\text{TE}}_\lambda = \lambda\boldsymbol{E}^{\text{TE}}_\lambda$ for

$$\boldsymbol{E}^{\text{TE}}_\lambda(\boldsymbol{r}) = \nabla \times [\boldsymbol{r} j_\ell(\sqrt{1+\lambda^{-1}}k|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)], \quad f^{\text{TE}}_\ell(1/\lambda) = 0,$$

and the eigenequation $\hat{\mathcal{G}}\boldsymbol{E}^{\text{TM}}_\lambda = \lambda\boldsymbol{E}^{\text{TM}}_\lambda$ for

$$\boldsymbol{E}^{\text{TM}}_\lambda(\boldsymbol{r}) = \nabla \times \nabla \times [\boldsymbol{r} j_\ell(\sqrt{1+\lambda^{-1}}k|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)], \quad f^{\text{TM}}_\ell(1/\lambda) = 0,$$

where the eigenvectors $\boldsymbol{E}^{\text{TE}}_\lambda$ and $\boldsymbol{E}^{\text{TM}}_\lambda$ represent free vibrations without incident field. This proves the set inclusion relation $\sigma^{\Phi}_p(\hat{\mathcal{G}}) \supset \sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) \cup \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$, where $\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}})$ and $\sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$ are given by Eq. 2.57.

To prove the set inclusion in the reverse direction $\sigma^{\Phi}_p(\hat{\mathcal{G}}) \subset \sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) \cup \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$, we first note that any eigenvector $\boldsymbol{E}_\lambda \in \Phi(O(\boldsymbol{0}, R); \mathbb{C}^3)$ subordinate to an eigenvalue $\lambda \in \sigma^{\Phi}_p(\hat{\mathcal{G}}) \subset \mathbb{C} \setminus \{0\}$ indeed must satisfy the Helmholtz equation $\nabla^2\boldsymbol{E}_\lambda(\boldsymbol{r}) + (1+\lambda^{-1})k^2\boldsymbol{E}_\lambda(\boldsymbol{r}) = \boldsymbol{0}, \forall \boldsymbol{r} \in O(\boldsymbol{0}, R)$, according to the smoothness argument carried out in Proposition 2.5. Meanwhile, a transverse vector field satisfying the Helmholtz equation can be expanded with respect to a complete orthogonal basis set

$$\{\nabla \times [\boldsymbol{r} j_\ell(\sqrt{1+\lambda^{-1}}k|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)], \nabla \times \nabla \times [\boldsymbol{r} j_\ell(\sqrt{1+\lambda^{-1}}k|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)] | \ell = 1, 2, \ldots; m = -\ell, \ldots, \ell\},$$

which is the totality of TE and TM waves with wave number $\sqrt{1+\lambda^{-1}}k$. Therefore, the eigenvector $\boldsymbol{E}_\lambda$ is either exactly parallel to a member of this complete orthogonal basis set, or it is a linear combination of TE or TM waves subordinate to the same eigenvalue λ. This proves that there are no eigenvalues of $\hat{\mathcal{G}} : \Phi(O(\boldsymbol{0}, R), \mathbb{C}^3) \longrightarrow \Phi(O(\boldsymbol{0}, R), \mathbb{C}^3)$ that lie outside the set $\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) \cup \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$.

Therefore, the equality $\sigma^{\Phi}_p(\hat{\mathcal{G}}) = \sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}}) \cup \sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$ must hold for $\sigma^{\Phi,\text{TE}}_p(\hat{\mathcal{G}})$ and $\sigma^{\Phi,\text{TM}}_p(\hat{\mathcal{G}})$ given in Eq. 2.57.

We have

$$\lim_{\chi\to-1} f_\ell^{\mathrm{TE}}(\chi) = -\frac{kRh_{\ell+1}^{(2)}(kR)}{(2\ell+1)!!},\quad \lim_{\chi\to-1} f_\ell^{\mathrm{TM}}(\chi) = -\frac{(\ell+1)h_{\ell}^{(2)}(kR)}{(2\ell+1)!!},$$

whereas the Wroński determinant

$$\det\begin{pmatrix} \operatorname{Re} h_\nu^{(2)}(x) & \operatorname{Im} h_\nu^{(2)}(x) \\ \mathrm{d}\operatorname{Re} h_\nu^{(2)}(x)/\,\mathrm{d}\,x & \mathrm{d}\operatorname{Im} h_\nu^{(2)}(x)/\,\mathrm{d}\,x \end{pmatrix} = -\frac{1}{x^2},$$

implies $h_\nu^{(2)}(kR) \neq 0$ for $kR \neq 0$ and $\nu \in \mathbb{C}$. This shows that $\chi = -1$ is not a root of $f_\ell^{\mathrm{TE}}(\chi)$ or $f_\ell^{\mathrm{TM}}(\chi)$, hence $-1 \notin \sigma_p^{\Phi,\mathrm{TE}}(\hat{\mathcal{G}}) \cup \sigma_p^{\Phi,\mathrm{TM}}(\hat{\mathcal{G}})$.

For each fixed ℓ, the two analytic functions $f_\ell^{\mathrm{TE}}(\chi)$ and $f_\ell^{\mathrm{TM}}(\chi)$ defined in Eqs. 2.56a and 2.56b are not identically vanishing, so each of them must have countably many roots without a finite accumulation point in $\mathbb{C}$ (see Ref. [89], p. 127). As a result, $\sigma_p^{\Phi}(\hat{\mathcal{G}})$ is the union of countably many countable sets, hence a countable set.

Using the definition of TE wave $\boldsymbol{r}\cdot\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}) = 0, |\boldsymbol{r}| < R$ and the transversality condition $\nabla\cdot\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}) = 0, |\boldsymbol{r}| < R$, we may rewrite the Born equation $(\hat{I} - \chi\hat{\mathcal{G}})\boldsymbol{E}^{\mathrm{TE}} = \boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TE}}$ as

$$\begin{aligned}\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TE}}(\boldsymbol{r}) &= \boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}) - \chi k^2 \iiint_{|\boldsymbol{r}'|<R} \frac{\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}' \\ &\quad +\chi \lim_{R'\to R-0}\left\{\nabla\left[\nabla \iiint_{R'<|\boldsymbol{r}'|<R} \frac{\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}'\right] - \frac{1}{R'}\oiint_{|\boldsymbol{r}'|=R'} \frac{\boldsymbol{r}'\cdot\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}\,S'\right\} \\ &= \boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}) - \chi k^2 \iiint_{|\boldsymbol{r}'|<R} \frac{\boldsymbol{E}^{\mathrm{TE}}(\boldsymbol{r}')e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \quad |\boldsymbol{r}| < R. \end{aligned}\tag{2.58}$$

This is effectively an equation $(\hat{I} - \chi k^2\hat{\mathcal{C}})\boldsymbol{E}^{\mathrm{TE}} = \boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TE}}$, where the convolution transform $\hat{\mathcal{C}} : L^2(O(\mathbf{0},R),\mathbb{C}^3) \longrightarrow L^2(O(\mathbf{0},R),\mathbb{C}^3)$ (as defined in Eq. 1.2) is a compact operator satisfying the Hilbert-Schmidt bound

$$\iiint_{|\boldsymbol{r}'|<R}\left(\iiint_{|\boldsymbol{r}|<R}\left|\frac{e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\right|^2\mathrm{d}^3\boldsymbol{r}\right)\mathrm{d}^3\boldsymbol{r}' < +\infty.$$

Therefore, we have $\sigma_p^{\Phi,\mathrm{TE}}(\hat{\mathcal{G}}) \subset \sigma_p(k^2\hat{\mathcal{C}})$, and the set $\sigma_p^{\Phi,\mathrm{TE}}(\hat{\mathcal{G}})$ may only accumulate at the point $\{0\}$, according to the Riesz-Schauder theory applicable to the compact operator $k^2\hat{\mathcal{C}}$. ▮

Remark For the specific case $kR = 1$, numerical evidence has been previously reported (see Fig. 4.1 in Ref. [16] and the accompanying text) that the roots of the denominators in the Mie series aggregate around the point $n^2 = 1 + \chi = -1$. This is consistent with the general conclusion (valid for all $kR > 0$) in Proposition 2.9, which pinpoints $\chi = -2 \Leftrightarrow \lambda = 1/\chi = -1/2$ as a potential accumulation point of eigenvalues.

In Fig. 1, we numerically illustrate the existence of accumulation points $\{0, -1/2\}$ in the physical point spectrum $\sigma_p^{\Phi}(\hat{\mathcal{G}})$ by two specific cases $kR = 1/2$ and $kR = 4\pi$. While the panoramic views of eigenvalues look different in the two cases, the locations of the accumulation points remain invariant. We use circular symbols ($\circ$) to plot the subset of eigenvalues $\sigma_p^{\Phi,\mathrm{TE}}(\hat{\mathcal{G}})$ related to TE waves, and use cross symbols ($\times$) to plot the subset of eigenvalues $\sigma_p^{\Phi,\mathrm{TM}}(\hat{\mathcal{G}})$ related to TM waves. It is graphically evident that $\sigma_p^{\Phi,\mathrm{TM}}(\hat{\mathcal{G}})$ may accumulate at the point $\{-1/2\}$, but $\sigma_p^{\Phi,\mathrm{TE}}(\hat{\mathcal{G}})$ may not.

Heuristically speaking, Eq. 2.56b hints at a universal accumulation point $1/\chi = -1/2$ in $\sigma_p^{\Phi,\mathrm{TM}}(\hat{\mathcal{G}})$ via the asymptotic expansion

$$(kR)^\ell f_\ell^{\mathrm{TM}}(\chi) = -\frac{i[\ell(\chi+2)+1]}{(2\ell+1)kR} + \frac{i\chi\{\ell[-\chi+2\ell(\chi+2)+4]+3\}kR}{2(2\ell-1)(2\ell+1)(2\ell+3)}\left[1+O\left(\frac{k^2R^2}{\ell}\right)\right].\tag{2.59}$$

Here, in passage to the limit $\ell\to\infty$, the right-hand side expression in Eq. 2.59 tends to zero if $\chi = -2$, and tends to a non-vanishing value if $\chi \neq -2$. Roughly speaking, the root of $f_\ell^{\mathrm{TM}}(\chi)$ (as approximately computed from the first two leading terms on the right-hand side of Eq. 2.59) asymptotically behaves like

$$\chi_\ell \sim -2 - \frac{1}{\ell}\left(1 + \frac{k^2R^2}{\ell}\right),\quad \ell\to\infty,$$

which can get arbitrarily close to the point $\{-2\}$, if ℓ is chosen to be sufficiently large. ▮



Figure 1 Plots of the eigenvalues $\sigma_p^\Phi(\hat{\mathcal{G}}) = \{\lambda \in \mathbb{C} \setminus \{0\} | f_\ell^{\mathrm{TE}}(1/\lambda) = 0 \text{ or } f_\ell^{\mathrm{TM}}(1/\lambda) = 0, \ell = 1, 2, \dots\}$ for two specific cases $kR = 1/2$ (a) and $kR = 4\pi$ (b). The complex roots of $f_\ell^{\mathrm{TE}}(1/\lambda)$ and $f_\ell^{\mathrm{TM}}(1/\lambda)$ are searched numerically and checked against the singularities in the phase diagrams of $\arg f_\ell^{\mathrm{TE}}(1/\lambda)$ and $\arg f_\ell^{\mathrm{TM}}(1/\lambda)$. An example of $\arg f_1^{\mathrm{TM}}(1/\lambda)$ is shown in the background of panel a. The values of ℓ are color-coded, as indicated by the scale bar accompanying panel b. Both panels a and b give graphical evidence of the accumulation points $\{0, -1/2\}$ (pinpointed by dashed arrows). In panel a, only a limited subset of $\sigma_p^\Phi(\hat{\mathcal{G}})$ is shown, corresponding to the eigenvalues located by numerical root-finding of $f_\ell^{\mathrm{TE}}(1/\lambda)$ and $f_\ell^{\mathrm{TM}}(1/\lambda)$ for $\ell = 1, \dots, 20$. Panel b further includes contributions from the roots of $f_\ell^{\mathrm{TM}}(1/\lambda), \ell = 24, \dots, 40$ near $\lambda = -1/2$.

The electric enhancement ratio (EER)

$$\mathrm{EER}[\boldsymbol{E}; \hat{\mathcal{B}}\boldsymbol{E}] := \iiint_V |\boldsymbol{E}(\boldsymbol{r})|^2 \,\mathrm{d}^3\boldsymbol{r} \Big/ \iiint_V |(\hat{\mathcal{B}}\boldsymbol{E})(\boldsymbol{r})|^2 \,\mathrm{d}^3\boldsymbol{r} \tag{2.60}$$

is a dimensionless quantity of practical interest in the problem of light-matter interaction described by the Born equation $\hat{\mathcal{B}}\boldsymbol{E} = (\hat{I} - \chi\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\mathrm{inc}}$. The functional EER (Eq. 2.60) evaluates the energy enhancement inside the volume V in the presence/absence of the dielectric. In the optical parlance, EER measures the "focusing power" of the dielectric medium.

Proposition 2.11 *For the light scattering problem concerning a spherical dielectric* $V = O(\mathbf{0}, R)$ *with* $-1/2 \in \sigma_c^\Phi(\hat{\mathcal{G}})$, *the electric enhancement is unbounded for* $\chi = -2$*:*

$$\sup_{0 < \iiint_V |\boldsymbol{E}(\boldsymbol{r})|^2 \,\mathrm{d}^3\boldsymbol{r} < +\infty} \mathrm{EER}[\boldsymbol{E}; (\hat{I} + 2\hat{\mathcal{G}})\boldsymbol{E}] = +\infty.$$

Proof From the definition of the continuous spectrum, the range of the operator $(\hat{I} + 2\hat{\mathcal{G}}) : \Phi(O(\mathbf{0}, R); \mathbb{C}^3) \longrightarrow \Phi(O(\mathbf{0}, R); \mathbb{C}^3)$ is dense in, but not equal to $\Phi(O(\mathbf{0}, R); \mathbb{C}^3)$. Thus, the operator $(\hat{I} + 2\hat{\mathcal{G}})$ does not have a bounded inverse, according to the open mapping theorem (see Ref. [44], p. 75). Written symbolically, this amounts to the relation $\sup_{0 < \|\boldsymbol{E}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} < +\infty} \mathrm{EER}[\boldsymbol{E}; (\hat{I} + 2\hat{\mathcal{G}})\boldsymbol{E}] = +\infty$. ▮

Remark From Eq. 2.59, it is also clear that when $\chi = -2$, we have the following asymptotic behavior of the noise amplification ratio for the inverse Born operator $\hat{\mathcal{B}}^{-1} = (\hat{I} - \chi\hat{\mathcal{G}})^{-1} = (\hat{I} + 2\hat{\mathcal{G}})^{-1}$:

$$\frac{\|(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{w}_{\ell m}^{\mathrm{TM}}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}}{\|\boldsymbol{w}_{\ell m}^{\mathrm{TM}}\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)}} \sim (2\ell + 1)kR, \quad \text{where} \quad \boldsymbol{w}_{\ell m}^{\mathrm{TM}}(\boldsymbol{r}) = \nabla \times \nabla \times [\boldsymbol{r} j_\ell(k|\boldsymbol{r}|) Y_{\ell m}(\theta, \phi)].$$



Figure 2 Plots of the function $A_\ell(\chi, kR) := 1/|(kR)^\ell f_\ell^{\mathrm{TM}}(\chi)|$ for two specific cases $kR = 1/2$ (a) and $kR = 4\pi$ (b). At the critical susceptibility $\chi = -2$, the function $A_\ell(-2, kR) \sim (2\ell+1)kR, \ell \gg (kR)^2$ has unbounded growth as $\ell \to +\infty$, indicating an unbounded electric enhancement ratio for $\chi = -2$. In contrast, for each non-accretive susceptibility $\operatorname{Im}\chi \le 0$ other than the critical point $\chi = -2$, the function $A_\ell(\chi, kR)$ is bounded for all $\ell = 1, 2, \dots$, with the limit value $\lim_{\ell\to\infty} A_\ell(\chi, kR) = 2kR/|\chi+2|$ (asymptotes shown in dashed lines). This is partly illustrated by two special situations $\chi = -1$ and $\chi = -i$.

Therefore, for a TM incident wave given by

$$\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}} = \sum_{\ell m} a_{\ell m}\boldsymbol{w}_{\ell m}^{\mathrm{TM}}, \quad \text{satisfying } \left\| \sum_{\ell m} a_{\ell m}\boldsymbol{w}_{\ell m}^{\mathrm{TM}} \right\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} < +\infty \text{ and } \left\| \sum_{\ell m} \ell a_{\ell m}\boldsymbol{w}_{\ell m}^{\mathrm{TM}} \right\|_{L^2(O(\mathbf{0},R);\mathbb{C}^3)} = +\infty, \tag{2.61}$$

we have $\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}} \in \Phi(O(\mathbf{0}, R), \mathbb{C}^3)$ but there is no such a square integrable field $\boldsymbol{E} \in L^2(O(\mathbf{0}, R), \mathbb{C}^3)$ satisfying $(\hat{I} + 2\hat{\mathcal{G}})\boldsymbol{E} = \boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}}$. This shows that the existence theorem fails when $\chi = -2$, along with an unbounded electric enhancement.

The critical behavior of $\chi = -2$ is illustrated by two explicit examples in Fig. 2. It is easy to verify that for TM waves with large ℓ values, the electric enhancement ratio has an asymptotic expression

$$\mathrm{EER}[(\hat{I} - \chi\hat{\mathcal{G}})^{-1}\boldsymbol{w}_{\ell m}^{\mathrm{TM}}; \boldsymbol{w}_{\ell m}^{\mathrm{TM}}] \sim \frac{1}{|(kR)^\ell f_\ell^{\mathrm{TM}}(\chi)|^2},$$

so we plot $A_\ell(\chi, kR) := 1/|(kR)^\ell f_\ell^{\mathrm{TM}}(\chi)|$ in Fig. 2 to appreciate the criticality of the point $\chi = -2$.

We note that the violence of existence for light scattering on a dielectric with $\chi = -2$ does not cause an “infinity catastrophe” (finite input causing infinite response) in reality, for several reasons.

Firstly, as an *ad hoc* explanation, we note that the pathological incident field given in Eq. 2.61 cannot analytically continue across the boundary, that is, the spherical harmonic expansion $\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}}(\boldsymbol{r}) = \sum_{\ell m} a_{\ell m}\boldsymbol{w}_{\ell m}^{\mathrm{TM}}(\boldsymbol{r})$ fails to converge for $|\boldsymbol{r}| > R$. This disqualifies $\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}}$ as an experimentally available incident field that has well-defined values outside the scattering medium. In other words, the failure of the existence theorem for $\chi = -2$ may well be an intrinsic defect of our admissibility criteria: we had only imposed the “energy admissibility” and “physical admissibility” on the internal field, but had not censored its extensibility to the space outside the dielectric. It was just remarkable that the (perhaps) glaring omission of such an extensibility censorship did not undermine the light scattering problem for all the physically available susceptibility values $\operatorname{Im}\chi \le 0$ other than the critical point $\chi = -2$.

Secondly, we might argue that even when the non-extensible localized incident field $\boldsymbol{E}_{\mathrm{inc}}^{\mathrm{TM}}$ (Eq. 2.61) can be experimentally realized, it does not cause an infinity catastrophe in the real world, where all the dielectric media are made of atoms and molecules that come in finite sizes. The Born equation formulated out of the macroscopic Maxwell equations with idealized dielectric geometry may well break down on a microscopic length scale, i.e. in the limit of $\ell \to \infty$. Therefore, a reasonable application of the Born equation to light scattering inevitably calls for a physical cut-off of large ℓ modes, thereby precluding the occurrence of any type of infinite response in a dielectric sphere.

Lastly, even for non-spherical dielectric media, where the above two rationales do not necessarily generalize well, there are physical mechanisms fulfilling the principle of "nature abhors infinity". Admittedly, when the susceptibility χ comes close to the critical point of $\{-2\}$, we have the resolvent estimate (see Ref. [40], p. 149):

$$\|(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\|_{\Phi(V;\mathbb{C}^3)} := \sup_{\forall \boldsymbol{E}_{\mathrm{inc}}\in\Phi(V,\mathbb{C}^3)\setminus\{0\}} \frac{\|(\hat{I}-\chi\hat{\mathcal{G}})^{-1}\boldsymbol{E}_{\mathrm{inc}}\|_{L^2(V;\mathbb{C}^3)}}{\|\boldsymbol{E}_{\mathrm{inc}}\|_{L^2(V;\mathbb{C}^3)}} \geq \frac{1}{|\chi|\operatorname{dist}(1/\chi,\sigma^{\Phi}(\hat{\mathcal{G}}))} \geq \frac{2}{|\chi+2|},$$

which suggests that a dielectric material with $\chi\approx -2$ can possess extreme capability to garner electromagnetic energy (that is, to form a "lens of super focusing power", in the optical parlance). However, it should be noted that the ansatz about linear dielectric response $\boldsymbol{D}=(1+\chi)\epsilon_0\boldsymbol{E}$ is reliable only for moderately large field intensities $|\boldsymbol{E}|^2$, beyond which the light scattering problem needs reformulation to accommodate non-linear optics effects (cf. the "non-linear saturation" argument in Ref. [78]). Even in the absence of non-linear optics effects, an extremely intense electromagnetic field may well cause electric leakage, induce chemical change, or pave way for pulverization/disintegration — there is still a variety of feedback mechanisms to drag the dielectric material away from the critical point $\chi=-2$, thus preventing an infinity catastrophe. ∎

S2.3.6. Resonance Modes in Acoustic and Electromagnetic Scattering

Here, we give a brief survey of acoustic and electromagnetic scattering problems posed in d-dimensional ($d=1,2,\dots$) Euclidean spaces $\mathbb{R}^d$. Our major aim is to compare the spectral properties of the scalar/vector wave scattering in various spatial dimensions and point out the distinctive feature (as well as the outstanding difficulty) of electromagnetic scattering.

Definition Let $u,u_{\mathrm{inc}}\in L^2(\Omega;\mathbb{C})$ be two square-integrable scalar fields defined in a bounded open set $\Omega\Subset\mathbb{R}^d$, $\tilde{u}\in L^2(\mathbb{R}^d;\mathbb{C})$ be the Fourier transform $\tilde{u}(\boldsymbol{q}):=\int\cdots\int_\Omega u(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^d\boldsymbol{r}$, then the acoustic scattering problem is given by

$$\begin{aligned} u_{\mathrm{inc}}(\boldsymbol{r}) &= u(\boldsymbol{r})-\lim_{\varepsilon\to0^+}\frac{\chi k^2}{(2\pi)^d}\fint\cdots\fint_{\mathbb{R}^d}\frac{\tilde{u}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^d\boldsymbol{q} \\ &= u(\boldsymbol{r})-\chi\int\cdots\int_\Omega u(\boldsymbol{r}')K^{(d)}(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^d\boldsymbol{r}',\quad \boldsymbol{r}\in\Omega, \end{aligned} \tag{2.62}$$

where the convolution kernel $K^{(d)}(\boldsymbol{r},\boldsymbol{r}')$ is given by

$$K^{(d)}(\boldsymbol{r},\boldsymbol{r}')=\frac{k^{(d+2)/2}H^{(2)}_{(d-2)/2}(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4i(2\pi|\boldsymbol{r}-\boldsymbol{r}'|)^{(d-2)/2}},$$

and $H^{(2)}_\ell$ denotes the ℓth order cylindrical Hankel function of the second kind.

Let $\boldsymbol{E},\boldsymbol{E}_{\mathrm{inc}}\in L^2(\Omega;\mathbb{C}^3)$ be two square-integrable vector fields defined in a bounded open set $\Omega\Subset\mathbb{R}^d$ ($d\leq3$), $\tilde{\boldsymbol{E}}\in L^2(\mathbb{R}^d;\mathbb{C}^3)$ be the Fourier transform $\tilde{\boldsymbol{E}}(\boldsymbol{q}):=\int\cdots\int_\Omega \boldsymbol{E}(\boldsymbol{r})e^{i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^d\boldsymbol{r}$, then the electromagnetic scattering problem is given by

$$\begin{aligned} \boldsymbol{E}_{\mathrm{inc}}(\boldsymbol{r}) &= \boldsymbol{E}(\boldsymbol{r})+\lim_{\varepsilon\to0^+}\frac{\chi}{(2\pi)^d}\fint\cdots\fint_{\mathbb{R}^d}\frac{\boldsymbol{q}[\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})]-k^2\tilde{\boldsymbol{E}}(\boldsymbol{q})}{|\boldsymbol{q}|^2-(k-i\varepsilon)^2}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}\,\mathrm{d}^d\boldsymbol{q} \\ &= \boldsymbol{E}(\boldsymbol{r})-\chi\int\cdots\int_\Omega \boldsymbol{E}(\boldsymbol{r}')K^{(d)}(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^d\boldsymbol{r}'-\frac{\chi}{k^2}\nabla\left[\nabla\cdot\int\cdots\int_\Omega \boldsymbol{E}(\boldsymbol{r}')K^{(d)}(\boldsymbol{r},\boldsymbol{r}')\,\mathrm{d}^d\boldsymbol{r}'\right],\quad \boldsymbol{r}\in\Omega, \end{aligned} \tag{2.63}$$

where the scalar product $\boldsymbol{q}\cdot\tilde{\boldsymbol{E}}(\boldsymbol{q})$ is evaluated with regard to $\boldsymbol{q}\in\mathbb{R}^d\subset\mathbb{R}^3$ and $\tilde{\boldsymbol{E}}\in\mathbb{C}^3$. In particular, Eq. 2.63 becomes the Born equation for light scattering when $d=3$. ∎

Remark In case $d=3$, Eq. 2.62 can be written as

$$u_{\mathrm{inc}}(\boldsymbol{r})=u(\boldsymbol{r})-\chi\iiint_\Omega u(\boldsymbol{r}')\frac{k^2e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{4\pi|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^3\boldsymbol{r}', \tag{2.64}$$

or symbolically as $(\hat{I}-\chi k^2\hat{\mathcal{C}})u=u_{\mathrm{inc}}$ (cf. Eq. 1.2). While Eq. 2.64 is primarily a model for scattering of scalar waves (e.g. acoustic waves, de Broglie waves among others) in three-dimensional Euclidean space $\mathbb{R}^3$, it has been sometimes used as a "scalar wave approximation" for light scattering (see Ref. [6], p. 198, as well as Refs. [90, 91]). Although the theories of the scalar wave scattering have been well-developed (see Ref. [56] or Ref. [57]), the fruits of acoustic scattering cannot be hastily transplanted to electromagnetic scattering problems, due to intrinsic discrepancies (cf. Theorem 2.1 and Proposition 2.12) between their spectral properties.

In case $d = 1, 2$, Eq. 2.63 can be used to describe the scattering of electromagnetic waves by dielectric structures with (idealized) translational symmetry along certain directions. For instance, in the $d = 1$ case, the bounded open set $\Omega \subset \mathbb{R}$ is a union of open intervals corresponding to the spatial region $\Omega \times \mathbb{R}^2$ occupied by parallel dielectric slabs; in the $d = 2$ case, the shape $\Omega \subset \mathbb{R}^2$ can be interpreted as the uniform cross-section of a cylindrical dielectric standing in the spatial region $\Omega \times \mathbb{R}$. To cast Eq. 2.63 with a well-defined Fourier transform, we have conveniently assumed that the electric field does not vary along the directions where the dielectric boasts translational symmetry, thus precluding the scenario of "oblique incidence".

When the electric polarization is perpendicular to Ω (say, both $\boldsymbol{E}$ and $\boldsymbol{E}_{\text{inc}}$ are parallel to the z-axis while $\Omega \subset \mathbb{R}^2$ sits in the xy-plane), then the term $\boldsymbol{q}[\boldsymbol{q} \cdot \tilde{\boldsymbol{E}}(\boldsymbol{q})]$ in Eq. 2.63 vanishes, and the vector wave scattering equation *exactly* degenerates to a scalar wave scattering equation. ∎

To ease further discussion, we will abbreviate Eq. 2.62 as $(\hat{I} - \chi\hat{\mathcal{C}}^{(d)})u = u_{\text{inc}}$ and Eq. 2.63 as $(\hat{I} - \chi\hat{\mathcal{G}}^{(d)})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}}$ by explicitly acknowledging the dimensionality in question. In parallel to Eq. 2.64, it is possible to spell out the equation $(\hat{I} - \chi\hat{\mathcal{C}}^{(d)})u = u_{\text{inc}}$ for $d = 1, 2$ (cf. Ref. [90]) as follows

$$u_{\text{inc}}(\boldsymbol{r}) = u(\boldsymbol{r}) - \chi \int_{\Omega\subset\mathbb{R}} u(\boldsymbol{r}') \frac{k e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{2i} \,\mathrm{d}\,\boldsymbol{r}' \tag{2.65}$$

$$u_{\text{inc}}(\boldsymbol{r}) = u(\boldsymbol{r}) - \chi \iint_{\Omega\subset\mathbb{R}^2} u(\boldsymbol{r}') \frac{k^2 H_0^{(2)}(k|\boldsymbol{r}-\boldsymbol{r}'|)}{4i} \,\mathrm{d}^2\boldsymbol{r}' \tag{2.66}$$

where $H_0^{(2)}$ denotes the zeroth order cylindrical Hankel function of the second kind, with logarithmic divergence at short distances: $H_0^{(2)}(k|\boldsymbol{r}-\boldsymbol{r}'|) \sim -2i[\ln(k|\boldsymbol{r}-\boldsymbol{r}'|)]/\pi + O(1)$. In general, for $d > 2$, the convolution kernel $K^{(d)}(\boldsymbol{r}, \boldsymbol{r}')$ related to $\hat{\mathcal{C}}^{(d)}$ exhibits power law divergence at short distances: $K^{(d)}(\boldsymbol{r}, \boldsymbol{r}') = O(|\boldsymbol{r}-\boldsymbol{r}'|^{2-d})$.

Proposition 2.12 (Acoustic Scattering and Universal Compactness) *The operators* $\hat{\mathcal{C}}^{(d)} : L^2(\Omega; \mathbb{C}) \longrightarrow L^2(\Omega; \mathbb{C})$ *are compact for any finite spatial dimensions* $d \geq 1$.

Proof For dimensions $d = 1, 2, 3$, direct computation reveals that the Hilbert-Schmidt bound is satisfied

$$\int \cdots \int_{\Omega} \left(\int \cdots \int_{\Omega} |K^{(d)}(\boldsymbol{r}, \boldsymbol{r}')|^2 \,\mathrm{d}^d\boldsymbol{r} \right) \mathrm{d}^d\boldsymbol{r}' < +\infty,$$

hence the corresponding operators $\hat{\mathcal{C}}^{(d)} : L^2(\Omega; \mathbb{C}) \longrightarrow L^2(\Omega; \mathbb{C})$ are compact. For $d > 3$, we may define the cut-off convolution kernel $K_\varepsilon^{(d)}(\boldsymbol{r}, \boldsymbol{r}'), \varepsilon > 0$ as

$$K_\varepsilon^{(d)}(\boldsymbol{r}, \boldsymbol{r}') = \begin{cases} K^{(d)}(\boldsymbol{r}, \boldsymbol{r}'), & |\boldsymbol{r}-\boldsymbol{r}'| > \varepsilon \\ 0, & |\boldsymbol{r}-\boldsymbol{r}'| \leq \varepsilon \end{cases}$$

and denote by $\hat{\mathcal{C}}_\varepsilon^{(d)} : L^2(\Omega; \mathbb{C}) \longrightarrow L^2(\Omega; \mathbb{C})$ the integral operator induced by the cut-off convolution kernel $K_\varepsilon^{(d)}(\boldsymbol{r}, \boldsymbol{r}')$. Using the Hilbert-Schmidt bounds, one can readily check the compactness of $\hat{\mathcal{C}}_\varepsilon^{(d)} : L^2(\Omega; \mathbb{C}) \longrightarrow L^2(\Omega; \mathbb{C})$ for any $\varepsilon > 0$. Meanwhile, according to the Young inequality (see Ref. [25], p. 271), we have the estimate

$$\|\hat{\mathcal{C}}_\varepsilon^{(d)} u - \hat{\mathcal{C}}^{(d)} u\|_{L^2(\Omega;\mathbb{C})} \leq \|u\|_{L^2(\Omega;\mathbb{C})} \int \cdots \int_{|\boldsymbol{r}|<\varepsilon} |K^{(d)}(\boldsymbol{r}, \boldsymbol{0})| \,\mathrm{d}^d\boldsymbol{r} = O(\varepsilon^2)\|u\|_{L^2(\Omega;\mathbb{C})},$$

hence the uniform convergence $\lim_{\varepsilon\to0^+} \|\hat{\mathcal{C}}_\varepsilon^{(d)} - \hat{\mathcal{C}}^{(d)}\|_{L^2(\Omega;\mathbb{C})} = 0$. Being the uniform limit of a sequence of compact operators, the operator $\hat{\mathcal{C}}^{(d)} : L^2(\Omega; \mathbb{C}) \longrightarrow L^2(\Omega; \mathbb{C})$ must be compact. ∎

Remark By appealing to the Riesz-Schauder theory of compact operators, we see that for any fixed $\Omega \Subset \mathbb{R}^d, d = 1, 2, \dots$, the acoustic scattering problem $(\hat{I} - \chi\hat{\mathcal{C}}^{(d)})u = u_{\text{inc}}$ is well-posed except for a countable set of "mechanical resonance modes" $\{\chi_s, s = 1, 2, \dots | 1/\chi \in \sigma_p(\hat{\mathcal{C}}^{(d)})\}$. Furthermore, this exceptional set has no finite accumulation points. ∎

Proposition 2.13 (Electromagnetic Scattering in One-Dimensional Dielectrics) *For a bounded and non-empty open set* $\Omega \subset \mathbb{R}$, *the operator* $\hat{\mathcal{G}}^{(1)} : L^2(\Omega;\mathbb{C}^3) \longrightarrow L^2(\Omega;\mathbb{C}^3)$ *has an orthogonal decomposition* $\hat{\mathcal{G}}^{(1)} = (-\hat{I}) \oplus (\hat{\mathcal{C}}^{(1)} \oplus \hat{\mathcal{C}}^{(1)}) : L^2(\Omega;\mathbb{C}) \oplus L^2(\Omega;\mathbb{C}^2) \longrightarrow L^2(\Omega;\mathbb{C}) \oplus L^2(\Omega;\mathbb{C}^2)$, *with spectrum* $\sigma(\hat{\mathcal{G}}^{(1)}) = \{-1\} \cup \sigma(\hat{\mathcal{C}}^{(1)})$.

Proof Without loss of generality, we may assume that $\Omega \subset \mathbb{R}$ sits along the x-axis, and decompose the polarizations of the electric fields into perpendicular (i.e. directed along the x-axis) and parallel (i.e. directed along the y, z-axes) components as $\boldsymbol{E} = \boldsymbol{E}_\perp + \boldsymbol{E}_\|$ and $\boldsymbol{E}_{\mathrm{inc}} = \boldsymbol{E}_{\mathrm{inc},\perp} + \boldsymbol{E}_{\mathrm{inc},\|}$. Here, the nomenclature for "perpendicular" and "parallel" refers to their respective relation to the boundary surface $\partial\Omega \times \mathbb{R}^2$ of the "dielectric slabs" $\Omega \times \mathbb{R}^2 \subset \mathbb{R}^3$. Under these assumptions, the electromagnetic scattering problem (Eq. 2.63) decouples into the following two equations for $\boldsymbol{r} \in \Omega$ within the dielectric volume:

$$\begin{aligned} \boldsymbol{E}_{\mathrm{inc},\perp}(\boldsymbol{r}) &= (1+\chi)\boldsymbol{E}_\perp(\boldsymbol{r}), \\ \boldsymbol{E}_{\mathrm{inc},\|}(\boldsymbol{r}) &= \boldsymbol{E}_\|(\boldsymbol{r}) - \chi \int_{\Omega\subset\mathbb{R}} \boldsymbol{E}_\|(\boldsymbol{r}') \frac{k e^{-ik|\boldsymbol{r}-\boldsymbol{r}'|}}{2i}\,\mathrm{d}\boldsymbol{r}'. \end{aligned}$$

In other words, we have an algebraic equation for the perpendicular electric field $[\hat{I} - \chi(-\hat{I})]\boldsymbol{E}_\perp = \boldsymbol{E}_{\mathrm{inc},\perp}$, and an analog of the scalar wave scattering equation for the parallel electric field $(\hat{I} - \chi\hat{\mathcal{C}}^{(1)})\boldsymbol{E}_\| = \boldsymbol{E}_{\mathrm{inc},\|}$. This results in the said orthogonal decomposition $\hat{\mathcal{G}}^{(1)} = (-\hat{I}) \oplus (\hat{\mathcal{C}}^{(1)} \oplus \hat{\mathcal{C}}^{(1)}) : L^2(\Omega;\mathbb{C}) \oplus L^2(\Omega;\mathbb{C}^2) \longrightarrow L^2(\Omega;\mathbb{C}) \oplus L^2(\Omega;\mathbb{C}^2)$. The spectrum of $\hat{\mathcal{G}}^{(1)} : L^2(\Omega;\mathbb{C}^3) \longrightarrow L^2(\Omega;\mathbb{C}^3)$ can be consequently derived from the orthogonal decomposition as $\sigma(\hat{\mathcal{G}}^{(1)}) = \{-1\} \cup \sigma(\hat{\mathcal{C}}^{(1)})$, which is a countable set with the only possible accumulation point being $\{0\}$. ∎

Remark We note that the singular behavior of one-dimensional electromagnetic scattering at $\chi = -1$ persists even if we confine the vector wave scattering problem to transverse fields and require that Ω has finitely many connected components, as can be seen from the representations

$$\Phi(\Omega;\mathbb{C}^3) = \{\boldsymbol{E}_\perp \in L^2(\Omega;\mathbb{C}^3) | \partial\boldsymbol{E}_\perp(\boldsymbol{r})/\partial x = 0, \forall \boldsymbol{r} \in \Omega\} \oplus L^2(\Omega;\mathbb{C}^2), \tag{2.67}$$

and

$$\begin{aligned} \dim\ker^{\Phi}(\hat{I} + \hat{\mathcal{G}}^{(1)}) &= \dim\{\boldsymbol{E}_\perp \in L^2(\Omega;\mathbb{C}^3) | \partial\boldsymbol{E}_\perp(\boldsymbol{r})/\partial x = 0, \forall \boldsymbol{r} \in \Omega\} + 2\dim\ker(\hat{I} + \hat{\mathcal{C}}^{(1)}) \\ &= \dim\mathcal{H}^0(\Omega;\mathbb{R}) + 2\dim\ker(\hat{I} + \hat{\mathcal{C}}^{(1)}). \end{aligned}$$

In other words, the dimensionality of the eigenspace $\ker^{\Phi}(\hat{I} + \hat{\mathcal{G}}^{(1)})$ is related to the number of connected components $N_\Omega = \dim\mathcal{H}^0(\Omega;\mathbb{R})$ in $\Omega \subset \mathbb{R}$. This persistent singular behavior does not contradict the conclusions in Proposition 2.4 or 2.8, because $\mathbb{R} \setminus (\Omega \cup \partial\Omega)$ is disconnected (and so is the volume external to the "parallel slabs" $\mathbb{R}^3 \setminus [(\Omega \times \mathbb{R}^2) \cup \partial(\Omega \times \mathbb{R}^2)]$).

Unlike the electromagnetic scattering problems in higher dimensions, the one-dimensional electromagnetic scattering problem is special in the sense that its spectrum $\sigma(\hat{\mathcal{G}}^{(1)}) = \{-1\} \cup \sigma(\hat{\mathcal{C}}^{(1)}) = \sigma_p(\hat{\mathcal{G}}^{(1)}) \cup \{0\}$ does not differ significantly from the "mechanical resonance modes" in acoustic scattering $\sigma(\hat{\mathcal{C}}^{(1)})$ — the spectrum of a compact operator $\hat{\mathcal{C}}^{(1)}$. If we confine our attention to "physically admissible vector fields" (Eq. 2.67) defined on a bounded open set $\Omega \subset \mathbb{R}$ with finitely many connected components (so that Ω is a one-dimensional "smooth" submanifold of $\mathbb{R}$), then the operator $\hat{\mathcal{G}}^{(1)} = (-\hat{I}) \oplus (\hat{\mathcal{C}}^{(1)} \oplus \hat{\mathcal{C}}^{(1)}) : \Phi(\Omega;\mathbb{C}^3) \longrightarrow \Phi(\Omega;\mathbb{C}^3)$ is compact, because the identity operator defined on a finite-dimensional space is compact. For light scattering problems concerning higher-dimensional bounded open sets $\Omega \Subset \mathbb{R}^d$ $(d = 2, 3)$ with smooth boundary $\partial\Omega$, the compactness of $\hat{\mathcal{G}}^{(d)} : \Phi(\Omega;\mathbb{C}^3) \longrightarrow \Phi(\Omega;\mathbb{C}^3)$ will be lost. ∎

Proposition 2.14 (Electromagnetic Scattering in Two-Dimensional Dielectrics) *For a bounded and non-empty open set* $\Omega \subset \mathbb{R}^2$, *the linear mapping* $\hat{I} - \chi\hat{\mathcal{G}}^{(2)} : L^2(\Omega;\mathbb{C}^3) \longrightarrow L^2(\Omega;\mathbb{C}^3)$ *is a Fredholm operator with index zero for* $\chi \notin (-\infty, -1]$, *while the operator* $\hat{\mathcal{G}}^{(2)} : L^2(\Omega;\mathbb{C}^3) \longrightarrow L^2(\Omega;\mathbb{C}^3)$ *is not (necessarily) compact. When* Ω *has finitely many connected components* $\Omega_1, \dots, \Omega_{N_\Omega}, N_\Omega < +\infty$, *and the boundary* $\partial\Omega = \bigsqcup_{j=1}^{N_\Omega} \partial\Omega_j \subset \mathbb{R}^2$ *is smooth, the operator* $\hat{\mathcal{G}}^{(2)}(\hat{I} + 2\hat{\mathcal{G}}^{(2)}) : \Phi(\Omega;\mathbb{C}^3) \longrightarrow \Phi(\Omega;\mathbb{C}^3)$ *is compact, where* $\Phi(\Omega;\mathbb{C}^3)$ *is the Hilbert subspace of* $L^2(\Omega;\mathbb{C}^3)$ *generated by the totality of transverse fields.*

Proof In parallel to the argument in Proposition 1.1, we split the operator $\hat{\mathcal{G}}^{(2)} : L^2(\Omega;\mathbb{C}^3) \longrightarrow L^2(\Omega;\mathbb{C}^3)$ into the sum of two parts $\hat{\mathcal{G}}^{(2)} = (\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)}) + \hat{\gamma}^{(2)}$, where

$$((\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)})\boldsymbol{E})(\boldsymbol{r}) = \nabla\left[\nabla \cdot \iint_\Omega \frac{\boldsymbol{E}(\boldsymbol{r}')}{2\pi} \ln\frac{1}{|\boldsymbol{r}-\boldsymbol{r}'|}\,\mathrm{d}^2\boldsymbol{r}'\right]$$

satisfies $\sigma(\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)}) \subset [-1, 0]$ (cf. Lemma 1.1). One can use the Hilbert-Schmidt bound to confirm that the operator $\hat{\gamma}^{(2)} : L^2(\Omega; \mathbb{C}^3) \longrightarrow L^2(\Omega; \mathbb{C}^3)$ is compact. Therefore, when $\chi \in \mathbb{C} \setminus (-\infty, -1]$, the linear mapping $\hat{I} - \chi\hat{\mathcal{G}}^{(2)} : L^2(\Omega; \mathbb{C}^3) \longrightarrow L^2(\Omega; \mathbb{C}^3)$ is the sum of an invertible operator $\hat{I} - \chi(\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)})$ and a compact operator $-\chi\hat{\gamma}^{(2)}$, hence it is a Fredholm operator with index zero: $\operatorname{ind}(\hat{I} - \chi\hat{\mathcal{G}}^{(2)}) = 0, \forall \chi \in \mathbb{C} \setminus (-\infty, -1]$.

Meanwhile, we recall the following series expansion of the two-dimensional Green function (see Ref. [9], p. 127) in polar coordinates $\boldsymbol{r}(\rho, \phi)$ and $\boldsymbol{r}'(\rho', \phi')$:

$$\frac{1}{2\pi} \ln \frac{1}{|\boldsymbol{r} - \boldsymbol{r}'|} = \frac{1}{2\pi} \ln \frac{1}{\rho_>} + \frac{1}{2\pi} \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{\rho_<}{\rho_>} \right)^m \cos(m\phi - m\phi'), \quad \rho_> = \max(|\boldsymbol{r}|, |\boldsymbol{r}'|), \rho_< = \min(|\boldsymbol{r}|, |\boldsymbol{r}'|).$$

Applying this expansion to the circular shape $\Omega = O(\mathbf{0}, R) \cap \mathbb{R}^2 := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 < R^2\}$, we can check that $\boldsymbol{f}_m(\rho, \phi) := \nabla(\rho^{|m|} e^{im\phi}), m \in \mathbb{Z} \setminus \{0\} = \{\pm1, \pm2, \dots\}$ satisfies the eigenequation $(\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)})\boldsymbol{f}_m = -\boldsymbol{f}_m/2$. This shows that when $\Omega = O(\mathbf{0}, R)$ is circular, we have a non-zero eigenvalue $-1/2 \in \sigma_p(\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)})$ with infinite multiplicity. As a non-zero eigenvalue of a compact operator always has finite multiplicity, we see that the operator $\hat{\mathcal{G}}^{(2)} - \hat{\gamma}^{(2)} : L^2(\Omega; \mathbb{C}^3) \longrightarrow L^2(\Omega; \mathbb{C}^3)$ is not (necessarily) compact, and neither is $\hat{\mathcal{G}}^{(2)} : L^2(\Omega; \mathbb{C}^3) \longrightarrow L^2(\Omega; \mathbb{C}^3)$.

The proof of the compactness for $\hat{\mathcal{G}}^{(2)}(\hat{I} + 2\hat{\mathcal{G}}^{(2)}) : \Phi(\Omega; \mathbb{C}^3) \longrightarrow \Phi(\Omega; \mathbb{C}^3)$ follows essentially the same spirit as in Theorem 2.1, with according changes from the boundary surface ∂V to the the boundary surface $\partial\Omega \times \mathbb{R}$ generated by the boundary curve $\partial\Omega \subset \mathbb{R}^2$, an application of the Rellich-Kondrachov theorem to the compact imbedding $W^{1,p} \Subset L^2$ where $dp/(d - p) > 2$, $d = 2$, $1 < p < 2$, and adoption of the Calderón-Zygmund criteria (Eq. 1.27) for $d = 2$. ▌

Remark When $\Omega = O(\mathbf{0}, R) \cap \mathbb{R}^2$ takes the circular shape sitting in the xy-plane, the problem posed as $(\hat{I} - \chi\hat{\mathcal{G}}^{(2)})\boldsymbol{E} = \boldsymbol{E}_{\text{inc}} \in \Phi(\Omega; \mathbb{C}^3)$ is just the light scattering on a circular cylinder with "perpendicular incidence" (i.e. $\partial\boldsymbol{E}/\partial z = 0$, see Ref. [6], Chap. 15). This problem can be solved following a similar vein as the Mie scattering for a dielectric sphere. Instead of the spherical counterparts, now the solution will be expressed in terms of cylindrical Bessel functions $J_m, m = \pm1, \pm2, \dots$, cylindrical Hankel functions of the second kind $H_m := H_m^{(2)}, m = \pm1, \pm2, \dots$ along with their respective derivatives J'_m and H'_m (see Ref. [6], p. 301). When $\boldsymbol{E}$ is polarized along the z-axis ("‖"-mode), the "cylindrical Mie series" will have summands proportional to[36]

$$\frac{\nabla \times [\boldsymbol{r} J_m(\sqrt{1+\chi} k|\boldsymbol{r}|) e^{im\phi}]}{\sqrt{1+\chi} J'_m(\sqrt{1+\chi} kR) H_m(kR) - J_m(\sqrt{1+\chi} kR) H'_m(kR)}, \quad m = \pm1, \pm2, \dots, \boldsymbol{r} \in O(\mathbf{0}, R) \cap \mathbb{R}^2,$$

which encounter singularities at

$$\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)}) = \left\{ \lambda \neq 0, -1 \,\middle|\, \sqrt{1+\lambda^{-1}} J'_m(\sqrt{1+\lambda^{-1}} kR) H_m(kR) - J_m(\sqrt{1+\lambda^{-1}} kR) H'_m(kR) = 0, m = 1, 2, \dots \right\} \tag{2.68}$$

Here, the pair of transcendental equations for $\pm|m|$ are linearly dependent. When $\nabla \times \boldsymbol{E}$ is polarized along the z-axis ("$\perp$"-mode), the "cylindrical Mie series" will have summands proportional to

$$\frac{1}{\sqrt{1+\chi}} \frac{\nabla \times \nabla \times [\boldsymbol{r} J_m(\sqrt{1+\chi} k|\boldsymbol{r}|) e^{im\phi}]}{J'_m(\sqrt{1+\chi} kR) H_m(kR) - \sqrt{1+\chi} J_m(\sqrt{1+\chi} kR) H'_m(kR)}, \quad m = \pm1, \pm2, \dots, \boldsymbol{r} \in O(\mathbf{0}, R) \cap \mathbb{R}^2,$$

which encounter singularities at

$$\sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)}) = \left\{ \lambda \neq 0, -1 \,\middle|\, J'_m(\sqrt{1+\lambda^{-1}} kR) H_m(kR) - \sqrt{1+\lambda^{-1}} J_m(\sqrt{1+\lambda^{-1}} kR) H'_m(kR) = 0, m = 1, 2, \dots \right\} \tag{2.69}$$

Here, the rationale behind $-1 \notin \sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)}) \cup \sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)})$ is the Wroński determinant

$$\det \begin{pmatrix} \operatorname{Re} H_\nu^{(2)}(x) & \operatorname{Im} H_\nu^{(2)}(x) \\ \mathrm{d} \operatorname{Re} H_\nu^{(2)}(x) / \mathrm{d}\, x & \mathrm{d} \operatorname{Im} H_\nu^{(2)}(x) / \mathrm{d}\, x \end{pmatrix} = -\frac{2}{\pi x},$$

which implies $H_\nu^{(2)}(kR) \neq 0$ for $kR \neq 0, \nu \in \mathbb{C}$, and consequently

$$\lim_{\chi \to -1} \frac{\sqrt{1+\chi} J'_m(\sqrt{1+\chi} kR) H_m(kR) - J_m(\sqrt{1+\chi} kR) H'_m(kR)}{(\sqrt{1+\chi} kR)^m} = \frac{kR H_{m+1}^{(2)}(kR)}{2^m m!} \neq 0,$$

$$\lim_{\chi \to -1} \sqrt{1+\chi} \frac{J'_m(\sqrt{1+\chi} kR) H_m(kR) - \sqrt{1+\chi} J_m(\sqrt{1+\chi} kR) H'_m(kR)}{(\sqrt{1+\chi} kR)^m} = \frac{H_m^{(2)}(kR)}{2^m (m-1)!} \neq 0,$$

---

[36] In principle, there is a term related to $m = 0$, but $\nabla \times [\boldsymbol{r} J_0(\sqrt{1+\chi} k|\boldsymbol{r}|)] = \mathbf{0}$.

for positive integers m.

Despite the similarities of the transcendental equations involved in Eqs. 2.68 and 2.69, the two sets $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})$ and $\sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)})$ may have dramatically different properties. The set $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})$ occurring in Eq. 2.68 arises from an analog of the scalar wave scattering equation $(\hat{I}-\chi\hat{\mathcal{C}}^{(2)})(\boldsymbol{e}_z\cdot\boldsymbol{E})=\boldsymbol{e}_z\cdot\boldsymbol{E}_{\text{inc}}$, so we have $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})\subset\sigma(\hat{\mathcal{C}}^{(2)})$. Consequently, the only possible accumulation point in $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})$ is $\{0\}$. In contrast, the set $\sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)})$ occurring in Eq. 2.69 may have two accumulation points $\{0,-1/2\}$. Akin to Eq. 2.59, we may give a heuristic illustration of the dramatic difference between $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})$ and $\sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)})$ using two respective asymptotic expansions:

$$\begin{aligned}&\frac{\sqrt{1+\chi}J_m'(\sqrt{1+\chi}kR)H_m(kR)-J_m(\sqrt{1+\chi}kR)H_m'(kR)}{(\chi+1)^{m/2}}\\ ={}&\frac{i\left(\sqrt{\chi+1}+1\right)}{\pi kR}-\frac{i\left[(m-1)\chi\left(\sqrt{\chi+1}m+m+2\right)+2\sqrt{\chi+1}-2\right]kR}{4m\left(m^2-1\right)\pi}\left[1+O\left(\frac{k^2R^2}{m}\right)\right]; \qquad (2.70\text{a})\\ &\frac{J_m'(\sqrt{1+\chi}kR)H_m(kR)-\sqrt{1+\chi}J_m(\sqrt{1+\chi}kR)H_m'(kR)}{(\chi+1)^{(m-1)/2}}\\ ={}&\frac{i(\chi+2)}{\pi kR}-\frac{i\chi[m(\chi+2)-\chi]kR}{4\left(m^2-1\right)\pi}\left[1+O\left(\frac{k^2R^2}{m}\right)\right]. \qquad (2.70\text{b})\end{aligned}$$

As $m\to+\infty$, Eq. 2.70a tends to a non-zero number for whatever $\chi\in\mathbb{C}$, suggesting that $\sigma_p^{\Phi,\parallel}(\hat{\mathcal{G}}^{(2)})$ cannot have an accumulation point other than $\{0\}$. Meanwhile, if $\chi=-2$, Eq. 2.70b asymptotically vanishes, or roughly speaking, the left-hand side of Eq. 2.70b has a root that asymptotically behaves as $\chi_m\sim-2-(kR/m)^2$. This suggests that $\sigma_p^{\Phi,\perp}(\hat{\mathcal{G}}^{(2)})$ may have an accumulation point corresponding to $1/\chi=-1/2$. ∎

To summarize, we have the following table for the spectral properties of acoustic and electromagnetic scattering problems in dimensions $d=1,2$ and 3:

<table>
<tr><th>d</th><th>Acoustic Scattering
$(\hat{I}-\chi\hat{\mathcal{C}}^{(d)})u=u_{\text{inc}}$</th><th colspan="2">Electromagnetic Scattering
$(\hat{I}-\chi\hat{\mathcal{G}}^{(d)})\boldsymbol{E}=\boldsymbol{E}_{\text{inc}}$</th></tr>
<tr><td>1</td><td rowspan="3">$\hat{\mathcal{C}}^{(d)}:L^2(\Omega;\mathbb{C})\longrightarrow L^2(\Omega;\mathbb{C})$ is a compact operator, $\sigma(\hat{\mathcal{C}}^{(d)})=\sigma_p(\hat{\mathcal{C}}^{(d)})\cup\{0\}$ possibly accumulates at $\{0\}$</td><td>$\sigma(\hat{\mathcal{G}}^{(1)})=\{-1\}\cup\sigma(\hat{\mathcal{C}}^{(1)})=\sigma_p(\hat{\mathcal{G}}^{(1)})\cup\{0\}$, $\hat{\mathcal{G}}^{(1)}:\Phi(\Omega;\mathbb{C}^3)\longrightarrow\Phi(\Omega;\mathbb{C}^3)$ is compact</td><td rowspan="3">$\hat{\mathcal{G}}^{(d)}(\hat{I}+2\hat{\mathcal{G}}^{(d)}):\Phi(\Omega;\mathbb{C}^3)\longrightarrow\Phi(\Omega;\mathbb{C}^3)$ is a compact operator for $d=1,2,3$, $\sigma^{\Phi}(\hat{\mathcal{G}}^{(d)})\subset\sigma_p^{\Phi}(\hat{\mathcal{G}}^{(d)})\cup\{0,-1/2\}$ possibly accumulates at $\{0,-1/2\}$</td></tr>
<tr><td>2</td><td rowspan="2">$\hat{\mathcal{G}}^{(d)}:\Phi(\Omega;\mathbb{C}^3)\longrightarrow\Phi(\Omega;\mathbb{C}^3)$ $(d=2,3)$ is *not* (necessarily) a compact operator</td></tr>
<tr><td>3</td></tr>
</table>

When $\Phi(\Omega;\mathbb{C}^3)$ appears in the above table, the "smoothness condition" is tacitly assumed, i.e. a one-dimensional dielectric $\Omega\subset\mathbb{R}$ has to have finitely many connected components, and a high-dimensional dielectric $\Omega\subset\mathbb{R}^d$ $(d=2,3)$ has to have finitely many connected components $\Omega_j, 1\leq j\leq N_\Omega<+\infty$, and a smooth boundary $\partial\Omega=\bigsqcup_{j=1}^{N_\Omega}\partial\Omega_j$.

References

[1] John William Strutt (Lord Rayleigh). On the light from the sky, its polarization and colour. *Phil. Mag.*, 41:107–120, 1871.

[2] John William Strutt (Lord Rayleigh). On the electromagnetic theory of light. *Phil. Mag.*, 12:81–101, 1881.

[3] Richard Gans. Strahlungsdiagramme ultramikroskopischer Teilchen. *Ann. Phys.*, 76:29–38, 1925.

[4] Max Born. *Optik: ein Lehrbuch der elektromagnetischen Lichttheorie.* Verlag von Julius Springer, Berlin, Germany, 1933.

[5] Gustav Mie. Beiträge zur Optik trüber Medien, speziell kolloidaler Metallösungen. *Ann. Phys.*, 25:377–455, 1908.

[6] Hendrik Christoffel van de Hulst. *Light Scattering by Small Particles.* Dover Publications, Mineola, NY, 2nd edition, 1981.

[7] Alexander A. Kokhanovsky. *Optics of Light Scattering Media: Problems and Solutions.* Springer-Praxis, Chicester, UK, 2nd edition, 2001.

[8] Claus Müller. *Foundations of the Mathematical Theory of Electromagnetic Waves*, volume 155 of *Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, Berlin, Germany, 1969. (in cooperation with T. P. Higgins, revised and enlarged translation of the German original: *Grundprobleme der mathematischen Theorie elektromagnetischer Schwingungen*, Springer-Verlag, Berlin, Germany, 1957).

[9] John David Jackson. *Classical Electrodynamics.* John Wiley & Sons, New York, NY, 3rd edition, 1999.

[10] Max Born and Emil Wolf. *Principles of Optics.* Cambridge University Press, Cambridge, UK, 7th edition, 1999.

[11] Bruce J. Berne and Robert Pecora. *Dynamic Light Scattering: With Applications to Chemistry, Biology, and Physics.* Dover Publications, Mineola, NY, 1976.

[12] Sergeĭ L'vovich Sobolev. *Partial Differential Equations of Mathematical Physics.* Dover Publications, Mineola, NY, 1989. (translated by E. R. Dawson from the 3rd Russian edition of *Uravneniia͡ matematicheskoĭ fiziki* = Уравнения математической физики, GITTL, Moscow, USSR, 1954; English translation edited by T. A. A. Broadbent).

[13] Michael Reed and Barry Simon. *Scattering Theory*, volume III of *Methods of Modern Mathematical Physics.* Academic Press, New York, NY, 1979.

[14] David L. Colton and Rainer Kress. *Inverse Acoustic and Electromagnetic Scattering Theory*, volume 93 of *Applied Mathematical Sciences.* Springer, Berlin, Germany, 2nd edition, 1998.

[15] Neil V. Budko and Alexander B. Samokhin. Spectrum of the volume integral operator of electromagnetic scattering. *SIAM J. Sci. Comput.*, 28:682–700, 2006.

[16] Jussi Rahola. On the eigenvalues of the volume integral operator of electromagnetic scattering. *SIAM J. Sci. Comput.*, 21:1740–1754, 2000.

[17] William K. Saunders. On solutions of Maxwell's equations in an exterior region. *Proc. Natl. Acad. Sci. USA*, 38:342–348, 1952.

[18] David L. Colton and Rainer Kress. *Integral Equation Methods in Scattering Theory.* John Wiley & Sons, New York, NY, 1983.

[19] George C. Hsiao and Ralph E. Kleinman. Mathematical foundations for error estimation in numerical solutions of integral equations in electromagnetics. *IEEE Trans. Antennas Propag.*, 45:316–328, 1997.

[20] P. Scott Carney, John C. Schotland, and Emil Wolf. Generalized optical theorem for reflection, transmission, and extinction of power for scalar fields. *Phys. Rev. E.*, 70:036611, 2004.

[21] Dalcio K. Dacol and Dilip G. Roy. Generalized optical theorem for scattering in inhomogeneous media. *Phys. Rev. E.*, 72:036609, 2005.

[22] David Halliday and Andrew Curtis. Generalized optical theorem for surface waves and layered media. *Phys. Rev. E.*, 79:056603, 2009.

[23] John L. Kelley. *General Topology*, volume 27 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, NY, 1975.

[24] James Munkres. *Topology.* Prentice Hall, Upper Saddle River, NJ, 2nd edition, 2000.

[25] Elias M. Stein. *Singular Integrals and Differentiability Properties of Functions*, volume 30 of *Princeton Mathematical Series.* Princeton University Press, Princeton, NJ, 1970.

[26] David Gilbarg and Neil S. Trudinger. *Elliptic Partial Differential Equations of Second Order*, volume 224 of *Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, Berlin, Germany, 1983.

[27] Michael Eugene Taylor. *Partial Differential Equations I: Basic Theory*, volume 115 of *Applied Mathematical Sciences.* Springer-Verlag, New York, NY, 1996.

[28] Lawrence C. Evans. *Partial Differential Equations*, volume 19 of *Graduate Studies in Mathematics.* American Mathematical Society, Providence, RI, 1998.

[29] Elliot H. Lieb and Michael Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics.* American Mathematical Society, Providence, RI, 2nd edition, 2001.

[30] Jürgen Jost. *Partial Differential Equations*, volume 214 of *Graduate Texts in Mathematics.* Springer, New York, NY, 2nd edition, 2007.

[31] Paul R. Halmos. *Measure Theory*, volume 18 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, NY, 1974.

[32] K. S. Shifrin. *Scattering of Light in a Turbid Medium.* NASA Tech. Transl. NASA TT F-477, Washington, D.C., 1968. (translated from the Russian original: *Rassei͡anie sveta v mutnoĭ srede* = Рассеяние света в мутной среде, Gostekhteorizdat, Moscow, USSR, 1951).

[33] J. van Bladel. Some remarks on Green's dyadic for infinite space. *IRE Trans. Antennas Propag.*, 9:563–566, 1961.

[34] A. R. Holt, N. K. Uzunoglu, and B. G. Evans. An integral equation solution to the scattering of electromagnetic radiation by dielectric spheroids and ellipsoids. *IEEE Trans. Antennas Propag.*, 26:706–712, 1978.

[35] George H. Goedecke and Sean G. O'Brien. Scattering by irregular inhomogeneous particles via the digitized Green's function algorithm. *Appl. Opt.*, 27:2431–2438, 1988.

[36] Michael I. Mishchenko, Warren J. Wiscombe, Joop W. Hovenier, and Larry D. Travis. Overview of scattering by nonspherical particles. In Michael I. Mishchenko, Joachim W. Hovenier, and Larry D. Travis, editors, *Light Scattering by Nonspherical Particles*, chapter 2, pages 29–60. Academic Press, San Diego, CA, 2000.

[37] M. A. Yurkin and A. G. Hoekstra. The discrete dipole approximation: An overview and recent developments. *J. Quant. Spect. & Rad. Trans.*, 106:558–589, 2007.

[38] Loukas Grafakos. *Classical Fourier Analysis*, volume 249 of *Graduate Texts in Mathematics.* Springer, New York, NY, 2nd edition, 2008.

[39] Reinhard Börger. A non-Jordan-measurable regularly open subset of the unit interval. *Archiv der Mathematik*, 73:262–264, 1999.

[40] Robert D. Richtmyer. *Principles of Advanced Mathematical Physics*, volume I. Springer-Verlag, New York, NY, 1978.

[41] Tosio Kato. *Perturbation Theory for Linear Operators*, volume 132 of *Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, Berlin, Germany, 2nd edition, 1980.

[42] John B. Conway. *A Course in Functional Analysis*, volume 96 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, NY, 1985.

[43] F. Riesz. Über lineare Funktionalgleichungen. *Acta Math.*, 41:71–98, 1916.

[44] Kôsaku Yosida. *Functional Analysis*, volume 123 of *Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, Berlin, Germany, 6th edition, 1980.

[45] Michael Reed and Barry Simon. *Functional Analysis*, volume I of *Methods of Modern Mathematical Physics.* Academic Press, New York, NY, 1972.

[46] Yuli Eidelman, Vitali Milman, and Antonis Tsolomitis. *Functional Analysis: An Introduction*, volume 66 of *Graduate Studies in Mathematics.* American Mathematical Society, Providence, RI, 2004.

[47] Max Born. Quantenmechanik der Stoßvorgänge. *Z. Phys.*, 38:803–827, 1926.

[48] A. P. Calderón and A. Zygmund. On the existence of certain singular integrals. *Acta Math.*, 88:85–139, 1952.

[49] Elias M. Stein. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, volume III of *Monographs in Harmonic Analysis.* Princeton University Press, Princeton, NJ, 1993.

[50] G. H. Hardy and J. E. Littlewood. Some properties of fractional integrals (1). *Math. Z.*, 27:565–606, 1928.

[51] G. H. Hardy and J. E. Littlewood. On certain inequalities connected with the calculus of variations. *J. London Math. Soc.*, 5:34–39, 1930.

[52] S. L. Sobolev. On a theorem of functional analysis. *Mat. Sb. (N.S.)*, 4:471–497, 1938. (English translation in Amer. Math. Soc. Transl. Ser. 2, 34:39-68, 1963).

[53] E. H. Lieb. Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities. *Ann. of Math.*, 118:349–374, 1983.

[54] Milton Abramowitz and Irene A. Stegun, editors. *Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables.* Dover Publications, New York, NY, 1965.

[55] George Neville Watson. *A Treatise on the Theory of Bessel Functions.* Cambridge University Press, Cambridge, UK, 2nd edition, 1952.

[56] Peter D. Lax and Ralph S. Phillips. *Scattering Theory*, volume 26 of *Pure and Applied Mathematics.* Academic Press, San Diego, CA, revised edition, 1989.

[57] Michael Eugene Taylor. *Partial Differential Equations II: Qualitative Studies of Linear Equations*, volume 116 of *Applied Mathematical Sciences.* Springer-Verlag, New York, NY, 1996.

[58] Einar Hille. *Functional Analysis and Semigroups*, volume 31 of *American Mathematical Society Colloquium Publications.* American Mathematical Society, New York, NY, 1948.

[59] Kôsaku Yosida. On the differentiability and the representation of one-parameter semigroups of linear operators. *J. Math. Soc. Japan*, 1:15–21, 1948.

[60] Klaus-Jochen Engel and Rainer Nagel. *One-Parameter Semigroups for Linear Evolution Equations*, volume 194 of *Graduate Texts in Mathematics.* Springer, New York, NY, 2000. (with contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli, and R. Schnaubelt).

[61] S. Bochner. Integration von Funktionen, deren Werte die Elemente eines Vektorraumes sind. *Fund. Math.*, 20:262–276, 1933.

[62] N. Dunford. Integration in general analysis. *Trans. Amer. Math. Soc.*, 37:441–453, 1935.

[63] Einar Hille. Representation of one-parameter semi-groups of linear transformations. *Proc. Nat. Acad. Sci. USA*, 28:175–178, 1942.

[64] G. Lumer and R. S. Phillips. Dissipative operators in a Banach space. *Pacific J. Math.*, 11:679–698, 1961.

[65] Robert Dautray and Jacques-Louis Lions. *Physical Origins and Classical Methods*, volume 1 of *Mathematical Analysis and Numerical Methods for Science and Technology.* Springer-Verlag, Berlin, Germany, 1990. (with the collaboration of Philippe Bénilan, Michel Cessenat, André Gervat, Alain Kavenoky, Hélène Lanchon, translated by Ian N. Sneddon from the French original: *Analyse mathématique et calcul numérique pour les sciences et les techniques*, S. A. Masson, Paris, France, 1984, 1985).

[66] Robert A. Adams and John J. F. Fournier. *Sobolev Spaces.* Academic Press, Amsterdam, the Netherlands, 2nd edition, 2003.

[67] Robert Dautray and Jacques-Louis Lions. *Spectral Theory and Applications*, volume 3 of *Mathematical Analysis and Numerical Methods for Science and Technology.* Springer-Verlag, Berlin, Germany, 1990. (with the collaboration of Michel Artola and Michel Cessenat, transl. by John C. Amson from the French original: *Analyse mathématique et calcul numérique pour les sciences et les techniques*, S. A. Masson, Paris, France, 1984, 1985).

[68] Jacques-Louis Lions and Enrico Magenes. *Non-Homogeneous Boundary Value Problems and Applications I*, volume 181 of *Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, Berlin, Germany, 1972. (translated by P. Kenneth from the French original: *Problèmes aux limites non homogènes et applications (tome I)*, S. A. Dunod, Paris, France, 1968).

[69] Robert Dautray and Jacques-Louis Lions. *Functional and Variational Methods*, volume 2 of *Mathematical Analysis and Numerical Methods for Science and Technology.* Springer-Verlag, Berlin, Germany, 1988. (with the collaboration of Michel Artola, Marc Authier, Philippe Bénilan, Michel Cessenat, Jean-Michel Combes, Hélène Lanchon, Bertrand Mercier, Claude Wild, Claude Zuily, translated by Ian N. Sneddon from the French original: *Analyse mathématique et calcul numérique pour les sciences et les techniques*, S. A. Masson, Paris, France, 1984, 1985).

[70] G. A. Bliss. An integral inequality. *J. London Math. Soc.*, 5:40–46, 1930.

[71] E. Rodemich. The Sobolev inequalities with best possible constants. In *Analysis Seminar at California Institute of Technology*, 1966.

[72] G. Talenti. Best constant in Sobolev inequality. *Ann. Mat. Pura Appl.*, 110:353–372, 1976.

[73] Neil V. Budko and Alexander B. Samokhin. Singular modes of the electromagnetic field. *J. Phys. A: Math. Theor.*, 40:6239–6250, 2007.

[74] Sheldon Axler, Paul Bourdon, and Wade Ramey. *Harmonic Function Theory*, volume 137 of *Graduate Texts in Mathematics.* Springer, New York, NY, 2nd edition, 2001.

[75] Stefan Bergman. *The Kernel Function and Conformal Mapping*, volume 5 of *Mathematical Surveys and Monographs.* American Mathematical Society, Providence, RI, 2nd edition, 1970.

[76] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory.* Addison-Wesley, Reading, MA, 1995.

[77] D. P. Craig and T. Thirunamachandran. *Molecular Quantum Electrodynamics: An Introduction to Radiation Molecule Interactions.* Dover Publications, Mineola, NY, 1998.

[78] Neil V. Budko and Alexander B. Samokhin. Classification of electromagnetic resonances in finite inhomogeneous three-dimensional structures. *Phys. Rev. Lett.*, 96:023904, 2006.

[79] Wolfgang Arendt and Charles J. K. Batty. Tauberian theorems for one-parameter semigroups. *Trans. Amer. Math. Soc.*, 306:837–852, 1988.

[80] Yu. I. Lyubich and Vũ Quốc Phóng. Asymptotic stability of linear differential equations on Banach spaces. *Studia Math.*, 88:37–42, 1988.

[81] Manfredo Perdigão do Carmo. *Differential Geometry of Curves and Surfaces.* Prentice-Hall, Englewood Cliffs, NJ, 1976.

[82] Hans Samelson. The orientability of hypersurfaces in $\mathbb{R}^n$. *Proc. Amer. Math. Soc.*, 22:301–302, 1969.

[83] P. Alexandroff and H. Hopf. *Topologie.* Springer, Berlin, Germany, 1935.

[84] V. Balakrishnan. Fractional powers of closed operators and the semi-groups generated by them. *Pacific J. Math.*, 10:419–437, 1960.

[85] Morris W. Hirsch. *Differential Topology*, volume 33 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, NY, 1976.

[86] A. S. Besicovitch. On the definition and value of the area. *Quarterly Journal of Mathematics*, 16:86–102, 1945.

[87] Raoul Bott and Loring W. Tu. *Differential Forms in Algebraic Topology*, volume 82 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, NY, 1982.

[88] Ib Henning Madsen and Jørgen Tornehave. *From Calculus to Cohomology.* Cambridge University Press, Cambridge, UK, 1997.

[89] Lars V. Ahlfors. *Complex Analysis.* McGraw-Hill, New York, NY, 3rd edition, 1979.

[90] R. E. Kleinman, G. F. Roach, and P. M. van den Berg. Convergent Born series for large refractive indices. *J. Opt. Soc. Am. A*, 7:890–897, 1990.

[91] T. W. Chen. Simple formula for light scattering by a large spherical dielectric. *Appl. Opt.*, 32:7568–7571, 1993.